

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 38/05, 31/675, 31/67, 31/66, 31/665, C07D 209/04, 213/16, 271/07, 307/36, 33/10, C07F 9/28	A1	(11) International Publication Number: WO 97/48409 (43) International Publication Date: 24 December 1997 (24.12.97)
(21) International Application Number: PCT/US97/10149 (22) International Filing Date: 13 June 1997 (13.06.97) (30) Priority Data: 08/665,775 17 June 1996 (17.06.96) US 08/864,545 28 May 1997 (28.05.97) US (71) Applicant: GUILFORD PHARMACEUTICALS INC. [US/US]; 6611 Tributary Street, Baltimore, MD 21224 (US). (72) Inventors: SLUSHER, Barbara, S.; Guilford Pharmaceuticals Inc., 6611 Tributary Street, Baltimore, MD 21224 (US). JACKSON, Paul, F.; Guilford Pharmaceuticals Inc., 6611 Tributary Street, Baltimore, MD 21224 (US). TAYS, Kevin, L.; Guilford Pharmaceuticals Inc., 6611 Tributary Street, Baltimore, MD 21224 (US). MACLIN, Keith, M.; Guilford Pharmaceuticals Inc., 6611 Tributary Street, Baltimore, MD 21224 (US). (74) Agent: NATH, Gary, M.; Nath & Associates, Suite 750, 1835 K Street, N.W., Washington, DC 20006-1203 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: METHODS OF CANCER TREATMENT USING NAALADASE INHIBITORS (57) Abstract The present disclosure relates to dipeptidase inhibitors, and more particularly, to novel methods of using phosphonate derivatives, hydroxyphosphinyl derivatives, and phosphoramidate derivatives to inhibit N-Acetylated α -Linked Acidic Dipeptidase (NAALADase) enzyme activity, and to treat prostate diseases, especially using the compounds of the present invention for the inhibition of the growth of prostate cancer cells.		

BEST AVAILABLE COPY

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

METHODS OF CANCER TREATMENT USING NAALADASE INHIBITORS

RELATED APPLICATIONS

This application is a continuation-in-part (CIP) of U.S. patent application serial number 08/665,775, filed June 17, 1996, entitled "Methods of Cancer Treatment Using NAALADase Inhibitors".

BACKGROUND OF THE INVENTION1. Field of the Invention

The present invention relates to novel methods of treating cancer, preventing tumor cell growth, inhibiting prostate tumor cell growth, and inhibiting NAALADase enzyme activity, by administering an effective amount of a NAALADase inhibitor.

2. Description of the Prior ArtProstate Cancer

Prostate cancer is the leading form of cancer and the second leading cause of death from cancer for men in the United States. The American Cancer Society has estimated that in 1996 alone, 317,100 new cases of prostate cancer were diagnosed and 41,400 deaths were caused by prostate cancer. The incidence rate of prostate cancer increased 65% between 1980 and 1990, and will continue to rise with improved screening tests and longer life expectancies. While most men used to die of other illnesses before prostate cancer had a chance to develop, higher prostate cancer mortality rates are expected as men live longer and the disease has more time to progress.

In 1993, the molecular cloning of Prostate Specific Membrane Antigen (PSMA) was reported as a potential prostate carcinoma marker and hypothesized to serve as a target for imaging and cytotoxic treatment modalities for prostate cancer. PSMA antibodies, particularly indium-111 labelled and tritium labelled PSMA antibodies, have been described and examined clinically for the diagnosis and treatment of prostate cancer. PSMA is expressed in prostatic ductal epithelium and is present in seminal plasma, prostatic fluid and urine. In 1996, it was found that the expression of PSMA cDNA confers the activity of NAALADase.

NAALADase Inhibitors

NAAG and NAALADase have been implicated in several human and animal pathological conditions relating to glutamate abnormalities and neurotoxicity. For example, it has been demonstrated that intra-hippocampal injections of NAAG elicit prolonged seizure activity. More recently, it was reported that rats genetically prone to epileptic seizures have a persistent increase in their basal level of NAALADase activity. These observations lend support the hypothesis that increased availability of synaptic glutamate elevates seizure susceptibility, and suggest that NAALADase inhibitors may provide anti-epileptic activity.

NAAG and NAALADase have also been implicated in the pathogenesis of ALS and in the pathologically similar animal disease called Hereditary Canine Spinal Muscular Atrophy (HCSMA). It has been shown that concentrations of NAAG and its metabolites -- NAA, glutamate and aspartate -- are

elevated two- to three-fold in the cerebrospinal fluid of ALS patients and HCSMA dogs. Additionally, NAALADase activity is significantly increased (two- to three-fold) in post-mortem spinal cord tissue from ALS patients and HCSMA dogs. As such, NAALADase inhibitors might be clinically useful in curbing the progression of ALS if an increased metabolism of NAAG is responsible for the alterations of CSF levels of these acidic amino acids and peptides.

Abnormalities in NAAG levels and NAALADase activity have also been documented in post-mortem schizophrenic brain, specifically in the prefrontal and limbic brain regions.

The findings described above suggest that NAALADase inhibitors could be useful in treating glutamate abnormalities. However, the present invention is directed to the surprising and unexpected discovery that the novel compounds of the present invention are not only effective NAALADase inhibitors but are effective in treating prostate diseases, particularly prostate cancer. Although the cancer data relate to prostate cancer cells, NAALADase inhibitors are expected to be equally effective in treating cancer of other tissues where NAALADase enzyme reside, such as the brain, kidney and testis.


While a few NAALADase inhibitors have been identified, they have only been used in non-clinical research. Examples of such inhibitors include general metallopeptidase inhibitors such as o-phenanthroline, metal chelators such as EGTA and EDTA, and peptide analogs such as quisqualic acid and β -NAAG. Accordingly, a need exists for more NAALADase inhibitors to be

identified and, particularly, for the treatment of prostate diseases such as prostate cancer.

SUMMARY OF THE INVENTION

5 The present invention is directed to novel methods for treating cancer, preventing tumor cell growth, inhibiting prostate tumor cell growth, and inhibiting NAALADase enzyme activity, in an animal, comprising administering an effective amount of a NAALADase inhibitor.

10 Preferred NAALADase inhibitors include compounds of formula I:



I

wherein

- 15 R_1 is hydrogen, C_1 - C_9 straight or branched chain alkyl, C_2 - C_9 straight or branched chain alkenyl group, C_3 - C_9 cycloalkyl, C_5 - C_9 cycloalkenyl, or Ar_1 ;
- X is CH_2 , O , or N ; and
- 20 R_2 is C_1 - C_9 straight or branched chain alkyl, C_2 - C_9 straight or branched chain alkenyl group, C_3 - C_9 cycloalkyl, C_5 - C_9 cycloalkenyl, or Ar_1 , wherein said alkyl, alkenyl, cycloalkyl, cycloalkenyl or aryl groups may be optionally substituted with carboxylic acid.

Preferably, the compound of formula I is present in an amount that is effective for inhibiting NAALADase enzyme activity, or treating a prostate disease in an animal.

The present invention further relates to a method of inhibiting NAALADase enzyme activity in an animal, comprising administering an effective amount of the compound of formula I to said animal.

BRIEF DESCRIPTION OF THE FIGURES

FIG. 1 is a bar graph plotting the growth of the prostate cancer cell line, LNCAP, against various concentrations of quisqualic acid, a NAALADase Inhibitor. FIG. 1 shows the effect of 7-day treatment with quisqualate on the growth of LNCAP cells. Concentrations ranging from 10 nM to 1 μ M of quisqualate show a sharp dose-dependent decrease of LNCAP cell proliferation as indicated by the significant decrease in the incorporation of [3H]thymidine.

FIG. 2 is a bar graph plotting the growth of the prostate cancer cell line, LNCAP, against various concentrations of 2-(phosphonomethyl)pentanedioic acid, a NAALADase Inhibitor. FIG. 2 shows the effect of 7-day treatment with 2-(phosphonomethyl)pentanedioic acid on the growth of LNCAP cells. Concentrations ranging from 100 pM to 10 nM of 2-(phosphonomethyl)pentanedioic acid show a sharp dose-dependent decrease of LNCAP cell proliferation as indicated by the significant decrease in the incorporation of [3H]thymidine.

FIG. 3 is a line graph of the response of LNCAP human

prostate tumors to daily treatment with 2-(phosphonomethyl)pentanedioic acid. Mean of individual tumor volumes are plotted as a function of time after the start of treatment. Error bars represent the SEM. Treatment with 2-(phosphonomethyl)pentanedioic acid for six weeks resulted in statistically significant difference between both the control group and animals given daily injections of drug ($p=0.04$), and the control group and animals implanted with polymer ($p=0.02$).

FIG. 4 is a line graph plotting the survival percentage of animals treated with injections against the number of days. FIG. 4 shows the higher mean survival percentage of animals injected with 2-(phosphonomethyl)pentanedioic acid mixed with polymer as compared to those animals only receiving intratumoral injections of 2-(phosphonomethyl)pentanedioic acid or a vehicle control. The graph shows that 88% of the animals treated with polymer were alive after 72 days, and those animals had small tumors.

FIG. 5 is a line graph plotting tumor growth against days following ratunning cell injections. Cells were injected, over a period of 84 days, with various dosages of 2-(phosphonomethyl)pentanedioic acid and a control vehicle. FIG. 5 shows that tumor growth slowed as a function of 2-(phosphonomethyl)pentanedioic acid dosage.

FIG. 6 is a line graph of the response of R3327 rat prostate tumors to daily treatment with 2-[[[(phenylmethyl)hydroxyphosphinyl)methyl]pentanedioic acid. Mean of individual tumor volumes expressed relative to the volume at the start of treatment (V/V_0) are plotted as a

function of time. Treatment with 2-

5 [((phenylmethyl)hydroxyphosphinyl)methyl]pentanedioic acid for 2.5 weeks resulted in a statistically significant difference between the control group and animals given daily injections of 1 μ g of drug ($p=0.02$).

DETAILED DESCRIPTION OF THE INVENTION

Definitions

10 "Compound 3" refers to 2-(phosphonomethyl)pentanedioic acid, a NAALADase inhibitor.

"Inhibition", in the context of enzymes, refers to reversible enzyme inhibition such as competitive, uncompetitive and non-competitive inhibition. Competitive, uncompetitive and non-competitive inhibition can be
15 distinguished by the effects of an inhibitor on the reaction kinetics of an enzyme. Competitive inhibition occurs when the inhibitor combines reversibly with the enzyme in such a way that it competes with a normal substrate for binding at the active site. The affinity between the inhibitor and the
20 enzyme may be measured by the inhibitor constant, K_i , which is defined as:

$$K_i = \frac{[E][I]}{[EI]}$$

25

wherein [E] is the concentration of the enzyme, [I] is the concentration of the inhibitor, and [EI] is the concentration of the enzyme-inhibitor complex formed by the reaction of the enzyme with the inhibitor. Unless otherwise specified, K_i as

used herein refers to the affinity between the inventive compounds and NAALADase. "IC50" is a related term used to define the concentration or amount of a compound which is required to cause a 50% inhibition of the target enzyme.


5 The term "inhibition", in the context of tumor growth or tumor cell growth, may be assessed by delayed appearance of primary or secondary tumors, slowed development of primary or secondary tumors, decreased occurrence of primary or secondary tumors, slowed or decreased severity of secondary effects of
10' disease, arrested tumor growth and regression of tumors, among others. In the extreme, complete inhibition, is referred to herein as prevention.

 "NAAG" refers to N-acetyl-aspartyl-glutamate, an important peptide component of the brain, with levels
15 comparable to the major inhibitor neurotransmitter gamma-aminobutyric acid (GABA). NAAG is neuron-specific, present in synaptic vesicles and released upon neuronal stimulation in several systems presumed to be glutamatergic. Studies suggest that NAAG may function as a neurotransmitter and/or
20 neuromodulator in the central nervous system, or as a precursor of the neurotransmitter glutamate.

 "NAALADase" refers to N-acetylated α -linked acidic dipeptidase, a membrane-bound metallopeptidase which catabolizes NAAG to N-acetylaspartate (NAA) and glutamate:

25

Catabolism of NAAG by NAALADase



5 NAALADase shows a high affinity for NAAG with a K_m of 540 nM. If NAAG is a bioactive peptide, then NAALADase may serve to inactivate NAAG'S synaptic action. Alternatively, if NAAG functions as a precursor for glutamate, the primary function of NAALADase may be to regulate synaptic glutamate

10 availability.

 "Pharmaceutically acceptable salt" refers to a salt of the inventive compounds which possesses the desired pharmacological activity and which is neither biologically nor otherwise undesirable. The salt can be formed with inorganic

15 acids such as acetate, adipate, alginate, aspartate, benzoate, benzenesulfonate, bisulfate butyrate, citrate, camphorate, camphorsulfonate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, fumarate, glucoheptanoate, glycerophosphate, hemisulfate heptanoate, hexanoate, hydro-

20 chloride hydrobromide, hydroiodide, 2-hydroxyethanesulfonate, lactate, maleate, methanesulfonate, 2-naphthalenesulfonate, nicotinate, oxalate, thiocyanate, tosylate and undecanoate. Examples of a base salt include ammonium salts, alkali metal

salts such as sodium and potassium salts, alkaline earth metal salts such as calcium and magnesium salts, salts with organic bases such as dicyclohexylamine salts, N-methyl-D-glucamine, and salts with amino acids such as arginine and lysine. The
5 basic nitrogen-containing groups can be quarternized with agents including lower alkyl halides such as methyl, ethyl, propyl and butyl chlorides, bromides and iodides; dialkyl sulfates such as dimethyl, diethyl, dibutyl and diamyl sulfates; long chain halides such as decyl, lauryl, myristyl
10 and stearyl chlorides, bromides and iodides; and aralkyl halides such as benzyl and phenethyl bromides.

The term "prevention", in relation to tumor growth or tumor cell growth, means no tumor or tumor cell growth if none had occurred, no further tumor or tumor cell growth if there
15 had already been growth.

The term "prostate disease" relates to prostate cancer such as adenocarcinoma or metastatic cancers, conditions characterized by abnormal growth of prostatic epithelial cells such as benign prostatic hyperplasia, and other conditions
20 requiring treatment by the compounds of the present invention.


"PSA" refers to Prostate Specific Antigen, a well known prostate cancer marker. It is a protein produced by prostate cells and is frequently present at elevated levels in the blood of men with prostate cancer. PSA correlates with tumor
25 burden, serves as an indicator of metastatic involvement, and provides a parameter for following a prostate cancer patient's response to surgery, irradiation and androgen replacement therapy.

"PSMA" refers to Prostate Specific Membrane Antigen, a potential prostate carcinoma marker that has been hypothesized to serve as a target for imaging and cytotoxic treatment modalities for prostate cancer. PSMA is expressed in
5 prostatic ductal epithelium and is present in seminal plasma, prostatic fluid and urine. It has been found that the expression of PSMA cDNA confers the activity of NAALADase.

The term "treatment" refers to any process, action, application, therapy, or the like, wherein an animal,
10 including a human being, is subject to medical aid with the object of improving the animal's condition, directly or indirectly.

Preferred NAALADase Inhibitors of the Present Invention

15 The present invention relates to a compound of formula I:



I


or a pharmaceutically acceptable salt, hydrate, or a mixture thereof, wherein:

- 20 R_1 is hydrogen, C_1 - C_8 straight or branched chain alkyl, C_2 - C_8 straight or branched chain alkenyl group, C_1 - C_8 cycloalkyl, C_5 - C_8 cycloalkenyl, or Ar_1 ;
 X is CH_2 , O , or NR_1 , where R_1 is defined above; and
 R_2 is C_1 - C_8 straight or branched chain alkyl, C_2 - C_8
25 straight or branched chain alkenyl group, C_1 - C_8

cycloalkyl, C₃-C₆ cycloalkenyl, or Ar₁, wherein said alkyl, alkenyl, cycloalkyl, cycloalkenyl or aryl group may be optionally substituted with carboxylic acid.

5 The present invention also contemplates that said alkyl, alkenyl, cycloalkyl, cycloalkenyl or aryl groups may be optionally substituted with C₁-C₆ cycloalkyl, C₁ or C₂ cycloalkyl, C₃-C₆ cycloalkenyl, C₁-C₄ alkyl, C₁-C₄ alkenyl, halo, hydroxy, carboxy, nitro, trifluoromethyl, C₁-C₆ straight or
10 branched chain alkyl or alkenyl, C₁-C₄ alkoxy, C₁-C₄ alkenyloxy, phenoxy, benzyloxy, amino, or Ar₁, and where Ar₁ is selected from the group consisting of 1-naphthyl, 2-naphthyl, 2-indolyl, 3-indolyl, 4-indolyl, 2-furyl, 3-furyl, tetrahydrofuranyl, 2-thienyl, 3-thienyl, 4-thienyl, 2-, 3-, or
15 4-pyridyl, or phenyl, having one to five substituents which are independently selected from the group consisting of hydrogen, halo, hydroxy, carboxy, nitro, trifluoromethyl, C₁-C₆ straight or branched alkyl or alkenyl, C₁-C₄ alkoxy or C₁-C₄ alkenyloxy, phenoxy, benzyloxy, and amino; or pharmaceutically
20 acceptable salts, hydrates, or mixtures thereof.

In a preferred embodiment, the compound is selected from the group of formula II:




II

25 wherein

R₁ is hydrogen, C₁-C₆ straight or branched chain alkyl,

C₂-C₈, straight or branched chain alkenyl group, C₁-C₈, cycloalkyl, C₅-C₇, cycloalkenyl, or Ar₁; and
R₂ is C₁-C₈, straight or branched chain alkyl, C₂-C₈, straight or branched chain alkenyl group, C₁-C₈, cycloalkyl, C₅-C₇, cycloalkenyl, or Ar₁, wherein said alkyl, alkenyl, cycloalkyl, cycloalkenyl or aryl group may be optionally substituted with carboxylic acid.

In another preferred embodiment, the R groups are either straight or branched aliphatic substituents or carbocyclic substituents illustrated by the compounds selected from the group of formula II:



wherein

R₁ is hydrogen, C₁-C₈, straight or branched chain alkyl, C₂-C₈, straight or branched chain alkenyl group, C₁-C₈, cycloalkyl, C₅-C₇, cycloalkenyl, 1-naphthyl, 2-naphthyl, or phenyl; and
R₂ is C₁-C₈, straight or branched chain alkyl, C₂-C₈, straight or branched chain alkenyl group, C₁-C₈, cycloalkyl, C₅-C₇, cycloalkenyl, 1-naphthyl, 2-naphthyl, or phenyl, wherein said alkyl, alkenyl, cycloalkyl, cycloalkenyl, 1-naphthyl, 2-naphthyl, or

phenyl group may be optionally substituted with carboxylic acid.

Especially preferred methods utilize compounds wherein R₁ is either a straight or branched aliphatic group or a carbocyclic group, R₂ is ethyl which is substituted with a carboxylic acid, and X is CH₂, are selected from the group consisting of:

- 2-[[methylhydroxyphosphinyl]methyl]pentanedioic acid;
- 10 2-[[ethylhydroxyphosphinyl]methyl]pentanedioic acid;
- 2-[[propylhydroxyphosphinyl]methyl]pentanedioic acid;
- 2-[[butylhydroxyphosphinyl]methyl]pentanedioic acid;
- 2-[[cyclohexylhydroxyphosphinyl]methyl]pentanedioic acid;
- 2-[[cyclohexyl)methylhydroxyphosphinyl]methyl]pentanedioic
- 15 acid;
- 2-[[phenylhydroxyphosphinyl]methyl]pentanedioic acid;
- 2-[[benzylhydroxyphosphinyl]methyl]pentanedioic acid;
- 2-[[phenylethylhydroxyphosphinyl]methyl]pentanedioic acid;
- 2-[[phenylpropylhydroxyphosphinyl]methyl]pentanedioic acid;
- 20 2-[[phenylbutylhydroxyphosphinyl]methyl]pentanedioic acid;
- 2-[[4-methylbenzyl)hydroxyphosphinyl]methyl]pentanedioic acid;
- 2-[[4-fluorobenzyl)hydroxyphosphinyl]methyl]pentanedioic acid;
- 25 2-[[2-fluorobenzyl)hydroxyphosphinyl]methyl]pentanedioic acid;
- 2-[[pentafluorobenzyl)hydroxyphosphinyl]methyl]pentanedioic acid;

- 2-[[(methoxybenzyl)hydroxyphosphinyl]methyl]pentanedioic acid;
2-[[(4-fluorophenyl)hydroxyphosphinyl]methyl]pentanedioic
acid;
2-[[((hydroxy)phenylmethyl)hydroxyphosphinyl]methyl]
5 pentanedioic acid;
2-[[(3-methylbenzyl)hydroxyphosphinyl]methyl]pentanedioic
acid;
2-(phosphonomethyl)pentanedioic acid;
2-[[(3-
10 trifluoromethylbenzyl)hydroxyphosphinyl]methyl]pentanedioic
acid;
2-[[(2, 3, 4-trimethoxyphenyl)hydroxyphosphinyl]methyl]
pentanedioic acid;
2-[[(1-naphthyl)hydroxyphosphinyl]methyl]pentanedioic acid;
15 2-[[(2-naphthyl)hydroxyphosphinyl]methyl]pentanedioic acid;
2-[[(1-naphthyl)methylhydroxyphosphinyl]methyl]pentanedioic
acid;
2-[[(2-naphthyl)methylhydroxyphosphinyl]methyl]pentanedioic
acid;
20 2-[[(1-naphthyl)ethylhydroxyphosphinyl]methyl]pentanedioic
acid;
2-[[(2-naphthyl)ethylhydroxyphosphinyl]methyl]pentanedioic
acid;
2-[[(1-naphthyl)propylhydroxyphosphinyl]methyl]pentanedioic
25 acid;
2-[[(2-naphthyl)propylhydroxyphosphinyl]methyl]pentanedioic
acid;
2-[[(1-naphthyl)butylhydroxyphosphinyl]methyl]pentanedioic

acid;

2-[[(2-naphthyl) butylhydroxyphosphinyl] methyl] pentanedioic
acid; and

2-[[(phenylprop-2-enyl) hydroxyphosphinyl] methyl] pentanedioic
5 acid.

Especially preferred methods utilize compounds wherein R₁
is either a straight or branched aliphatic group or a
carbocyclic group, R₂ is ethyl which is substituted with a
10 carboxylic acid, and X is CH₂, are selected from the group
consisting of:

2-[(benzylhydroxyphosphinyl) methyl] pentanedioic acid;

2-[(phenylhydroxyphosphinyl) methyl] pentanedioic acid;

2-[[(hydroxy) phenylmethyl] hydroxyphosphinyl] methyl]
15 pentanedioic acid;

2-[(butylhydroxyphosphinyl) methyl] pentanedioic acid;

2-[[(3-methylbenzyl) hydroxyphosphinyl] methyl] pentanedioic
acid;

2-[[(3-phenylpropyl) hydroxyphosphinyl] methyl] pentanedioic acid;

20 2-[[(4-fluorophenyl) hydroxyphosphinyl] methyl] pentanedioic
acid;

2-[(methylhydroxyphosphinyl) methyl] pentanedioic acid;

2-[(phenylethylhydroxyphosphinyl) methyl] pentanedioic acid;

2-[[(4-methylbenzyl) hydroxyphosphinyl] methyl] pentanedioic
25 acid;

2-[[(4-fluorobenzyl) hydroxyphosphinyl] methyl] pentanedioic
acid;

2-[[(4-methoxybenzyl) hydroxyphosphinyl] methyl] pentanedioic

acid;

2-[[(2-fluorobenzyl)hydroxyphosphinyl]methyl]pentanedioic
acid;

2-[[(pentafluorobenzyl)hydroxyphosphinyl]methyl]pentanedioic
acid;

2-(phosphonomethyl)pentanedioic acid; and

2-[[(3-
trifluoromethylbenzyl)hydroxyphosphinyl]methyl]pentanedioic
acid.

10

Especially preferred methods utilize compounds wherein R_1 is a straight or branched aliphatic group or a carbocyclic group, R_2 is phenyl, and X is CH_2 , are selected from the group consisting of:

- 15 3-(methylhydroxyphosphinyl)-2-phenylpropanoic acid;
3-(ethylhydroxyphosphinyl)-2-phenylpropanoic acid;
3-(propylhydroxyphosphinyl)-2-phenylpropanoic acid;
3-(butylhydroxyphosphinyl)-2-phenylpropanoic acid;
3-(cyclohexylhydroxyphosphinyl)-2-phenylpropanoic acid;
20 3-((cyclohexyl)methylhydroxyphosphinyl)-2-phenylpropanoic
acid;
3-(phenylhydroxyphosphinyl)-2-phenylpropanoic acid;
3-(benzylhydroxyphosphinyl)-2-phenylpropanoic acid;
3-(phenylethylhydroxyphosphinyl)-2-phenylpropanoic acid;
25 3-(phenylpropylhydroxyphosphinyl)-2-phenylpropanoic acid;
3-(phenylbutylhydroxyphosphinyl)-2-phenylpropanoic acid;
3-[(2, 3, 4-trimethoxyphenyl)-3-hydroxyphosphinyl]-2-
phenylpropanoic acid;

3-[(1-naphthyl)hydroxyphosphinyl]-2-phenylpropanoic acid;
3-[(2-naphthyl)hydroxyphosphinyl]-2-phenylpropanoic acid;
3-[(1-naphthyl)methylhydroxyphosphinyl]-2-phenylpropanoic
acid;
5 3-[(2-naphthyl)methylhydroxyphosphinyl]-2-phenylpropanoic
acid;
3-[(1-naphthyl)ethylhydroxyphosphinyl]-2-phenylpropanoic acid;
3-[(2-naphthyl)ethylhydroxyphosphinyl]-2-phenylpropanoic acid;
3-[(1-naphthyl)propylhydroxyphosphinyl]-2-phenylpropanoic
10 acid;
3-[(2-naphthyl)propylhydroxyphosphinyl]-2-phenylpropanoic
acid;
3-[(1-naphthyl)butylhydroxyphosphinyl]-2-phenylpropanoic acid;
3-[(2-naphthyl)butylhydroxyphosphinyl]-2-phenylpropanoic acid;
15 and
3-[phenylprop-2-enylhydroxyphosphinyl]-2-phenylpropanoic acid.

Although not limited to any one species, a highly
preferred species where R_1 is a straight or branched aliphatic
20 group or a carbocyclic group, R_2 is ethyl which is substituted
with carboxylic acid, and X is CH_2 is
2-[[benzylhydroxyphosphinyl)methyl]pentanedioic acid.

Other preferred compounds of the present invention are
25 selected from the group consisting of:
hydroxyphosphinyl derivatives wherein X is CH_2 , R_1 is a
straight or branched aliphatic group or a carbocyclic group,
and R_2 is an C_2 - C_9 alkyl or alkenyl chain which is substituted

with a carboxylic acid. Exemplary species include:

- 2-[(methylhydroxyphosphinyl)methyl]hexanedioic acid;
- 2-[(benzylhydroxyphosphinyl)methyl]hexanedioic acid;
- 2-[(methylhydroxyphosphinyl)methyl]heptanedioic acid;
- 5 2-[(benzylhydroxyphosphinyl)methyl]heptanedioic acid;
- 2-[(methylhydroxyphosphinyl)methyl]octanedioic acid;
- 2-[(benzylhydroxyphosphinyl)methyl]octanedioic acid;
- 2-[(methylhydroxyphosphinyl)methyl]nonanedioic acid;
- 2-[(benzylhydroxyphosphinyl)methyl]nonanedioic acid;
- 10 2-[(methylhydroxyphosphinyl)methyl]decanedioic acid; and
- 2-[(benzylhydroxyphosphinyl)methyl]decanedioic acid.

Other preferred compounds are selected from the group consisting of:

- 15 hydroxyphosphinyl derivatives wherein X is CH_2 , R_1 is benzyl, and R_2 is a straight or branched aliphatic group or a carbocyclic group. Exemplary species include:
- 3-(benzylhydroxyphosphinyl)-2-methylpropanoic acid;
- 3-(benzylhydroxyphosphinyl)-2-ethylpropanoic acid;
- 20 3-(benzylhydroxyphosphinyl)-2-propylpropanoic acid;
- 3-(benzylhydroxyphosphinyl)-2-butylpropanoic acid;
- 3-(benzylhydroxyphosphinyl)-2-cyclohexylpropanoic acid;
- 3-(benzylhydroxyphosphinyl)-2-(cyclohexyl)methylpropanoic acid;
- 25 3-(benzylhydroxyphosphinyl)-2-phenylpropanoic acid;
- 3-(benzylhydroxyphosphinyl)-2-benzylpropanoic acid;
- 3-(benzylhydroxyphosphinyl)-2-phenylethylpropanoic acid;
- 3-(benzylhydroxyphosphinyl)-2-phenylpropylpropanoic acid;

3-(benzylhydroxyphosphinyl)-2-phenylbutylpropanoic acid;

3-(benzylhydroxyphosphinyl)-2-(2, 3, 4-trimethoxyphenyl)
propanoic acid;

3-(benzylhydroxyphosphinyl)-2-(1-naphthyl)propanoic acid;

5 3-(benzylhydroxyphosphinyl)-2-(2-naphthyl)propanoic acid;

3-(benzylhydroxyphosphinyl)-2-(1-naphthyl)methylpropanoic
acid;

3-(benzylhydroxyphosphinyl)-2-(2-naphthyl)methylpropanoic
acid;

10 3-(benzylhydroxyphosphinyl)-2-(1-naphthyl)ethylpropanoic acid;

3-(benzylhydroxyphosphinyl)-2-(2-naphthyl)ethylpropanoic acid;

3-(benzylhydroxyphosphinyl)-2-(1-naphthyl)propylpropanoic
acid;

3-(benzylhydroxyphosphinyl)-2-(2-naphthyl)propylpropanoic
15 acid;

3-(benzylhydroxyphosphinyl)-2-(1-naphthyl)butylpropanoic acid;

3-(benzylhydroxyphosphinyl)-2-(2-naphthyl)butylpropanoic acid;
and

3-(benzylhydroxyphosphinyl)-2-phenylprop-2-enylpropanoic acid.

20

Especially preferred methods use compounds wherein R₁ is
said alkyl, alkenyl, cycloalkyl, or aryl group which is
substituted with a heterocyclic group, R₂ is ethyl which is
substituted with a carboxylic acid, and X is CH₂, are selected
25 from the group consisting of:

2-[[[(2-pyridyl)methylhydroxyphosphinyl]methyl]pentanedioic
acid;

2-[[[(3-pyridyl)methylhydroxyphosphinyl]methyl]pentanedioic

acid;

2-[[(4-pyridyl)methylhydroxyphosphinyl]methyl]pentanedioic
acid;

5 2-[[(3-pyridyl)ethylhydroxyphosphinyl]methyl]pentanedioic
acid;

2-[[(3-pyridyl)propylhydroxyphosphinyl]methyl]pentanedioic
acid;

2-[[(tetrahydrofuranyl)methylhydroxyphosphinyl]methyl]
pentanedioic acid;

10 2-[[(tetrahydrofuranyl)ethylhydroxyphosphinyl]methyl]
pentanedioic acid;

2-[[(tetrahydrofuranyl)propylhydroxyphosphinyl]methyl]
pentanedioic acid;

15 2-[[(2-indolyl)methylhydroxyphosphinyl]methyl]pentanedioic
acid;

2-[[(3-indolyl)methylhydroxyphosphinyl]methyl]pentanedioic
acid;

2-[[(4-indolyl)methylhydroxyphosphinyl]methyl]pentanedioic
acid;

20 2-[[(3-indolyl)ethylhydroxyphosphinyl]methyl]pentanedioic
acid;

2-[[(3-indolyl)propylhydroxyphosphinyl]methyl]pentanedioic
acid;

25 2-[[(2-thienyl)methylhydroxyphosphinyl]methyl]pentanedioic
acid;

2-[[(3-thienyl)methylhydroxyphosphinyl]methyl]pentanedioic
acid;

2-[[(4-thienyl)methylhydroxyphosphinyl]methyl]pentanedioic

acid;

2-[[[(3-thienyl)ethylhydroxyphosphinyl]methyl]pentanedioic
acid; and

2-[[[(3-thienyl)propylhydroxyphosphinyl]methyl]pentanedioic
5 acid.

Especially preferred methods use compounds wherein R₁ is
said alkyl, alkenyl, cycloalkyl, or aryl group which is
substituted with a heterocyclic group, R₂ is phenyl, and X is
10 CH₂, are selected from the group consisting of:

3-[(2-pyridyl)methylhydroxyphosphinyl]-2-phenylpropanoic acid;

3-[(3-pyridyl)methylhydroxyphosphinyl]-2-phenylpropanoic acid;

3-[(4-pyridyl)methylhydroxyphosphinyl]-2-phenylpropanoic acid;

3-[(3-pyridyl)ethylhydroxyphosphinyl]-2-phenylpropanoic acid;

15 3-[(3-pyridyl)propylhydroxyphosphinyl]-2-phenylpropanoic acid;

3-[(tetrahydrofuranyl)methylhydroxyphosphinyl]-2-phenyl
propanoic acid;

3-[(tetrahydrofuranyl)ethylhydroxyphosphinyl]-2-phenyl
propanoic acid;

20 3-[(tetrahydrofuranyl)propylhydroxyphosphinyl]-2-phenyl
propanoic acid;

3-[(2-indolyl)methylhydroxyphosphinyl]-2-phenylpropanoic acid;

3-[(3-indolyl)methylhydroxyphosphinyl]-2-phenylpropanoic acid;

3-[(4-indolyl)methylhydroxyphosphinyl]-2-phenylpropanoic acid;

25 3-[(3-indolyl)ethylhydroxyphosphinyl]-2-phenylpropanoic acid;

3-[(3-indolyl)propylhydroxyphosphinyl]-2-phenylpropanoic acid;

3-[(2-thienyl)methylhydroxyphosphinyl]-2-phenylpropanoic acid;

3-[(3-thienyl)methylhydroxyphosphinyl]-2-phenylpropanoic acid;

3-[(4-thienyl)methylhydroxyphosphinyl]-2-phenylpropanoic acid;
3-[(3-thienyl)ethylhydroxyphosphinyl]-2-phenylpropanoic acid;
and
3-[(3-thienyl)propylhydroxyphosphinyl]-2-phenylpropanoic acid.


5

Especially preferred methods use compounds wherein R₁ is benzyl, R₂ is said alkyl, alkenyl, cycloalkyl, or aryl group which is substituted with a heterocyclic group, and X is CH₂, are selected from the group consisting of:

- 10 3-(benzylhydroxyphosphinyl)-2-(2-pyridyl)methylpropanoic acid;
3-(benzylhydroxyphosphinyl)-2-(3-pyridyl)methylpropanoic acid;
3-(benzylhydroxyphosphinyl)-2-(4-pyridyl)methylpropanoic acid;
3-(benzylhydroxyphosphinyl)-2-(3-pyridyl)ethylpropanoic acid;
3-(benzylhydroxyphosphinyl)-2-(3-pyridyl)propylpropanoic acid;
15 3-(benzylhydroxyphosphinyl)-2-(tetrahydrofuranyl)methyl
propanoic acid;
3-(benzylhydroxyphosphinyl)-2-(tetrahydrofuranyl)ethyl
propanoic acid;
3-(benzylhydroxyphosphinyl)-2-(tetrahydrofuranyl)propyl
20 propanoic acid;
3-(benzylhydroxyphosphinyl)-2-(2-indolyl)methylpropanoic acid;
3-(benzylhydroxyphosphinyl)-2-(3-indolyl)methylpropanoic acid;
3-(benzylhydroxyphosphinyl)-2-(4-indolyl)methylpropanoic acid;
3-(benzylhydroxyphosphinyl)-2-(3-indolyl)ethylpropanoic acid;
25 3-(benzylhydroxyphosphinyl)-2-(3-indolyl)propylpropanoic acid;
3-(benzylhydroxyphosphinyl)-2-(2-thienyl)methylpropanoic acid;
3-(benzylhydroxyphosphinyl)-2-(3-thienyl)methylpropanoic acid;
3-(benzylhydroxyphosphinyl)-2-(4-thienyl)methylpropanoic acid;

3-(benzylhydroxyphosphinyl)-2-(3-thienyl)ethylpropanoic acid;
and
3-(benzylhydroxyphosphinyl)-2-(3-thienyl)propylpropanoic acid.

5 In another preferred embodiment, the R groups are heterocyclic substituents illustrated by the compounds selected from the group having Formula II:



II

10 wherein

R₁ is Ar₁; and

R₂ is C₁-C₉ straight or branched chain alkyl, C₂-C₉ straight or branched chain alkenyl group, C₃-C₈ cycloalkyl, C₅-C₇ cycloalkenyl, or Ar₁, wherein said
15 alkyl, alkenyl, cycloalkyl, cycloalkenyl or aryl group may be optionally substituted with carboxylic acid.

Especially preferred methods use compounds wherein R₁ is a
20 heterocyclic group, R₂ is ethyl which is substituted with carboxylic acid, and X is CH₂, are selected from the group consisting of:

2-[[(2-pyridyl)hydroxyphosphinyl]methyl]pentanedioic acid;

2-[[(3-pyridyl)hydroxyphosphinyl]methyl]pentanedioic acid;

25 2-[[(4-pyridyl)hydroxyphosphinyl]methyl]pentanedioic acid;

2-[[[(tetrahydrofuranyl)hydroxyphosphinyl]methyl]pentanedioic acid;

2-[[[(2-indolyl)hydroxyphosphinyl]methyl]pentanedioic acid;

2-[[[(3-indolyl)hydroxyphosphinyl]methyl]pentanedioic acid;

5 2-[[[(4-indolyl)hydroxyphosphinyl]methyl]pentanedioic acid;

2-[[[(2-thienyl)hydroxyphosphinyl]methyl]pentanedioic acid;

2-[[[(3-thienyl)hydroxyphosphinyl]methyl]pentanedioic acid; and

2-[[[(4-thienyl)hydroxyphosphinyl]methyl]pentanedioic acid.

10 Compounds of the present invention wherein R₁ is a heterocyclic group, R₂ is phenyl, and X is CH₂, are selected from the group consisting of:

3-[(2-pyridyl)hydroxyphosphinyl]-2-phenylpropanoic acid;

3-[(3-pyridyl)hydroxyphosphinyl]-2-phenylpropanoic acid;

15 3-[(4-pyridyl)hydroxyphosphinyl]-2-phenylpropanoic acid;

3-[(tetrahydrofuranyl)hydroxyphosphinyl]-2-phenylpropanoic acid;

3-[(2-indolyl)hydroxyphosphinyl]-2-phenylpropanoic acid;

3-[(3-indolyl)hydroxyphosphinyl]-2-phenylpropanoic acid;


20 3-[(4-indolyl)hydroxyphosphinyl]-2-phenylpropanoic acid;

3-[(2-thienyl)hydroxyphosphinyl]-2-phenylpropanoic acid;

3-[(3-thienyl)hydroxyphosphinyl]-2-phenylpropanoic acid; and

3-[(4-thienyl)hydroxyphosphinyl]-2-phenylpropanoic acid.

25 Compounds are also preferably selected from the group of formula II:



wherein

II

R_1 is hydrogen, C_1 - C_8 straight or branched chain alkyl,
 C_2 - C_8 straight or branched chain alkenyl group, C_3 - C_8
cycloalkyl, C_5 - C_8 cycloalkenyl, or Ar_1 ; and
 R_2 is Ar_1 , wherein said aryl group may be optionally
substituted with carboxylic acid.


Particular species wherein R_2 is heterocyclic may be easily made and used by persons of ordinary skill in the art in accordance with the teachings provided herein and known in the art.

Compounds wherein R_1 is benzyl, R_2 is heterocyclic, and X is CH_2 are selected from the group consisting of:

3-(benzylhydroxyphosphinyl)-2-(2-pyridyl)propanoic acid;
3-(benzylhydroxyphosphinyl)-2-(3-pyridyl)propanoic acid;
3-(benzylhydroxyphosphinyl)-2-(4-pyridyl)propanoic acid;
3-(benzylhydroxyphosphinyl)-2-(tetrahydrofuranyl)propanoic
acid;
3-(benzylhydroxyphosphinyl)-2-(2-indolyl)propanoic acid;
3-(benzylhydroxyphosphinyl)-2-(3-indolyl)propanoic acid;
3-(benzylhydroxyphosphinyl)-2-(4-indolyl)propanoic acid;
3-(benzylhydroxyphosphinyl)-2-(2-thienyl)propanoic acid;
3-(benzylhydroxyphosphinyl)-2-(3-thienyl)propanoic acid; and

3-(benzylhydroxyphosphinyl)-2-(4-thienyl)propanoic acid.

Preferred compounds are also selected from formula III:



III


5 wherein

R_1 is hydrogen, C_1 - C_8 straight or branched chain alkyl, C_2 - C_8 straight or branched chain alkenyl group, C_3 - C_8 cycloalkyl, C_5 - C_8 cycloalkenyl, or Ar_1 ; and

10 R_2 is C_1 - C_8 straight or branched chain alkyl, C_2 - C_8 straight or branched chain alkenyl group, C_3 - C_8 cycloalkyl, C_5 - C_8 cycloalkenyl, or Ar_1 , wherein said alkyl, alkenyl, cycloalkyl, cycloalkenyl or aryl group may be optionally substituted with carboxylic acid.

15

In another preferred embodiment, the R groups are straight or branched or carbocyclic substituents illustrated by the compounds selected from the group of formula III:



20

III

wherein

R₁ is hydrogen, C₁-C₈ straight or branched chain alkyl, C₂-C₈ straight or branched chain alkenyl group, C₃-C₈ cycloalkyl, C₅-C₈ cycloalkenyl, 1-naphthyl, 2-naphthyl, or phenyl; and

5 R₂ is C₁-C₈ straight or branched chain alkyl, C₂-C₈ straight or branched chain alkenyl group, C₃-C₈ cycloalkyl, C₅-C₈ cycloalkenyl, 1-naphthyl, 2-naphthyl, or phenyl, wherein said alkyl, alkenyl, cycloalkyl, cycloalkenyl, 1-naphthyl, 2-naphthyl, or
10 phenyl group may be optionally substituted with carboxylic acid.

Especially preferred compounds of Formula III of the present invention wherein R₁ is a straight or branched
15 aliphatic group or a carbocyclic group and R₂ is ethyl which is substituted with a carboxylic acid are selected from the group consisting of:

2-[[methylhydroxyphosphinyl]oxy]pentanedioic acid;

2-[[ethylhydroxyphosphinyl]oxy]pentanedioic acid;

20 2-[[propylhydroxyphosphinyl]oxy]pentanedioic acid;

2-[[butylhydroxyphosphinyl]oxy]pentanedioic acid;

2-[[cyclohexylhydroxyphosphinyl]oxy]pentanedioic acid;

2-[[[(cyclohexyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;

2-[[phenylhydroxyphosphinyl]oxy]pentanedioic acid;

25 2-[[benzylhydroxyphosphinyl]oxy]pentanedioic acid;

2-[[phenylethylhydroxyphosphinyl]oxy]pentanedioic acid;

2-[[phenylpropylhydroxyphosphinyl]oxy]pentanedioic acid;

2-[[phenylbutylhydroxyphosphinyl]oxy]pentanedioic acid;

- 2-[[[(4-methylbenzyl)hydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(4-fluorobenzyl)hydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(2-fluorobenzyl)hydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(pentafluorobenzyl)hydroxyphosphinyl]oxy]pentanedioic
5 acid;
2-[[[(methoxybenzyl)hydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(4-fluorophenyl)hydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(hydroxy)phenylmethyl]hydroxyphosphinyl]oxy]pentanedioic
acid;
10 2-[[[(3-methylbenzyl)hydroxyphosphinyl]oxy]pentanedioic acid;
2-(phosphono)oxy]pentanedioic acid;
2-[[[(3-
trifluoromethylbenzyl)hydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(2, 3, 4-trimethoxyphenyl)hydroxyphosphinyl]oxy]
15 pentanedioic acid;
2-[[[(1-naphthyl)hydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(2-naphthyl)hydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(1-naphthyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(2-naphthyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;
20 2-[[[(1-naphthyl)ethylhydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(2-naphthyl)ethylhydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(1-naphthyl)propylhydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(2-naphthyl)propylhydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(1-naphthyl)butylhydroxyphosphinyl]oxy]pentanedioic acid;
25 2-[[[(2-naphthyl)butylhydroxyphosphinyl]oxy]pentanedioic acid;
and
2-[[[(phenylprop-2-enyl)hydroxyphosphinyl]oxy]pentanedioic
acid.

Especially preferred compounds of Formula III of the present invention wherein R_1 is a straight or branched aliphatic group or a carbocyclic group and R_2 is ethyl which is substituted with a carboxylic acid are selected from the group consisting of:

- 2-[(benzylhydroxyphosphinyl)oxy]pentanedioic acid;
- 2-[(phenylhydroxyphosphinyl)oxy]pentanedioic acid;
- 2-[[[(hydroxy)phenylmethyl]hydroxyphosphinyl]oxy]pentanedioic acid;
- 2-[(butylhydroxyphosphinyl)oxy]pentanedioic acid;
- 2-[[[3-methylbenzyl]hydroxyphosphinyl]oxy]pentanedioic acid;
- 2-[(3-phenylpropylhydroxyphosphinyl)oxy]pentanedioic acid;
- 2-[[[4-fluorophenyl]hydroxyphosphinyl]oxy]pentanedioic acid;
- 2-[(methylhydroxyphosphinyl)oxy]pentanedioic acid;
- 2-[(phenylethylhydroxyphosphinyl)oxy]pentanedioic acid;
- 2-[[[4-methylbenzyl]hydroxyphosphinyl]oxy]pentanedioic acid;
- 2-[[[4-fluorobenzyl]hydroxyphosphinyl]oxy]pentanedioic acid;
- 2-[[[4-methoxybenzyl]hydroxyphosphinyl]oxy]pentanedioic acid;
- 2-[[[2-fluorobenzyl]hydroxyphosphinyl]oxy]pentanedioic acid;
- 2-[[[pentafluorobenzyl]hydroxyphosphinyl]oxy]pentanedioic acid;
- 2-[(phosphono)oxy]pentanedioic acid; and
- 2-[[[3-trifluoromethylbenzyl]hydroxyphosphinyl]oxy]pentanedioic acid.

Compounds of the present invention wherein R_1 is a straight or branched aliphatic group or a carbocyclic group, R_2 is phenyl, and X is oxygen are selected from the group

consisting of:

- 2-[[methylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
2-[[ethylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
2-[[propylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
5 2-[[butylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
2-[[cyclohexylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
2-[[[(cyclohexyl)methylhydroxyphosphinyl]oxy]-2-phenylethanoic
acid;
2-[[phenylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
10 2-[[benzylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
2-[[phenylethylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
2-[[phenylpropylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
2-[[phenylbutylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
2-[[[(2, 3, 4-trimethoxyphenyl)-3-hydroxyphosphinyl]oxy]-2-
15 phenylethanoic acid;
2-[[[(1-naphthyl)hydroxyphosphinyl]oxy]-2-phenylethanoic acid;
2-[[[(2-naphthyl)hydroxyphosphinyl]oxy]-2-phenylethanoic acid;
2-[[[(1-naphthyl)methylhydroxyphosphinyl]oxy]-2-phenylethanoic
acid;
20 2-[[[(2-naphthyl)methylhydroxyphosphinyl]oxy]-2-phenylethanoic
acid;
2-[[[(1-naphthyl)ethylhydroxyphosphinyl]oxy]-2-phenylethanoic
acid;
2-[[[(2-naphthyl)ethylhydroxyphosphinyl]oxy]-2-phenylethanoic
25 acid;
2-[[[(1-naphthyl)propylhydroxyphosphinyl]oxy]-2-phenylethanoic
acid;
2-[[[(2-naphthyl)propylhydroxyphosphinyl]oxy]-2-phenylethanoic

acid;

2-[[[(1-naphthyl)butylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;

2-[[[(2-naphthyl)butylhydroxyphosphinyl]oxy]-2-phenylethanoic acid; and

2-[[[phenylprop-2-enylhydroxyphosphinyl]oxy]-2-phenylethanoic acid.

Although not limited to any one particular species, a highly preferred species where R_1 is a straight or branched aliphatic group or a carbocyclic group, R_2 is ethyl which is substituted with carboxylic acid, and X is oxygen is 2-[[benzylhydroxyphosphinyl]oxy]pentanedioic acid.

Other especially preferred compounds of the present invention are selected from the group consisting of: phosphonate derivatives wherein X is oxygen, R_1 is a straight or branched aliphatic group or a carbocyclic group, and R_2 is an C_2 - C_8 alkyl or alkenyl chain which is substituted with a carboxylic acid. Exemplary species include:

2-[(methylhydroxyphosphinyl)oxy]hexanedioic acid;

2-[(benzylhydroxyphosphinyl)oxy]hexanedioic acid;

2-[(methylhydroxyphosphinyl)oxy]heptanedioic acid;

2-[(benzylhydroxyphosphinyl)oxy]heptanedioic acid;

2-[(methylhydroxyphosphinyl)oxy]octanedioic acid;

2-[(benzylhydroxyphosphinyl)oxy]octanedioic acid;

2-[(methylhydroxyphosphinyl)oxy]nonanedioic acid;

2-[(benzylhydroxyphosphinyl)oxy]nonanedioic acid;

2-[(methylhydroxyphosphinyl)oxy]decanedioic acid; and
2-[(benzylhydroxyphosphinyl)oxy]decanedioic acid.

Other especially preferred compounds are selected from
5 the group consisting of:
phosphonate derivatives wherein X is oxygen, R₁ is benzyl, and
R₂ is a straight or branched aliphatic group or a carbocyclic
group. Exemplary species include:

2-[(benzylhydroxyphosphinyl)oxy]-2-methylethanoic acid;
10 2-[(benzylhydroxyphosphinyl)oxy]-2-ethylethanoic acid;
2-[(benzylhydroxyphosphinyl)oxy]-2-propylethanoic acid;
2-[(benzylhydroxyphosphinyl)oxy]-2-butylethanoic acid;
2-[(benzylhydroxyphosphinyl)oxy]-2-cyclohexylethanoic acid;
2-[(benzylhydroxyphosphinyl)oxy]-2-(cyclohexyl) methylethanoic
15 acid;
2-[(benzylhydroxyphosphinyl)oxy]-2-phenylethanoic acid;
2-[(benzylhydroxyphosphinyl)oxy]-2-benzylethanoic acid;
2-[(benzylhydroxyphosphinyl)oxy]-2-phenylethylethanoic acid;
2-[(benzylhydroxyphosphinyl)oxy]-2-phenylpropylethanoic acid;
20 2-[(benzylhydroxyphosphinyl)oxy]-2-phenylbutylethanoic acid;
2-[(benzylhydroxyphosphinyl)oxy]-2-(2, 3, 4-
trimethoxyphenyl)ethanoic acid;
2-[(benzylhydroxyphosphinyl)oxy]-2-(1-naphthyl)ethanoic acid;
2-[(benzylhydroxyphosphinyl)oxy]-2-(2-naphthyl)ethanoic acid;
25 2-[(benzylhydroxyphosphinyl)oxy]-2-(1-naphthyl)methylethanoic
acid;
2-[(benzylhydroxyphosphinyl)oxy]-2-(2-naphthyl)methylethanoic
acid;

2-[[benzylhydroxyphosphinyl]oxy]-2-(1-naphthyl)ethylethanoic acid;

2-[[benzylhydroxyphosphinyl]oxy]-2-(2-naphthyl)ethylethanoic acid;

5 2-[[benzylhydroxyphosphinyl]oxy]-2-(1-naphthyl)propylethanoic acid;

2-[[benzylhydroxyphosphinyl]oxy]-2-(2-naphthyl)propylethanoic acid;

10 2-[[benzylhydroxyphosphinyl]oxy]-2-(1-naphthyl)butylethanoic acid;

2-[[benzylhydroxyphosphinyl]oxy]-2-(2-naphthyl)butylethanoic acid; and

2-[[benzylhydroxyphosphinyl]oxy]-2-phenylprop-2-enylethanoic acid.

15

Especially preferred methods utilize compounds wherein R_1 is said alkyl, alkenyl, cycloalkyl, or aryl group which is substituted with a heterocyclic group, R_2 is ethyl which is substituted with a carboxylic acid, and X is oxygen are
20 selected from the group consisting of:

2-[[[(2-pyridyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;

2-[[[(3-pyridyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;

2-[[[(4-pyridyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;

2-[[[(3-pyridyl)ethylhydroxyphosphinyl]oxy]pentanedioic acid;

25 2-[[[(3-pyridyl)propylhydroxyphosphinyl]oxy]pentanedioic acid;

2-[[[(tetrahydrofuranyl)methylhydroxyphosphinyl]oxy]
pentanedioic acid;

2-[[[(tetrahydrofuranyl)ethylhydroxyphosphinyl]oxy]

pentanedioic acid;

2-[[[(tetrahydrofuranyl)propylhydroxyphosphinyl]oxy]

pentanedioic acid;

2-[[[(2-indolyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;

5 2-[[[(3-indolyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;

2-[[[(4-indolyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;

2-[[[(3-indolyl)ethylhydroxyphosphinyl]oxy]pentanedioic acid;

2-[[[(3-indolyl)propylhydroxyphosphinyl]oxy]pentanedioic acid;

2-[[[(2-thienyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;

10 2-[[[(3-thienyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;

2-[[[(4-thienyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;

2-[[[(3-thienyl)ethylhydroxyphosphinyl]oxy]pentanedioic acid;

and

2-[[[(3-thienyl)propylhydroxyphosphinyl]oxy]pentanedioic acid.

15

Especially preferred methods utilize compounds wherein R_1 is said alkyl, alkenyl, cycloalkyl, or aryl group which is substituted with a heterocyclic group, R_2 is phenyl, and X is oxygen are selected from the group consisting of:

20 2-[[[(2-pyridyl)methylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;

2-[[[(3-pyridyl)methylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;

25 2-[[[(4-pyridyl)methylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;

2-[[[(3-pyridyl)ethylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;

2-[[[(3-pyridyl)propylhydroxyphosphinyl]oxy]-2-phenylethanoic

acid;

2-[[[(tetrahydrofuranyl)methylhydroxyphosphinyl]oxy]-2-phenyl
ethanoic acid;

2-[[[(tetrahydrofuranyl)ethylhydroxyphosphinyl]oxy]-2-phenyl
5 ethanoic acid;

2-[[[(tetrahydrofuranyl)propylhydroxyphosphinyl]oxy]-2-phenyl
ethanoic acid;

2-[[[(2-indolyl)methylhydroxyphosphinyl]oxy]-2-phenylethanoic
acid;

10 2-[[[(3-indolyl)methylhydroxyphosphinyl]oxy]-2-phenylethanoic
acid;

2-[[[(4-indolyl)methylhydroxyphosphinyl]oxy]-2-phenylethanoic
acid;

2-[[[(3-indolyl)ethylhydroxyphosphinyl]oxy]-2-phenylethanoic
15 acid;

2-[[[(3-indolyl)propylhydroxyphosphinyl]oxy]-2-phenylethanoic
acid;

2-[[[(2-thienyl)methylhydroxyphosphinyl]oxy]-2-phenylethanoic
acid;

20 2-[[[(3-thienyl)methylhydroxyphosphinyl]oxy]-2-phenylethanoic
acid;

2-[[[(4-thienyl)methylhydroxyphosphinyl]oxy]-2-phenylethanoic
acid;

2-[[[(3-thienyl)ethylhydroxyphosphinyl]oxy]-2-phenylethanoic
25 acid; and

2-[[[(3-thienyl)propylhydroxyphosphinyl]oxy]-2-phenylethanoic
acid.

Especially preferred methods utilize compounds wherein R_1 is benzyl, R_2 is said alkyl, alkenyl, cycloalkyl, or aryl group which is substituted with a heterocyclic group, and X is oxygen are selected from the group consisting of:

5 2-[[benzylhydroxyphosphinyl]oxy]-2-(2-pyridyl)methylethanoic acid;

2-[[benzylhydroxyphosphinyl]oxy]-2-(3-pyridyl)methylethanoic acid;

10 2-[[benzylhydroxyphosphinyl]oxy]-2-(4-pyridyl)methylethanoic acid;

2-[[benzylhydroxyphosphinyl]oxy]-2-(3-pyridyl)ethylethanoic acid;

2-[[benzylhydroxyphosphinyl]oxy]-2-(3-pyridyl)propylethanoic acid;

15 2-[[benzylhydroxyphosphinyl]oxy]-2-(tetrahydrofuranyl)methylethanoic acid;

2-[[benzylhydroxyphosphinyl]oxy]-2-(tetrahydrofuranyl)ethylethanoic acid;

20 2-[[benzylhydroxyphosphinyl]oxy]-2-(tetrahydrofuranyl)propylethanoic acid;

2-[[benzylhydroxyphosphinyl]oxy]-2-(2-indolyl)methylethanoic acid;

2-[[benzylhydroxyphosphinyl]oxy]-2-(3-indolyl)methylethanoic acid;

25 2-[[benzylhydroxyphosphinyl]oxy]-2-(4-indolyl)methylethanoic acid;

2-[[benzylhydroxyphosphinyl]oxy]-2-(3-indolyl)ethylethanoic acid;

2-[[benzylhydroxyphosphinyl]oxy]-2-(3-indolyl)propylethanoic acid;

2-[[benzylhydroxyphosphinyl]oxy]-2-(2-thienyl)methylethanoic acid;


5 2-[[benzylhydroxyphosphinyl]oxy]-2-(3-thienyl)methylethanoic acid;

2-[[benzylhydroxyphosphinyl]oxy]-2-(4-thienyl)methylethanoic acid;

10 2-[[benzylhydroxyphosphinyl]oxy]-2-(3-thienyl)ethylethanoic acid; and

2-[[benzylhydroxyphosphinyl]oxy]-2-(3-thienyl)propylethanoic acid.

15 In another preferred embodiment, R_1 is a heterocyclic substituent illustrated by the compounds selected from the group of formula III:



wherein

III

20 R_1 is Ar_1 ; and

R_2 is $\text{C}_1\text{-C}_9$ straight or branched chain alkyl, $\text{C}_2\text{-C}_9$ straight or branched chain alkenyl group, $\text{C}_1\text{-C}_8$ cycloalkyl, $\text{C}_5\text{-C}_9$ cycloalkenyl, or Ar_2 , wherein said alkyl, alkenyl, cycloalkyl, cycloalkenyl or aryl group may be optionally substituted with carboxylic

25

acid.

Especially preferred compounds wherein R_1 is a heterocyclic group, R_2 is ethyl which is substituted with carboxylic acid, and X is oxygen are selected from the group consisting of:


- 2-[[[(2-pyridyl)hydroxyphosphinyl]oxy]pentanedioic acid;
- 2-[[[(3-pyridyl)hydroxyphosphinyl]oxy]pentanedioic acid;
- 2-[[[(4-pyridyl)hydroxyphosphinyl]oxy]pentanedioic acid;
- 10 2-[[[(tetrahydrofuranyl)hydroxyphosphinyl]oxy]pentanedioic acid;
- 2-[[[(2-indolyl)hydroxyphosphinyl]oxy]pentanedioic acid;
- 2-[[[(3-indolyl)hydroxyphosphinyl]oxy]pentanedioic acid;
- 2-[[[(4-indolyl)hydroxyphosphinyl]oxy]pentanedioic acid;
- 15 2-[[[(2-thienyl)hydroxyphosphinyl]oxy]pentanedioic acid;
- 2-[[[(3-thienyl)hydroxyphosphinyl]oxy]pentanedioic acid; and
- 2-[[[(4-thienyl)hydroxyphosphinyl]oxy]pentanedioic acid.

Compounds of the present invention wherein R_1 is a heterocyclic group, R_2 is phenyl, and X is oxygen are selected from the group consisting of:

- 2-[[[(2-pyridyl)hydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[[(3-pyridyl)hydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[[(4-pyridyl)hydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 25 2-[[[(tetrahydrofuranyl)hydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[[(2-indolyl)hydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[[(3-indolyl)hydroxyphosphinyl]oxy]-2-phenylethanoic acid;

- 2-[[[(4-indolyl)hydroxyphosphinyl]oxy]-2-phenylethanoic acid;
2-[[[(2-thienyl)hydroxyphosphinyl]oxy]-2-phenylethanoic acid;
2-[[[(3-thienyl)hydroxyphosphinyl]oxy]-2-phenylethanoic acid;
and
5 2-[[[(4-thienyl)hydroxyphosphinyl]oxy]-2-phenylethanoic acid.

Compounds are also preferably selected from the group of formula III:



10

wherein

III


- R_1 is hydrogen, C_1 - C_8 straight or branched chain alkyl, C_2 - C_8 straight or branched chain alkenyl group, C_3 - C_8 cycloalkyl, C_5 - C_7 cycloalkenyl, or Ar_1 ; and
15 R_2 is Ar_1 , wherein said aryl group may be optionally substituted with carboxylic acid.

Particular species wherein R_2 is heterocyclic may be easily made and used by persons of ordinary skill in the art
20 in accordance with the teachings provided herein and known in the art.

Compounds of the present invention wherein R_1 is benzyl, R_2 is a heterocyclic group, and X is oxygen are selected from the group consisting of:
25 2-[[benzylhydroxyphosphinyl]oxy]-2-(2-pyridyl)ethanoic acid;

2-[[benzylhydroxyphosphinyl]oxy]-2-(3-pyridyl)ethanoic acid;
 2-[[benzylhydroxyphosphinyl]oxy]-2-(4-pyridyl)ethanoic acid;
 2-[[benzylhydroxyphosphinyl]oxy]-2-(tetrahydrofuranyl)ethanoic
 acid;
 5 2-[[benzylhydroxyphosphinyl]oxy]-2-(2-indolyl)ethanoic acid;
 2-[[benzylhydroxyphosphinyl]oxy]-2-(3-indolyl)ethanoic acid;
 2-[[benzylhydroxyphosphinyl]oxy]-2-(4-indolyl)ethanoic acid;
 2-[[benzylhydroxyphosphinyl]oxy]-2-(2-thienyl)ethanoic acid;
 2-[[benzylhydroxyphosphinyl]oxy]-2-(3-thienyl)ethanoic acid;
 10 and
 2-[[benzylhydroxyphosphinyl]oxy]-2-(4-thienyl)ethanoic acid.

Preferred phosphoramidate compounds are selected from
 formula IV:



15


IV

wherein

R₁ is hydrogen, C₁-C₈ straight or branched chain alkyl,
 C₂-C₈ straight or branched chain alkenyl group, C₃-C₈
 20 cycloalkyl, C₅-C₇ cycloalkenyl, or Ar₁; and
 R₂ is C₁-C₈ straight or branched chain alkyl, C₂-C₈
 straight or branched chain alkenyl group, C₃-C₈
 cycloalkyl, C₅-C₇ cycloalkenyl, or Ar₁, wherein said
 alkyl, alkenyl, cycloalkyl, cycloalkenyl or aryl
 25 group may be optionally substituted with carboxylic

acid.

In a preferred embodiment, the R groups are straight
 branched aliphatic or carbocyclic substituents illustrated by
 5 the compounds selected from the group of formula IV:



IV

wherein

- 10 R_1 is hydrogen, C_1 - C_8 straight or branched chain alkyl, C_2 - C_8 straight or branched chain alkenyl group, C_3 - C_8 cycloalkyl, C_5 - C_8 cycloalkenyl, 1-naphthyl, 2-naphthyl, or phenyl; and
- 15 R_2 is C_1 - C_8 straight or branched chain alkyl, C_2 - C_8 straight or branched chain alkenyl group, C_3 - C_8 cycloalkyl, C_5 - C_8 cycloalkenyl, 1-naphthyl, 2-naphthyl, or phenyl, wherein said alkyl, alkenyl, cycloalkyl, cycloalkenyl, 1-naphthyl, 2-naphthyl, or phenyl group may be optionally substituted with
- 20 carboxylic acid.

Especially preferred compounds of Formula IV of the
 present invention wherein R_1 is a straight or branched
 aliphatic group or a carbocyclic group, R_2 is ethyl which is
 substituted with a carboxylic acid, and the NR_1 is amino are
 25 selected from the group consisting of:

- 2-[[methylhydroxyphosphinyl]amino]pentanedioic acid;
2-[[ethylhydroxyphosphinyl]amino]pentanedioic acid;
2-[[propylhydroxyphosphinyl]amino]pentanedioic acid;
2-[[butylhydroxyphosphinyl]amino]pentanedioic acid;
5 2-[[cyclohexylhydroxyphosphinyl]amino]pentanedioic acid;
2-[[[(cyclohexyl)methylhydroxyphosphinyl]amino]pentanedioic
acid;
2-[[phenylhydroxyphosphinyl]amino]pentanedioic acid;
2-[[benzylhydroxyphosphinyl]amino]pentanedioic acid;
10 2-[[phenylethylhydroxyphosphinyl]amino]pentanedioic acid;
2-[[phenylpropylhydroxyphosphinyl]amino]pentanedioic acid;
2-[[phenylbutylhydroxyphosphinyl]amino]pentanedioic acid;
2-[[[(4-methylbenzyl)hydroxyphosphinyl]amino]pentanedioic acid;
2-[[[(4-fluorobenzyl)hydroxyphosphinyl]amino]pentanedioic acid;
15 2-[[[(2-fluorobenzyl)hydroxyphosphinyl]amino]pentanedioic acid;
2-[[[(pentafluorobenzyl)hydroxyphosphinyl]amino]pentanedioic
acid;
2-[[[(methoxybenzyl)hydroxyphosphinyl]amino]pentanedioic acid;
2-[[[(4-fluorophenyl)hydroxyphosphinyl]amino]pentanedioic acid;
20 2-[[[(hydroxy)phenylmethyl]hydroxyphosphinyl]amino]
pentanedioic acid;
2-[[[(3-methylbenzyl)hydroxyphosphinyl]amino]pentanedioic acid;
2-[(phosphono)amino]pentanedioic acid;
2-[[[(3-
25 trifluoromethylbenzyl)hydroxyphosphinyl]amino]pentanedioic
acid;
2-[[[(2, 3, 4-trimethoxyphenyl)hydroxyphosphinyl]amino]
pentanedioic acid;

2-[[[(1-naphthyl)hydroxyphosphinyl]amino]pentanedioic acid;
2-[[[(2-naphthyl)hydroxyphosphinyl]amino]pentanedioic acid;
2-[[[(1-naphthyl)methylhydroxyphosphinyl]amino]pentanedioic
acid;

5 2-[[[(2-naphthyl)methylhydroxyphosphinyl]amino]pentanedioic
acid;

2-[[[(1-naphthyl)ethylhydroxyphosphinyl]amino]pentanedioic
acid;

2-[[[(2-naphthyl)ethylhydroxyphosphinyl]amino]pentanedioic
10 acid;

2-[[[(1-naphthyl)propylhydroxyphosphinyl]amino]pentanedioic
acid;

2-[[[(2-naphthyl)propylhydroxyphosphinyl]amino]pentanedioic
acid;

15 2-[[[(1-naphthyl)butylhydroxyphosphinyl]amino]pentanedioic
acid;

2-[[[(2-naphthyl)butylhydroxyphosphinyl]amino]pentanedioic
acid; and

2-[[[(phenylprop-2-enyl)hydroxyphosphinyl]amino]pentanedioic
20 acid.

Especially preferred compounds of Formula IV of the
present invention wherein R_1 is a straight or branched
aliphatic group or a carbocyclic group, R_2 is ethyl which is
25 substituted with a carboxylic acid, and the NR_1 is amino are
selected from the group consisting of:

2-[(benzylhydroxyphosphinyl)amino]pentanedioic acid;

2-[(phenylhydroxyphosphinyl)amino]pentanedioic acid;

2-[[[(hydroxy)phenylmethyl]hydroxyphosphinyl]amino]
pentanedioic acid;

2-[(butylhydroxyphosphinyl)amino]pentanedioic acid;

2-[[[(3-methylbenzyl)hydroxyphosphinyl]amino]pentanedioic acid;

5 2-[(3-phenylpropylhydroxyphosphinyl)amino]pentanedioic acid;

2-[[[(4-fluorophenyl)hydroxyphosphinyl]amino]pentanedioic acid;

2-[(methylhydroxyphosphinyl)amino]pentanedioic acid;

2-[(phenylethylhydroxyphosphinyl)amino]pentanedioic acid;

2-[[[(4-methylbenzyl)hydroxyphosphinyl]amino]pentanedioic acid;

10 2-[[[(4-fluorobenzyl)hydroxyphosphinyl]amino]pentanedioic acid;

2-[[[(4-methoxybenzyl)hydroxyphosphinyl]amino]pentanedioic
acid;

2-[[[(2-fluorobenzyl)hydroxyphosphinyl]amino]pentanedioic acid;

15 2-[[[(pentafluorobenzyl)hydroxyphosphinyl]amino]pentanedioic
acid;

2-[(phosphono)amino]pentanedioic acid; and

2-[[[3-

trifluoromethylbenzyl)hydroxyphosphinyl]amino]pentanedioic
acid.

20

Compounds of the present invention wherein R_1 is a
straight or branched aliphatic group or a carbocyclic group, R_2
is phenyl, and X is amino are selected from the group
consisting of:

25 2-[[methylhydroxyphosphinyl]amino]-2-phenylethanoic acid;

2-[[ethylhydroxyphosphinyl]amino]-2-phenylethanoic acid;

2-[[propylhydroxyphosphinyl]amino]-2-phenylethanoic acid;

2-[[butylhydroxyphosphinyl]amino]-2-phenylethanoic acid;

2-[[cyclohexylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
2-[[cyclohexylmethylhydroxyphosphinyl]amino]-2-phenyl
ethanoic acid;

2-[[phenylhydroxyphosphinyl]amino]-2-phenylethanoic acid;

5 2-[[benzylhydroxyphosphinyl]amino]-2-phenylethanoic acid;

2-[[phenylethylhydroxyphosphinyl]amino]-2-phenylethanoic acid;

2-[[phenylpropylhydroxyphosphinyl]amino]-2-phenylethanoic
acid;

2-[[phenylbutylhydroxyphosphinyl]amino]-2-phenylethanoic acid;

10 2-[[2, 3, 4-trimethoxyphenyl]-3-hydroxyphosphinyl]amino]-2-
phenylethanoic acid;

2-[[1-naphthyl]hydroxyphosphinyl]amino]-2-phenylethanoic
acid;

15 2-[[2-naphthyl]hydroxyphosphinyl]amino]-2-phenylethanoic
acid;

2-[[1-naphthyl]methylhydroxyphosphinyl]amino]-2-phenyl
ethanoic acid;

2-[[2-naphthyl]methylhydroxyphosphinyl]amino]-2-phenyl
ethanoic acid;

20 2-[[1-naphthyl]ethylhydroxyphosphinyl]amino]-2-phenylethanoic
acid;

2-[[2-naphthyl]ethylhydroxyphosphinyl]amino]-2-phenylethanoic
acid;

25 2-[[1-naphthyl]propylhydroxyphosphinyl]amino]-2-phenyl
ethanoic acid;

2-[[2-naphthyl]propylhydroxyphosphinyl]amino]-2-phenyl
ethanoic acid;

2-[[1-naphthyl]butylhydroxyphosphinyl]amino]-2-phenylethanoic

acid;

2-[[(2-naphthyl)butylhydroxyphosphinyl]amino]-2-phenylethanoic acid; and

2-[[phenylprop-2-enylhydroxyphosphinyl]amino]-2-phenylethanoic acid.

Although not limited to any one particular species, a highly preferred phosphoramidate species where R_1 is aliphatic or carbocyclic, R_2 is ethyl which is substituted with carboxylic acid, and X is amino is 2-[[benzylhydroxyphosphinyl]amino]pentanedioic acid.

Other especially preferred compounds are selected from the group consisting of:

phosphoramidate derivatives wherein X is amino, R_1 is a straight or branched aliphatic group or a carbocyclic group, and R_2 is an C_2 - C_8 alkyl or alkenyl chain which is substituted with a carboxylic acid. Exemplary species include:

2-[(methylhydroxyphosphinyl)amino]hexanedioic acid;

2-[(benzylhydroxyphosphinyl)amino]hexanedioic acid;

2-[(methylhydroxyphosphinyl)amino]heptanedioic acid;

2-[(benzylhydroxyphosphinyl)amino]heptanedioic acid;

2-[(methylhydroxyphosphinyl)amino]octanedioic acid;

2-[(benzylhydroxyphosphinyl)amino]octanedioic acid;

2-[(methylhydroxyphosphinyl)amino]nonanedioic acid;

2-[(benzylhydroxyphosphinyl)amino]nonanedioic acid;

2-[(methylhydroxyphosphinyl)amino]decanedioic acid; and

2-[(benzylhydroxyphosphinyl)amino]decanedioic acid.

Other especially preferred compounds are selected from the group consisting of:

phosphoramidate derivatives wherein X is amino, R₁ is benzyl, and R₂ is a straight or branched aliphatic group or a carbocyclic group. Exemplary species include:

2-[[benzylhydroxyphosphinyl]amino]-2-methylethanoic acid;

2-[[benzylhydroxyphosphinyl]amino]-2-ethylethanoic acid;

2-[[benzylhydroxyphosphinyl]amino]-2-propylethanoic acid;

2-[[benzylhydroxyphosphinyl]amino]-2-butylethanoic acid;

2-[[benzylhydroxyphosphinyl]amino]-2-cyclohexylethanoic acid;

2-[[benzylhydroxyphosphinyl]amino]-2-(cyclohexyl)methyl ethanoic acid;

2-[[benzylhydroxyphosphinyl]amino]-2-phenylethanoic acid;

2-[[benzylhydroxyphosphinyl]amino]-2-benzylethanoic acid;

2-[[benzylhydroxyphosphinyl]amino]-2-phenylethylethanoic acid;

2-[[benzylhydroxyphosphinyl]amino]-2-phenylpropylethanoic acid;

2-[[benzylhydroxyphosphinyl]amino]-2-phenylbutylethanoic acid;

2-[[benzylhydroxyphosphinyl]amino]-2-(2, 3, 4-

trimethoxyphenyl)ethanoic acid;

2-[[benzylhydroxyphosphinyl]amino]-2-(1-naphthyl)ethanoic acid;

2-[[benzylhydroxyphosphinyl]amino]-2-(2-naphthyl)ethanoic acid;

2-[[benzylhydroxyphosphinyl]amino]-2-(1-naphthyl)methyl ethanoic acid;

2-[[benzylhydroxyphosphinyl]amino]-2-(2-naphthyl)methyl ethanoic acid;

2-[[benzylhydroxyphosphinyl]amino]-2-(1-naphthyl)ethylethanoic acid;

2-[[benzylhydroxyphosphinyl]amino]-2-(2-naphthyl)ethylethanoic acid;

5 2-[[benzylhydroxyphosphinyl]amino]-2-(1-naphthyl)propyl ethanoic acid;

2-[[benzylhydroxyphosphinyl]amino]-2-(2-naphthyl)propyl ethanoic acid;

10 2-[[benzylhydroxyphosphinyl]amino]-2-(1-naphthyl)butylethanoic acid;

2-[[benzylhydroxyphosphinyl]amino]-2-(2-naphthyl)butylethanoic acid; and

2-[[benzylhydroxyphosphinyl]amino]-2-phenylprop-2-enylethanoic acid.

15

Especially preferred methods utilize compounds wherein R₁ is said alkyl, alkenyl, cycloalkyl, or aryl group which is substituted with a heterocyclic group, R₂ is ethyl which is substituted with carboxylic acid, and X is amino are selected from the group consisting of:

20 2-[[[(2-pyridyl)methylhydroxyphosphinyl]amino]pentanedioic acid;

2-[[[(3-pyridyl)methylhydroxyphosphinyl]amino]pentanedioic acid;

25 2-[[[(4-pyridyl)methylhydroxyphosphinyl]amino]pentanedioic acid;

2-[[[(3-pyridyl)ethylhydroxyphosphinyl]amino]pentanedioic acid;

2-[[[(3-pyridyl)propylhydroxyphosphinyl]amino]pentanedioic

acid;

2-[[[(tetrahydrofuranyl)methylhydroxyphosphinyl]amino]
pentanedioic acid;

2-[[[(tetrahydrofuranyl)ethylhydroxyphosphinyl]amino]
5 pentanedioic acid;

2-[[[(tetrahydrofuranyl)propylhydroxyphosphinyl]amino]
pentanedioic acid;

2-[[[(2-indolyl)methylhydroxyphosphinyl]amino]pentanedioic
acid;

10 2-[[[(3-indolyl)methylhydroxyphosphinyl]amino]pentanedioic
acid;

2-[[[(4-indolyl)methylhydroxyphosphinyl]amino]pentanedioic
acid;

2-[[[(3-indolyl)ethylhydroxyphosphinyl]amino]pentanedioic acid;

15 2-[[[(3-indolyl)propylhydroxyphosphinyl]amino]pentanedioic
acid;

2-[[[(2-thienyl)methylhydroxyphosphinyl]amino]pentanedioic
acid;

2-[[[(3-thienyl)methylhydroxyphosphinyl]amino]pentanedioic
20 acid;

2-[[[(4-thienyl)methylhydroxyphosphinyl]amino]pentanedioic
acid;

2-[[[(3-thienyl)ethylhydroxyphosphinyl]amino]pentanedioic acid;
and

25 2-[[[(3-thienyl)propylhydroxyphosphinyl]amino]pentanedioic
acid.

Especially preferred methods utilize compounds wherein R₁

is said alkyl, alkenyl, cycloalkyl, or aryl group which is substituted with a heterocyclic group, R₂ is phenyl, and X is amino are selected from the group consisting of:

- 5 2-[[[(2-pyridyl)methylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
- 2-[[[(3-pyridyl)methylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
- 2-[[[(4-pyridyl)methylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
- 10 2-[[[(3-pyridyl)ethylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
- 2-[[[(3-pyridyl)propylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
- 15 2-[[[(tetrahydrofuranyl)methylhydroxyphosphinyl]amino]-2-phenyl ethanoic acid;
- 2-[[[(tetrahydrofuranyl)ethylhydroxyphosphinyl]amino]-2-phenyl ethanoic acid;
- 2-[[[(tetrahydrofuranyl)propylhydroxyphosphinyl]amino]-2-phenyl ethanoic acid;
- 20 2-[[[(2-indolyl)methylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
- 2-[[[(3-indolyl)methylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
- 25 2-[[[(4-indolyl)methylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
- 2-[[[(3-indolyl)ethylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
- 2-[[[(3-indolyl)propylhydroxyphosphinyl]amino]-2-phenylethanoic

acid;

2-[[(2-thienyl)methylhydroxyphosphinyl]amino]-2-phenylethanoic acid;

2-[[(3-thienyl)methylhydroxyphosphinyl]amino]-2-phenylethanoic acid;

2-[[(4-thienyl)methylhydroxyphosphinyl]amino]-2-phenylethanoic acid;

2-[[(3-thienyl)ethylhydroxyphosphinyl]amino]-2-phenylethanoic acid; and

2-[[(3-thienyl)propylhydroxyphosphinyl]amino]-2-phenylethanoic acid.

Especially preferred methods utilize compounds wherein R_1 is benzyl, R_2 is said alkyl, alkenyl, cycloalkyl, or aryl group which is substituted with a heterocyclic group, and X is amino are selected from the group consisting of:

2-[[benzylhydroxyphosphinyl]amino]-2-(2-pyridyl)methylethanoic acid;

2-[[benzylhydroxyphosphinyl]amino]-2-(3-pyridyl)methylethanoic acid;

2-[[benzylhydroxyphosphinyl]amino]-2-(4-pyridyl)methylethanoic acid;

2-[[benzylhydroxyphosphinyl]amino]-2-(3-pyridyl)ethylethanoic acid;

2-[[benzylhydroxyphosphinyl]amino]-2-(3-pyridyl)propylethanoic acid;

2-[[benzylhydroxyphosphinyl]amino]-2-(tetrahydrofuranyl)methylethanoic acid;

2-[[benzylhydroxyphosphinyl]amino]-2-(tetrahydrofuranyl)ethyl
ethanoic acid;

2-[[benzylhydroxyphosphinyl]amino]-2-(tetrahydrofuranyl)
propylethanoic acid;

5 2-[[benzylhydroxyphosphinyl]amino]-2-(2-indolyl)methylethanoic
acid;

2-[[benzylhydroxyphosphinyl]amino]-2-(3-indolyl)methylethanoic
acid;

10 2-[[benzylhydroxyphosphinyl]amino]-2-(4-indolyl)methylethanoic
acid;

2-[[benzylhydroxyphosphinyl]amino]-2-(3-indolyl)ethylethanoic
acid;

2-[[benzylhydroxyphosphinyl]amino]-2-(3-indolyl)propylethanoic
acid;

15 2-[[benzylhydroxyphosphinyl]amino]-2-(2-thienyl)methylethanoic
acid;

2-[[benzylhydroxyphosphinyl]amino]-2-(3-thienyl)methylethanoic
acid;


20 2-[[benzylhydroxyphosphinyl]amino]-2-(4-thienyl)methylethanoic
acid;

2-[[benzylhydroxyphosphinyl]amino]-2-(3-thienyl)ethylethanoic
acid; and

2-[[benzylhydroxyphosphinyl]amino]-2-(3-thienyl)propylethanoic
acid.

25

In another preferred embodiment, R₁ is a heterocyclic
substituent illustrated by the compounds selected from the
group of formula IV:



wherein

IV

R_1 is Ar_1 ; and

R_2 is C_1 - C_8 straight or branched chain alkyl, C_2 - C_8 straight or branched chain alkenyl group, C_3 - C_8 cycloalkyl, C_5 - C_8 cycloalkenyl, or Ar_1 , wherein said alkyl, alkenyl, cycloalkyl, cycloalkenyl or aryl group may be optionally substituted with carboxylic acid.

Especially preferred compounds wherein R_1 is a heterocyclic group, R_2 is ethyl which is substituted with carboxylic acid, and X is amino are selected from the group consisting of:

2-[[(2-pyridyl)hydroxyphosphinyl]amino]pentanedioic acid;
 2-[[(3-pyridyl)hydroxyphosphinyl]amino]pentanedioic acid;
 2-[[(4-pyridyl)hydroxyphosphinyl]amino]pentanedioic acid;
 2-[[(tetrahydrofuranyl)hydroxyphosphinyl]amino]pentanedioic acid;


2-[[(2-indolyl)hydroxyphosphinyl]amino]pentanedioic acid;
 2-[[(3-indolyl)hydroxyphosphinyl]amino]pentanedioic acid;
 2-[[(4-indolyl)hydroxyphosphinyl]amino]pentanedioic acid;
 2-[[(2-thienyl)hydroxyphosphinyl]amino]pentanedioic acid;
 2-[[(3-thienyl)hydroxyphosphinyl]amino]pentanedioic acid; and

2-[[[(4-thienyl)hydroxyphosphinyl]amino]pentanedioic acid.

Compounds of the present invention wherein R₁ is a heterocyclic group, R₂ is phenyl, and X is amino are selected from the group consisting of:

2-[[[(2-pyridyl)hydroxyphosphinyl]amino]-2-phenylethanoic acid;
 2-[[[(3-pyridyl)hydroxyphosphinyl]amino]-2-phenylethanoic acid;
 2-[[[(4-pyridyl)hydroxyphosphinyl]amino]-2-phenylethanoic acid;
 2-[[[(tetrahydrofuran-2-yl)hydroxyphosphinyl]amino]-2-phenylethanoic acid;
 2-[[[(2-indolyl)hydroxyphosphinyl]amino]-2-phenylethanoic acid;
 2-[[[(3-indolyl)hydroxyphosphinyl]amino]-2-phenylethanoic acid;
 2-[[[(4-indolyl)hydroxyphosphinyl]amino]-2-phenylethanoic acid;
 2-[[[(2-thienyl)hydroxyphosphinyl]amino]-2-phenylethanoic acid;
 2-[[[(3-thienyl)hydroxyphosphinyl]amino]-2-phenylethanoic acid;
 and
 2-[[[(4-thienyl)hydroxyphosphinyl]amino]-2-phenylethanoic acid.

Compounds are also preferably selected from the group of formula IV:



wherein

IV

R₁ is hydrogen, C₁-C₈ straight or branched chain alkyl, C₂-C₈ straight or branched chain alkenyl group, C₁-C₈

cycloalkyl, C₅-C₇ cycloalkenyl, or Ar₁; and

R₂ is Ar₁, wherein said aryl group may be optionally substituted with carboxylic acid.

5 Particular species wherein R₂ is heterocyclic may be easily made and used by persons of ordinary skill in the art in accordance with the teachings provided herein and known in the art.

Compounds of the present invention wherein R₁ is benzyl, R₂ is heterocyclic, and X is amino are selected from the group consisting of:

2-[[benzylhydroxyphosphinyl]amino]-2-(2-pyridyl)ethanoic acid;

2-[[benzylhydroxyphosphinyl]amino]-2-(3-pyridyl)ethanoic acid;

2-[[benzylhydroxyphosphinyl]amino]-2-(4-pyridyl)ethanoic acid;

15 2-[[benzylhydroxyphosphinyl]amino]-2-(tetrahydrofuranyl)ethanoic acid;

2-[[benzylhydroxyphosphinyl]amino]-2-(2-indolyl)ethanoic acid;

2-[[benzylhydroxyphosphinyl]amino]-2-(3-indolyl)ethanoic acid;

2-[[benzylhydroxyphosphinyl]amino]-2-(4-indolyl)ethanoic acid;

20 2-[[benzylhydroxyphosphinyl]amino]-2-(2-thienyl)ethanoic acid;

2-[[benzylhydroxyphosphinyl]amino]-2-(3-thienyl)ethanoic acid;

and

2-[[benzylhydroxyphosphinyl]amino]-2-(4-thienyl)ethanoic acid.


25

Synthesis of Compounds

The compounds of the present invention can be readily prepared by standard techniques of organic chemistry, utilizing the general synthetic pathways depicted below (see

Schemes I-IX). Precursor compounds may be prepared by methods known in the art, such as those described in the method of Jackson et al. (*J. Med. Chem.* 39(2), 619-622, Design, Synthesis, and Biological Activity of a Potent Inhibitor of the Neuropeptidase N-Acetylated α -Linked Acidic Dipeptidase) and, for example, in Froestl et al. (*J. Med. Chem.*, 1995, 38, 3313-3331, Phosphinic Acid Analogues of GABA).



SCHEME I



Production of compounds containing the R group substitutions can be easily made utilizing known methods. Further methods of synthesizing phosphinic acid esters are also described in *J. Med. Chem.*, 1988, 31, 204-212, and may be found in Scheme II, below.


SCHEME II

Method A



- | | | | | |
|----|----|--|----|---|
| 5 | A. | R' = (CH ₂) ₃ Ph | H. | n-C ₇ H ₁₅ |
| | B. | (CH ₂) ₄ Ph | I. | n-C ₉ H ₁₇ |
| | C. | (CH ₂) ₅ Ph | J. | n-C ₉ H ₁₉ |
| | D. | (CH ₂) ₄ (P-F-Ph) | K. | n-C ₁₀ H ₂₁ |
| | E. | (CH ₂) ₄ -(3-pyridyl) | L. | CH ₂ (CH)(CH ₃)C ₄ H ₉ |
| 10 | F. | n-C ₅ H ₁₁ | M. | CH ₂ (CH ₃)CH(CH ₃) ₂ |
| | G. | n-C ₅ H ₁₃ | | |

Method B




- 15
- N. R' = n-C₄H₉
 O. CH(CH₃)C₅H₁₁

20

Starting with the aforementioned phosphinic acid esters, there are a variety of routes that can be used to prepare the compounds of the present invention. For example, a general route was recently described in *J. Med. Chem.*, 1996, 39, 619-

622, and is set forth below in Scheme III.




SCHEME III


5

Another route for preparing the compounds of the present invention is as set forth below in scheme IV and Scheme V. Scheme IV and Scheme V also show a phosphinic acid derivative as a starting material to prepare the compounds of the present invention and the R group is contemplated as including any reasonable chemical substitution and includes without limitation the R groups listed in Scheme II and within the specification.

10




SCHEME IV




SCHEME V

Another route of preparing the compounds of the present invention allows for aromatic substitution at R1 and is set forth below in Scheme VI.




SCHEME VI

Another route of preparing the compounds of the present invention allows for aromatic substitution at the R2 position and is set forth below in Scheme VII.




SCHEME VII

Preparation of the compounds of the present invention wherein X is NR1 is set forth below in Scheme VIII.



SCHEME VIII

Preparation of the compounds of the present invention wherein X is oxygen is set forth below in Scheme IX.



SCHEME IX

Pharmaceutical Compositions of the Present Invention

The present invention also relates to a pharmaceutical composition comprising:

- (i) a therapeutically effective amount of a compound of formulas I, II, III or IV; and
- (ii) a pharmaceutically acceptable carrier.

In another preferred embodiment, the pharmaceutical composition further comprises a therapeutic agent selected from the group consisting of therapeutic hormones, chemotherapeutic agents, monoclonal antibodies, anti-angiogenesis agents, radiolabelled compounds, antineoplastic

agents and mixtures thereof. Examples of therapeutic hormones include diethylstilbestrol (DES), leuprolide, flutamide, cyproterone acetate, ketoconazole and amino glutethimide are preferred. Examples of antineoplastic agents include 5-fluorouracil, vinblastine sulfate, estramustine phosphate, suramin and strontium-89. Examples of chemotherapeutic agents include buserelin, chlorotranisene, chromic phosphate, cisplatin, cyclophosphamide, dexamethasone, doxorubicin, estradiol, estradiol valerate, estrogens conjugated and esterified, estrone, ethinyl estradiol, floxuridine, goserelin, hydroxyurea, melphalan, methotrexate, mitomycin and prednisone.

In a further preferred embodiment, the compound of formula I, II, III or IV is present in an amount that is effective for inhibiting NAALADase activity in an animal or treating a prostate disease in an animal.

Process for Preparing Pharmaceutical Compositions

In yet another preferred embodiment, a process for preparing a pharmaceutical composition or medicament containing a compound of the present invention for treating a disease is also contemplated.

Methods of Use of the Present Invention

i) Method of inhibiting NAALADase enzyme activity

The present invention further relates to a method of inhibiting NAALADase enzyme activity in an animal, comprising

administering an effective amount of a compound of formula I, II, III or IV to said animal.

ii) Method of treating a prostate disease

5 The present invention also relates to a method of treating a prostate disease in an animal, comprising administering an effective amount of a compound of formula I, II, III or IV to said animal.

10 In a preferred embodiment, said prostate disease is prostate cancer such as prostatic adenocarcinoma, benign prostatic hyperplasia, or conditions involving the prostate requiring administration of the compounds of the present invention, such as prostatic intraepithelial neoplasia (PIN).

15 iii) Method of treating cancer

20 In addition to prostate cancer, other forms of cancer that may be treated with the compounds of the present invention include without limitation: ACTH-producing tumors, acute lymphocytic leukemia, acute nonlymphocytic leukemia, cancer of the adrenal cortex, bladder cancer, brain cancer, breast cancer, cervix cancer, chronic lymphocytic leukemia, chronic myelocytic leukemia, colorectal cancer, cutaneous T-cell lymphoma, endometrial cancer, esophageal cancer, Ewing's sarcoma, gallbladder cancer, hairy cell leukemia, head & neck
25 cancer, Hodgkin's lymphoma, Kaposi's sarcoma, kidney cancer, liver cancer, lung cancer (small and/or non-small cell), malignant peritoneal effusion, malignant pleural effusion, melanoma, mesothelioma, multiple myeloma, neuroblastoma, non-

Hodgkin's lymphoma, osteosarcoma, ovary cancer, ovary (germ cell) cancer, pancreatic cancer, penis cancer, retinoblastoma, skin cancer, soft-tissue sarcoma, squamous cell carcinomas, stomach cancer, testicular cancer, thyroid cancer, trophoblastic neoplasms, cancer of the uterus, vaginal cancer, cancer of the vulva and Wilm's tumor.

The compounds of the present invention are particularly useful in treating cancer of tissues where NAALADase enzymes reside. Such tissues include the prostate as well as the brain, kidney and testis.

For patients who initially present without advanced or metastatic cancer, NAALADase inhibitor based drugs are used as an immediate initial therapy prior to surgery and radiation therapy, and as a continuous post-treatment therapy in patients at risk for recurrence or metastasis (based upon high PSA, high Gleason's score, locally extensive disease, and/or pathological evidence of tumor invasion in the surgical specimen). The goal in these patients is to inhibit the growth of potentially metastatic cells from the primary tumor during surgery or radiotherapy and inhibit the growth of tumor cells from undetectable residual primary tumor.

For patients who initially present with advanced or metastatic cancer, NAALADase inhibitor based drugs are used as a continuous supplement to, or possible as a replacement for hormonal ablation. The goal in these patients is to slow tumor cell growth from both the untreated primary tumor and from the existing metastatic lesions.

In addition, the invention may be particularly

efficacious during post-surgical recovery, where the present compositions and methods may be particularly effective in lessening the chances of recurrence of a tumor engendered by shed cells that cannot be removed by surgical intervention.

5

iv) Diagnostic kits

The present invention also includes a diagnostic kit for performing the methods of the present invention and may contain compounds and/or compositions containing the compounds of the present invention. Radiolabelled compounds and monoclonal antibodies may be used in a manner so as to provide diagnostic information. Examples of diagnostic information and uses include determining the type of disease, the progress of the particular disease, the location of cells targeted by a NAALADase inhibitor, radiolabelled compound or monoclonal antibody, and similar diagnostic uses known to persons skilled in the art.

15

Route of Administration

20

In the methods of the present invention, the compounds may be administered orally, parenterally, by inhalation spray, topically, rectally, nasally, buccally, vaginally or via an implanted reservoir in dosage formulations containing conventional non-toxic pharmaceutically-acceptable carriers, adjuvants and vehicles. The term parenteral as used herein includes subcutaneous, intravenous, intramuscular, intraperitoneal, intrathecal, intraventricular, intrasternal or intracranial injection and infusion techniques. Invasive

25

techniques are preferred, particularly direct administration to damaged neuronal tissue.

To be effective therapeutically as central nervous system targets, the compounds of the present invention should readily
5 penetrate the blood-brain barrier when peripherally administered. Compounds which cannot penetrate the blood-brain barrier can be effectively administered by an intraventricular route.

The compounds may also be administered in the form of
10 sterile injectable preparations, for example, as sterile injectable aqueous or oleaginous suspensions. These suspensions can be formulated according to techniques known in the art using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparations may
15 also be sterile injectable solutions or suspensions in non-toxic parenterally-acceptable diluents or solvents, for example, as solutions in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution. In addition,
20 sterile fixed oils are conventionally employed as solvents or suspending mediums. For this purpose, any bland fixed oil such as a synthetic mono- or di-glyceride may be employed. Fatty acids such as oleic acid and its glyceride derivatives, including olive oil and castor oil, especially in their
25 polyoxyethylated forms, are useful in the preparation of injectables. These oil solutions or suspensions may also contain long-chain alcohol diluents or dispersants.

Additionally, the compounds may be administered orally in

the form of capsules, tablets, aqueous suspensions or solutions. Tablets may contain carriers such as lactose and corn starch, and/or lubricating agents such as magnesium stearate. Capsules may contain diluents including lactose and
5 dried corn starch. Aqueous suspensions may contain emulsifying and suspending agents combined with the active ingredient. The oral dosage forms may further contain sweetening and/or flavoring and/or coloring agents.

The compounds may further be administered rectally in the
10 form of suppositories. These compositions can be prepared by mixing the drug with suitable non-irritating excipients which are solid at room temperature, but liquid at rectal temperature such that they will melt in the rectum to release the drug. Such excipients include cocoa butter, beeswax and
15 polyethylene glycols.

Moreover, the compounds may be administered topically, especially when the conditions addressed for treatment involve areas or organs readily accessible by topical application, including neurological disorders of the eye, the skin or the
20 lower intestinal tract.

For topical application to the eye, or ophthalmic use, the compounds can be formulated as micronized suspensions in isotonic, pH adjusted sterile saline or, preferably, as a solution in isotonic, pH adjusted sterile saline, either with
25 or without a preservative such as benzylalkonium chloride. Alternatively, the compounds may be formulated into ointments, such as petrolatum.

For topical application to the skin, the compounds can be

formulated into suitable ointments containing the compounds suspended or dissolved in, for example, mixtures with one or more of the following: mineral oil, liquid petrolatum, white petrolatum, propylene glycol, polyoxyethylene polyoxypropylene compound, emulsifying wax and water. Alternatively, the compounds can be formulated into suitable lotions or creams containing the active compound suspended or dissolved in, for example, a mixture of one or more of the following: mineral oil, sorbitan monostearate, polysorbate 60, cetyl ester wax, cetearyl alcohol, 2-octyldodecanol, benzyl alcohol and water.

Topical application to the lower intestinal tract can be effected in rectal suppository formulations (see above) or in suitable enema formulations.

The compounds of the present invention may be administered by a single dose, multiple discrete doses or continuous infusion. Since the compounds are small, easily diffusible and relatively stable, they are well suited to continuous infusion. Pump means, particularly subcutaneous pump means, are preferred for continuous infusion.

Compositions and methods of the invention also may utilize controlled release technology. Thus, for example, NAALADase inhibitors may be incorporated into a polymer matrix for controlled release over a period of days. Such controlled release films are well known to the art. Examples of polymers commonly employed for this purpose that may be used in the present invention include nondegradable ethylene-vinyl acetate copolymer and degradable lactic acid-glycolic acid copolymers. Certain hydrogels such as poly(hydroxyethylmethacrylate) or

poly(vinylalcohol) also may be useful.

Dosage

Dose levels on the order of about 0.1 mg to about 10,000 mg of the active ingredient compound are useful in the treatment of the above conditions, with preferred levels being about 0.1 mg to about 1,000 mg. The specific dose level for any particular patient will vary depending upon a variety of factors, including the activity of the specific compound employed; the age, body weight, general health, sex and diet of the patient; the time of administration; the rate of excretion; drug combination; the severity of the particular disease being treated; and the form of administration. Typically, in vitro dosage-effect results provide useful guidance on the proper doses for patient administration. Studies in animal models are also helpful, particularly in determining effective doses for treating cancer. The considerations for determining the proper dose levels are well known in the art.

In a preferred embodiment, the compounds of the present invention are administered in lyophilized form. In this case, 1 to 100 mg of a compound of the present invention may be lyophilized in individual vials, together with a carrier and a buffer, such as mannitol and sodium phosphate. The compound may be reconstituted in the vials with bacteriostatic water before administration.

As previously mentioned, the compounds of the present invention may be administered in combination with one or more

therapeutic agents, including chemotherapeutic agents. TABLE I provides known median dosages for selected chemotherapeutic agents. Specific dose levels for these agents will depend upon considerations such as those identified above for the

5 compounds of the present invention.

TABLE I

CHEMOTHERAPEUTIC AGENT	MEDIAN DOSAGE
Asparaginase	10,000 units
Bleomycin Sulfate	15 units
5 Carboplatin	50-450 mg
Carmustine	100 mg
Cisplatin	10-50 mg
Cladribine	10 mg
10 Cyclophosphamide (lyophilized)	100 mg-2 gm
Cyclophosphamide (non-lyophilized)	100 mg-2 gm
Cytarabine (lyophilized powder)	100 mg-2 gm
15 Dacarbazine	100 mg-200 mg
Dactinomycin	0.5 mg
Daunorubicin	20 mg
Diethylstilbestrol	250 mg
Doxorubicin	10-150 mg
20 Etidronate	300 mg
Etoposide	100 mg
Floxuridine	500 mg
Fludarabine Phosphate	50 mg
Fluorouracil	500 mg-5 gm
25 Goserelin	3.6 mg
Granisetron Hydrochloride	1 mg
Idarubicin	5-10 mg
Ifosfamide	1-3 gm
Leucovorin Calcium	50-350 mg
30 Leuprolide	3.75-7.5 mg
Mechlorethamine	10 mg
Medroxyprogesterone	1 gm
Melphalan	50 gm
Methotrexate	20 mg-1 gm

CHEMOTHERAPEUTIC AGENT	MEDIAN DOSAGE
Mitomycin	5-40 mg
Mitoxantrone	20-30 mg
Ondansetron Hydrochloride	40 mg
Paclitaxel	30 mg
Pamidronate Disodium	30-90 mg
Pegaspargase	750 units
Plicamycin	2,500 mcgm
Streptozocin	1 gm
Thiotepa	15 mg
Teniposide	50 mg
Vinblastine	10 mg
Vincristine	1-5 mg
Aldesleukin	22 million units
Epoetin Alpha	2,000-10,000 units
Filgrastim	300-480 mcgm
Immune Globulin	500 mg-10 gm
Interferon Alpha-2a	3-36 million units
Interferon Alpha-2b	3-50 million units
Levamisole	50 mg
Octreotide	1,000-5,000 mcgm
Sargramostim	250-500 mcgm

Administration regimen

For the methods of the present invention, any administration regimen regulating the timing and sequence of drug delivery can be used and repeated as necessary to effect treatment. Such regimen may include pretreatment and/or co-administration with additional therapeutic agents.

For patients with prostate cancer that is neither advanced nor metastatic, the compounds of the present invention may be administered (i) prior to surgery or radiation treatment to reduce the risk of metastasis; (ii) during surgery or in conjunction with radiation treatment; and/or (iii) after surgery or radiation therapy to reduce the risk of recurrence and to inhibit the growth of any residual tumorous cells.

For patients with advanced or metastatic prostate cancer, the compounds of the present invention may be administered as a continuous supplement to, or as a replacement for, hormonal ablation in order to slow tumor cell growth in both the untreated primary tumor and the existing metastatic lesions.

The methods of the present invention are particularly useful where shed cells could not be removed by surgical intervention. After post-surgical recovery, the methods of the present invention would be effective in reducing the chances of recurrence of a tumor engendered by such shed cells.

Combination with Other Treatments

(i) Surgery and Radiation Treatment

In general, surgery and radiation treatment are employed as potentially curative therapies for patients with localized prostate cancer who are under 70 years of age and are expected to live at least 10 more years.

Approximately 70% of newly diagnosed prostate cancer patients fall into this category. Approximately 90% of these

patients (65% of total patients) undergo surgery, while approximately 10% of these patients (7% of total patients) undergo radiation treatment.

Histopathological examination of surgical specimens
5 reveals that approximately 63% of patients undergoing surgery (40% of total patients) have locally extensive tumors or regional (lymph node) metastasis that was undetected at initial diagnosis. These patients are at a significantly greater risk of recurrence. Approximately 40% of these
10 patients will actually develop recurrence within five years after surgery. Results after radiation treatment are even less encouraging. Approximately 80% of patients who have undergone radiation treatment as their primary therapy have disease persistence or develop recurrence or metastasis within
15 five years after treatment.

Currently, most prostate cancer patients undergoing surgery and radiation treatment do not receive any immediate follow-up therapy. Rather, they are monitored frequently for elevated Prostate Specific Antigen ("PSA"), which is the
20 primary indicator of recurrence or metastasis.

Based on the above statistics, there is considerable opportunity to use the present invention in conjunction with surgery and/or radiation treatment.

(ii) Hormonal Therapy

25 Hormonal ablation is the most effective palliative treatment for the 10% of patients with metastatic prostate cancer. Hormonal ablation by medication and/or orchiectomy is used to block hormones that promote further growth and

metastasis of prostate cancer. With time, both the primary and metastatic tumors of virtually all of these patients become hormone-independent and resistant to therapy.

Approximately 50% of patients with metastatic cancer die within three years after initial diagnosis, and 75% of such patients die within five years after diagnosis. Continuous supplementation with the compounds of the present invention may be used to prevent or reverse this potentially metastasis-permissive state.

(iii) Chemotherapy

While chemotherapy has been successful in treating some forms of cancer, it has shown slight therapeutic value in treating prostate cancer where it is generally reserved as a last resort. Accordingly, the opportunity to treat prostate cancer by combining chemotherapy with the methods of the present invention will be rare. When combined, however, such treatments should be more effective than chemotherapy alone in controlling prostate cancer.

(iv) Immunotherapy

The compounds of the present invention may also be used in combination with monoclonal antibodies to treat prostate cancer. Such combined treatment is particularly effective for patients with pelvic lymph node involvement, of which only 34% survive after 5 years. An example of such monoclonal antibodies is cell membrane-specific anti-prostate antibody.

The present invention may also be used with immunotherapies based on polyclonal or monoclonal antibody-

derived reagents. Monoclonal antibody-derived reagents are preferred. These reagents are well known in the art, and include radiolabelled monoclonal antibodies such as monoclonal antibodies conjugated with strontium-89.

5 (v) Cryotherapy

The methods of the present invention may also be used in conjunction with cryotherapy for treatment of prostate cancer.

Experimental Studies

10 The following experimental studies of compounds of the present invention and of structurally related compounds provide strong evidence that the compounds of the present invention are non-toxic and are effective in inhibiting NAALADase activity, treating glutamate abnormalities and
15 treating prostate diseases.

In Vivo Toxicity of NAALADase Inhibitors

To examine the toxicological effect of NAALADase inhibition in vivo, a group of mice were injected with 2-
20 (phosphonomethyl)pentanedioic acid, a NAALADase inhibitor of high activity, in doses of 1, 5, 10, 30, 100, 300 and 500 mg/kg body weight. The mice were subsequently observed two times per day for 5 consecutive days. The survival rate at each dose level is provided in TABLE II below. The results
25 show that the NAALADase inhibitor is non-toxic to mice, suggesting that the compounds of the present invention would be similarly non-toxic to humans when administered at therapeutically effective amounts.

TABLE II

TOXICOLOGICAL EFFECTS OF NAALADASE INHIBITORS							
Dose (mg/kg)	1	5	10	30	100	300	500
Survival Rate After 5 days (%)	100	100	100	100	100	100	66.7

In Vitro Assay of NAALADase Activity

The following compounds were tested for *in vitro* inhibition of NAALADase activity. The results are provided in Tables III(a), III(b), and III(c) below.

TABLE III(a)

IN VITRO ACTIVITY OF NAALADASE INHIBITORS

Compound	K_i (nM)
2-(phosphonomethyl)pentanedioic acid	0.275 ± 0.08
2-(phosphonomethyl)succinic acid	700.00 ± 67.3
2-[[2-carboxyethyl)hydroxyphosphinyl]-methyl]pentanedioic acid)	1.89 ± 0.19

2-(phosphonomethyl)pentanedioic acid showed a high level of NAALADase inhibiting activity, with a K_i of 0.27 nM (Table III(a)). The activity of this compound is >1000 times more potent than that of previously described inhibitors. Since 2-(phosphonomethyl)pentanedioic acid is similar in structure to the compounds of the present invention, the results suggest that the compounds of the present invention would also be potent NAALADase inhibitors. By comparison, 2-(phosphonomethyl)succinic acid exhibits much lower NAALADase

inhibiting activity, suggesting that a glutamate analog attached to the phosphonic acid contributes to its NAALADase inhibiting activity. The results also show that 2-[[2-carboxyethyl)-hydroxyphosphinyl]methyl]pentanedioic acid, which has an additional carboxylic acid side chain similar to the aspartate residue found in NAAG, exhibits a lower NAALADase inhibiting activity than 2-(phosphonomethyl)-pentanedioic acid.

10










Table III (b)

Other compounds demonstrating inhibition of NAALADase activity are set forth below in Table III(b). Results of the nine compounds in Table III(b) shows the remarkable K_i activity of a variety of compounds of the present invention. These compounds show NAALADase inhibitory ability wherein R1 comprises an aliphatic group, a aliphatic which is substituted, an aromatic group, and aromatic which is substituted.

15

Table III(b)

In vitro Activity of NAALADase Inhibitors

COMPOUND	Ki(nM)	COMPOUND	Ki(nM)
	34		231
	36		532
	54		1100
	148		148
	190		






5

Further results, provided in Table III(c), show the remarkable Ki activity of the compounds of the present invention. These compounds show NAALADase inhibition wherein R1 comprises a substituted aliphatic (benzyl) which is further substituted.

10

Table III(c) *in vitro* Activity of NAALADase Inhibitors

5

<u>Compound</u>	<u>Ki Value (nM)</u>
	Ki = 68nM
	Ki = 70nM
	Ki = 90nM
	Ki = 175nM
	Ki = 38nM

Protocol for In Vitro Assay of NAALADase Activity

The amount of [³H]Glu liberated from [³H]NAAG in 50 mM Tris-Cl buffer was measured for 15 minutes at 37° C using 30-50 µg of synaptosomal protein. Substrate and product were
5 resolved by anion-exchange liquid chromatography. Duplicate assays were performed so that no more than 20% of the NAAG was digested, representing the linear range of peptidase activity. Quisqualate (100 µM) was included in parallel assay tubes to confirm the specificity of the measurements.

10

In Vitro Assay of NAALADase Inhibitors
on Cancer

Referring now to FIGS. 1 and 2, the effect of NAALADase inhibitors on cancer cell line were examined. LNCAP cells (a
15 prostate cancer cell line) were treated with quisqualate acid (in concentrations ranging from 10 nM to 1 µM) and 2-(phosphonomethyl)pentanedioic acid (in concentrations ranging from 100 pM to 10 nM). The 3H-thymidine measurement for each concentration of quisqualate acid and 2-
20 (phosphonomethyl)pentanedioic acid is also provided in TABLE IV below. FIGS. 1 and 2 present this data graphically and particularly illustrate the decrease in proliferation and thymidine uptake of cells treated with NAALADase inhibitors.

25

TABLE IV

<u>3H-Thymidine Incorporation (dpm/well)</u>			
<u>Dose</u>	<u>Quisqualic Acid</u>	<u>2-(phosphonomethyl)-pentanedioic acid</u>	
Control	4813 \pm 572	4299 \pm 887	
10 pM	--	3078 \pm 1006	
100 pM	--	2062 \pm 595	
1 nM	3668 \pm 866	1001 \pm 52	
10 nM	2137 \pm 764	664 \pm 366	
100 nM	1543 \pm 312	--	
1 μ M	1295 \pm 181	--	

The results show that LNCAP cell proliferation (as measured by the incorporation of 3H-thymidine) decreased significantly as the concentration of the NAALADase inhibitors increased, suggesting that the compounds of the present invention would be effective in treating cancer, particularly prostate cancer.

Protocol for In Vitro Cancer Assay

Cells in RPMI 1640 medium containing 10% Fetal Calf Serum (FCS) are plated in 24 well plates and allowed to adhere for 24 hours before addition of quisqualic acid (10^{-9} M to 10^{-6} M) or 2-(phosphonomethyl)pentanedioic acid (10^{-11} M to 10^{-8} M) for 7 days. On the 7th day, the cells are pulsed with 3H-thymidine for 4 hours, harvested and measured for radioactivity. Values represent means \pm SEM of 6 separate cell wells for each treatment. All experiments are performed

at least twice.

To control for non-specific cytostatic effects of quisqualate acid and 2-(phosphonomethyl)pentanedioic acid, the agents are simultaneously evaluated on a non-NAALADase
5 containing prostate cell line, DU145 (Carter et al., Proc. Natl. Acad. Sci. USA, (93) 749-753, 1996). If the treatments with quisqualate acid and 2-(phosphonomethyl)pentanedioic have no significant effect on cell growth, the NAALADase inhibiting activity of the agents are uniquely responsible for their
10 cytostatic effects on NAALADase containing prostate carcinoma cell lines.

Cell Lines and Tissue Culture

LNCAP cells are obtained from Dr. William Nelson at the Johns Hopkins School of Medicine in Baltimore, MD. DU145
15 cells are obtained from American Type Culture Collection (Rockville, MD). Cells are grown in RPMI-1640 media supplemented with 10% heat-inactivated fetal calf serum, 2 mM-glutamine, 100 units/ml penicillin, and 100 µg/ml streptomycin (Paragon) in a humidified incubator at 37°C in a 5% CO₂/95% O₂
20 atmosphere.

[3H] Thymidine Incorporation Assays

The cells are suspended at 1×10^5 cells/ml in RPMI-1640 media and seeded into 24-well plates at 500 µl per well. After 24 hours, various concentrations of quisqualic acid
25 (Sigma) or the potent NAALADase inhibitor 2-(phosphonomethyl)pentanedioic acid (synthesized according to the methods of Jackson et al., J Med Chem 39(2) 619-622) is added to the wells and the plates are returned to the

incubator. On days 3, 5 and 7, media and drug are refreshed. On the 8th day following seeding, each well is pulsed with 1 μ Ci 3 H-thymidine (New England Nuclear) for 4 hours. Media is then removed and the wells washed 2 times with phosphate buffered saline (pH=7.4). The contents of each well is subsequently solubilized with 250 μ l of 0.2 N NaOH and transferred to scintillation vials. 5 ml UltimaGold (Packard) scintillation cocktail is added and radioactivity is quantitated using a Beckman LS6001 scintillation counter.

The purity and/or identity of all synthetic compounds is ascertained by thin layer chromatography, High Pressure Liquid Chromatography (HPLC), mass spectrometry, and elemental analysis. Proton Nuclear Magnetic Resonance (NMR) spectra are obtained using a Bruker spectrometer. Chemical shifts are reported in parts per million relative to tetramethylsilane as internal standard. Analytical thin-layer chromatography (TLC) is conducted on prelayered silica gel GHLF plates (Analtech, Newark, DE). Visualization of the plates is accomplished by using UV light, phosphomolybdic acid-ethanol, and/or iodoplatinate charring. Flash chromatography is conducted on Kieselgel 60, 230-400 mesh (E. Merck, Darmstadt, West Germany). Solvents are either reagent or HPLC grade. Reactions are run at ambient temperature and under a nitrogen atmosphere unless otherwise noted. Solutions are evaporated under reduced pressure on a Buchi rotary evaporator.

In vivo LNCaP Tumor Xenograft Assay and Results

Referring now to FIGS. 3 and 4, LNCaP human prostate cancer

cells were injected subcutaneously into the right flank of male nude mice. 2-(phosphonomethyl)pentanedioic acid, a NAALADase inhibitor, was administered by daily intratumoral injection (0.25 μ g/day) beginning when the tumors reached a volume of approximately 50-70 mm³. An additional group was included using a silicon polymer containing 2-(phosphonomethyl)pentanedioic acid which released approximately 0.25 μ g/day of drug locally into the tumor. The 2-(phosphonomethyl)pentanedioic acid polymer was changed two times per week. Tumor volumes were monitored for 42 days after the beginning of treatment.

EXPERIMENTAL PROCEDURES

Cell Lines

LNCaP is a human prostate cancer cell line that was established in 1973 from a pleural effusion of a patient who had been treated with 5-FU, doxorubicin, methotrexate, and CTX in the 3 months before the cell line was initiated. This line is androgen receptor positive and has been used in screening anticancer drugs that are targeted as hormone antagonists. LNCaP was grown in RPMI with 1.5 g NaHCO₃/L, 10% fetal bovine serum (FBS), and 2 mM L-glutamine and was kept at 37°C in a humidified 5% CO₂/O₂ incubator. Antibiotics were not added to the medium.

Animal Tumor Model

NCr nude (nu/nu) male mice, age 4-5 weeks, were purchased from Taconic (Germantown, NY). The animals were housed four per cage in sterile filter-topped cages in a ventilated cage rack. Upon arrival, they were quarantined for four working days before

use. Temperature was maintained at $72 \pm 5^{\circ}\text{F}$ and relative humidity at 35-70%, and a 12-hr light/dark cycle is used. The mice were fed sterile, autoclavable, certified Purina rodent chow *ad libitum*. Drinking water was acidified and autoclaved, and the source water was recirculated, deionized, UV-treated, and 5- μm filtered.

After the animals were released from quarantine, the mice were injected subcutaneously in the right flank with 1×10^7 LNCaP cells in Matrigel™ (0.1-ml injection volume). Tumor dimensions and body weight were measured twice weekly. Vernier calipers were used to measure tumors in three planes, and tumor volume (V) was calculated as follows: $V = \pi(x \times y \times z)/6$, where x, y, and z were the tumor measurements minus skin thickness. At the end of the experiment, the mice were sacrificed by CO₂ inhalation followed by cervical dislocation.

Pharmaceuticals

2-(phosphonomethyl)pentanedioic acid was made up in water at a concentration of 2.5 mg/ml. Polymer containing 2-(phosphonomethyl)pentanedioic acid was made up by grinding 140 mg NaCl to a fine powder then mixing with 5mg 2-(phosphonomethyl)pentanedioic acid and 350 mg silicone gel. The mixture was spread to a thin film and allowed to dry for 24 hours. The material was cut into 1-1.5 mg pieces for subcutaneous implantation.

Treatment Protocol

When the tumor volumes reached a predetermined size (mean

tumor volume 50-70 mm³), mice were added randomly into treatment groups of six to eight mice each. All treatments were administered daily for at least 4 weeks. 2-(phosphonomethyl)pentanedioic acid was administered
5 intratumorally daily in a volume of 0.05 ml containing 0.025 µg 2-(phosphonomethyl)pentanedioic acid per injection.

Polymer containing 2-(phosphonomethyl)pentanedioic acid (10 µg drug/mg polymer) was implanted subcutaneously. Mice were anaesthetized with metaflane, and a small (<2mm) incision was made
10 near the tumor site. Following implantation, the incision was closed with a wound clip. Polymer was replaced twice weekly.

The tumor were measured twice weekly for at least 8 weeks after the first treatment. The mean tumor volume for each group was calculated for each time point. Comparisons between groups
15 at specific times were made using an unpaired, two-tailed t-test, and the results were analyzed using analysis of variance (ANOVA).

Systemic toxicity was assessed from reductions in body weight after treatment. The mice were sacrificed at the end of the follow-up period, or earlier if their tumor volumes reached
20 1600 mm³ or the tumors ulcerated.

Statistical Analysis

Statistical analysis as described above was performed using JMP (SAS Institute Inc., Cary, NC)

25

In vivo Rat Dunning R3327 Model

Referring now to FIGS. 5 and 6, Dunning R3327-G prostate cancer cells were injected subcutaneously into both flanks of

syngeneic male rats. In the first study, the anti-tumor growth activity of 2-(phosphonomethyl)pentanedioic acid was tested following daily subcutaneous injections of the drug (1, 3, 10 and 30 mg/kg). 2-(phosphonomethyl)pentanedioic acid injections and tumor measurements were continued for 12 weeks. In the second study, the anti-tumor growth activity of 2-[[phenylmethyl]hydroxyphosphinyl]methyl]pentanedioic acid was tested following daily intra-tumoral injections of the drug (0.1, 1, 10, 100 μ g) after the tumor reached an initial volume of 80-290 mm³. Tumor volumes were subsequently monitored for 42 days after the beginning of drug treatment.

EXPERIMENTAL PROCEDURES

Cell Lines

R3327-G is a cell line derived from an androgen-sensitive papillary adenocarcinoma derived from a spontaneously forming tumor in the rat prostate. R3327-G cells were grown in RPMI, 10% fetal bovine serum (FBS), 2 mM L-glutamine, and 10⁻⁸ M dexamethasone. Cultures were kept at 37°C in a humidified 5% CO₂/O₂ incubator. Antibiotics were not added to the medium.

Animal Tumor Model

Copenhagen male rats, age 8-10 weeks, were purchased from Harlan Sprague Dawley (Indianapolis, IN). The animals were housed two per cage. Upon arrival, they were quarantined for four working days before use. Temperature was maintained at 72 \pm 5°F and relative humidity at 35-70%, and a 12-hr light/dark cycle was used. The rats were fed certified Purina rodent chow

and water *ad libitum*.

After the animals were released from quarantine, the rats were injected subcutaneously in both flanks with 1×10^7 R3327-G cells (0.1-ml injection volume). Tumor dimensions and body weight were measured twice weekly. Vernier calipers were used to measure tumors in three planes, and tumor volume (V) was calculated as follows: $V = \pi(x \times y \times z)/6$, where x, y, and z were the tumor measurements minus skin thickness. Tumors began to appear 4-5 weeks after tumor cell injection. At the end of the experiment, the rats were sacrificed by CO₂ inhalation.

Pharmaceuticals

2-(phosphonomethyl)pentanedioic acid was made up in physiological saline fresh each day prior to injection. A stock solution of 2-[[phenylmethyl]hydroxyphosphinyl]methyl]pentanedioic acid was made up in water at a concentration 2.5 mg/ml; ten-fold serial dilutions were made fresh weekly for injections.

Treatment Protocol

In the 2-(phosphonomethyl)pentanedioic acid study, the rats were given daily subcutaneous injections of drug beginning the 14 days following tumor cell implantation and continued for 12 weeks. In the 2-[[phenylmethyl]hydroxyphosphinyl]methyl]pentanedioic acid study, the drug was not administered until the tumor volumes reached a predetermined size (mean tumor volume 90-290 mm³). At this time, the rats were divided into treatment groups of five rats each. All treatments of 2-

[[phenylmethyl]hydroxyphosphinyl]methyl] pentanedioic acid were subsequently administered intra-tumorally daily for 6 weeks.

The tumors were measured twice weekly. The mean tumor volume for each group was calculated for each time point.

5 Comparisons between groups at specific times were made using an unpaired, two-tailed t-test, and the results were analyzed using analyzed of variance (ANOVA). For the 2-

10 [[phenylmethyl]hydroxyphosphinyl]methyl] pentanedioic acid study, individual tumor volumes (V) were expressed as a fraction of the tumor volume on Day 0, the first day of treatment (V₀). For each group, the mean of the ratio V/V₀ was plotted as a function of time after treatment.

Statistical Analysis

15 Statistical analysis as described above was performed using JMP (SAS Institute, Inc. Cary, NC).

EXAMPLES

20 The following examples are illustrative of preferred embodiments of methods of use and preparation of compounds of the invention and are not to be construed as limiting the invention thereto. Unless otherwise indicated, all percentages are based upon 100% of the final formulations.

25 EXAMPLE 1

Preparation of 2-[(methyldihydroxyphosphinyl)methyl] pentanedioic acid

Scheme IV R=CH₃, R₁=CH₂Ph

Methyl-O-benzylphosphinic acid

Dichloromethylphosphite (10.0 g, 77 mmol) in 80 mL of dry diethyl ether was cooled to -20°C under an atmosphere of nitrogen. A solution of benzyl alcohol (23 g, 213 mmol) and triethylamine (10.2 g, 100 mmol) in 40 mL of diethyl ether was added dropwise over 1 hour while maintaining an internal temperature range of 0°C to 10°C. Once addition was complete the mixture was warmed to room temperature and stirred overnight. The mixture was filtered and the solid cake washed with 200 mL of diethyl ether. The organics were combined and evaporated under reduced pressure to give 25 g of a clear and colorless liquid. The liquid was purified by flash chromatography and eluted with a 1:1 hexane/ethyl acetate to ethyl acetate gradient. The desired fractions were collected and evaporated to give methyl O-benzylphosphinic acid (1, R=CH₃, R₁=CH₂Ph, 6.5 g, 50%) as a clear and colorless oil. R_f 0.1 (1:1, Hexane/EtOAc).
¹H NMR (d₆-DMSO): 7.4 ppm (m, 5H), 7.1 ppm (d, 1H), 5.0 ppm (dd, 2H), 1.5 ppm (d, 3H)

2,4-Di(benzylloxycarbonyl)butyl(methyl)-O-benzylphosphinic acid

Methyl-O-benzylphosphinic acid (3.53 g, 20.7 mmol) in 200 mL of dichloromethane was cooled to -5°C under an atmosphere of nitrogen. Triethylamine (3.2 g, 32 mmol) was added via syringe followed by trimethylsilyl chloride (2.9 g, 27 mmol). The reaction mixture was stirred and warmed to room temperature over 1 hour. Dibenzyl 2-methylenepentanedioate (2, 6.0 g, 18.5 mmol) in 10 mL of dichloromethane was added. The mixture was then stirred at room temperature overnight. The reaction mixture was

cooled to 0°C and trimethylaluminum (9 mL, 18 mmol, 2.0 M in dichloromethane) was added. The flask was warmed and stirred for 72 hours. The clear light yellow solution was cooled to 5°C and quenched by the slow addition of 5% hydrochloric acid. The quenched reaction mixture was warmed to room temperature and the organic layer removed. The organic layer was washed with 5% hydrochloric acid and with water. The organics were dried (MgSO₄) and evaporated under reduced pressure to give 8 g of a clear light yellow oil. The oil was purified on silica gel and eluted with a gradient of 1:1 hexanes/ethyl acetate to 100% ethyl acetate. The desired fractions were collected and evaporated to give 2,4-di(benzyloxycarbonyl)buty(methyl)-O-benzylphosphinic acid (3, R=CH₃, R₁=CH₂Ph 0.8 g, 8%) as a clear and colorless oil. R_f 0.5 (ethyl acetate).

¹H NMR (CDCl₃): 7.4 ppm (m, 15H), 5.1 ppm (m, 6H), 3.0 ppm (m, 1H), 2.4 ppm (m, 3H), 2.1 ppm (m, 3H), 1.5 ppm (dd, 3H)

Elemental Analysis Calculated C₂₈H₃₁O₆P.0.5H₂O:C 68.01, H 6.32

Found: C 66.85, H 6.35

20 2-[(Methylhydroxyphosphinyl)methyl]pentanedioic acid

2,4-di(benzyloxycarbonyl)buty(methyl)-O-benzylphosphinic acid (0.8 g, 1.6 mmol) in 20 mL of water containing 100 mg of 10% Pd/C was hydrogenated at 40 psi for 4 hours. The mixture was filtered over a pad of Celite and evaporated at high vacuum to give 2-[(methylhydroxyphosphinyl)methyl]pentanedioic acid (4, R=CH₃, 0.28 g, 78% as a clear and colorless viscous oil.

¹H NMR (D₂O): 2.5 ppm (m, 1H), 2.2 ppm (t, 2H), 2.0 ppm (m, 1H), 1.7 ppm (m, 3H), 1.3 ppm (d, 3H)

Elemental Analysis Calculated $C_7H_{13}O_5P \cdot 0.2 H_2O$: C 36.92 H 5.93

Found: C 37.06 H 6.31

EXAMPLE 2

5 Preparation of 2-[(butylhydroxyphosphinyl)methyl] pentanedioic acid

Scheme IV R=n-butyl, R1=H

Butylphosphinic Acid

10 Diethyl chlorophosphite (25g, 0.16mol) in 60 mL of dry ether was cooled to 0°C under an atmosphere of nitrogen.

Butylmagnesium chloride (80 mL, 0.16 mol, 2.0 M solution in ether) was added dropwise over a period of 2 hours while maintaining the internal temperature at 0°C. Once addition

15 was complete the thick white slurry was heated to 30°C for 1 hour. The suspension was filtered under a nitrogen atmosphere and the filtrate evaporated under reduced pressure. The clear light yellow liquid was then brought up in 15 mL of water and stirred at room temperature. Concentrated hydrochloric acid

20 (0.5 mL) was then added and an exothermic reaction was observed. The mixture was stirred an additional 15 minutes and extracted with two 75 mL portions of ethyl acetate. The organics were combined, dried ($MgSO_4$) and evaporated to give a clear and colorless liquid. The liquid was treated with NaOH

25 (40 mL, 20 M) and stirred for 1 hour. The mixture was then washed with diethyl ether and acidified to pH 1.0. The desired material was extracted from the acidified extract with two 100 mL portions of ethyl acetate. The organics were

combined, dried (MgSO_4) and evaporated under reduced pressure to give butylphosphinic acid (1, R=n-butyl, R1=H, 10g, 51%) as a clear and colorless liquid.

^1H NMR (d_6 -DMSO): 6.9 ppm(d, 1H), 1.6 ppm(m, 2H), 1.4 ppm(m, 4H), 0.9 ppm(t, 3H)

Butyl[2,4-di(benzyloxycarbonyl)butyl]phosphinic acid

Butylphosphinic acid (2.0g, 16mmol) in 80 mL of dry dichloromethane was cooled to 0°C under an atmosphere of nitrogen. Triethylamine (6.7 g, 66 mmol) was added followed by trimethylsilyl chloride (58 mL, 58 mmol, 1.0 M in dichloromethane). The mixture was stirred at 0°C for 10 minutes and dibenzyl 2-methylenepentanedioate (2) (6.4 g, 20 mmol) in 20 mL of dichloromethane was added. The cold bath was removed and the reaction warmed to room temperature and stirred overnight. The mixture was then cooled to 0°C and quenched by the slow addition of 5% hydrochloric acid. The dichloromethane layer was then removed and washed with 5% hydrochloric acid and with brine. The organic layer was dried (MgSO_4) and evaporated to give a clear light golden liquid. The liquid was purified by flash chromatography and eluted with 3:1 hexane/ethyl acetate containing 5% acetic acid. The desired fractions were combined and evaporated to give butyl[2,4-di(benzyloxycarbonyl)butyl]phosphinic acid (3, R=n-butyl, R1=H) (2.9 g, 40%) as a clear and colorless oil.

Rf 0.12 (3:1, Hex./EtOAc 5% AcOH).

^1H NMR (d_6 -DMSO): 7.3 ppm (m, 10), 5.0 ppm (s, 4H), 2.7 ppm (m, 1H) 2.3 ppm (y, 2H), 1.8 ppm (m, 2H), 1.3 ppm (m, 4H), 0.8 ppm

(t, 3H)

2-[(Butylhydroxyphosphinyl)methyl]pentanedioic acid

Butyl[2,4-di(benzyloxycarbonyl)butyl]phosphinic acid (2.9 g,
5 6.5 mmol) in 30 mL of water containing 0.32 g 10% Pd/C was
hydrogenated on a Parr hydrogenator at 40 psi for 4.5 hours.
The mixture was filtered through a pad of Celite and
evaporated under high vacuum to give 2-

10 [(butylhydroxyphosphinyl)methyl]pentanedioic acid (4, R=n-
butyl) (0.75 g, 43%) as a clear and colorless viscous oil.

¹H NMR (D₂O): 2.4 ppm (m, 1H), 2.1 ppm (t, 2H), 1.9 ppm (m,
1H), 1.6 ppm (m, 3H), 1.4 ppm (m, 2H), 1.1 ppm (m, 4H), 0.6
ppm (t, 3H)

Elemental Analysis Calculated C₁₀H₁₉O₆P. 0.5 H₂O: C 43.64, H
15 7.32: Found C 43.25, H 7.12

EXAMPLE 3

Preparation of 2-[(benzylhydroxyphosphinyl)methyl]pentanedioic
acid

20 Scheme IV R=CH₂Ph, R₁=H

Benzylphosphinic acid

Diethylchlorophosphite (25 g, 0.16 mol) in 100 mL of dry
diethyl ether was cooled to 0°C under an atmosphere of
25 nitrogen. Benzylmagnesium chloride (80 mL, 0.16 mol, 2.0 M
solution in Et₂O) was added dropwise over two hours while
maintaining a temperature below 10°C. A thick white slurry
formed and stirring was continued at room temperature for 1

hour. The mixture was filtered under a nitrogen atmosphere and the filtrate evaporated under reduced pressure to give a clear and colorless liquid. The liquid was stirred as 15 mL of water was added followed by 0.5ml concentrated hydrochloric acid. An exothermic reaction was observed and stirring was continued for an additional 30 minutes followed by extraction with ethyl acetate. The organics were combined, washed with brine, dried (MgSO_4) and evaporated. The clear light golden liquid was added to sodium hydroxide (50 mL, 2.0 M NaOH), stirred for one hour and washed with diethyl ether. The aqueous layer was acidified to pH 1.0 with concentrated hydrochloric acid and extracted with ethyl acetate. The organics were combined, dried (MgSO_4) and evaporated to give benzylphosphinic acid (1, $\text{R}=\text{CH}_2\text{Ph}$, R_1+H) (8 g, 32%) as a clear light golden oil.

^1H NMR (d_6 -DMSO): 7.3 ppm (m, 5H), 6.9 ppm (d, 1H), 3.1 ppm (d, 2H)

Benzyl[2,4-di(benzylloxycarbonyl)butyl]phosphinic acid

Benzylphosphinic acid (2.3 g, 15 mmol) in 150 mL of dry dichloromethane was cooled to 0°C under a nitrogen atmosphere. Triethylamine (6.5 g, 65mmol) was added followed by trimethylsilyl chloride (5.8 g, 54 mmol) while the reaction temperature was maintained at 0°C . After 30 minutes dibenzyl 2-methylenepentanedioate (2) in 20 mL of dichloromethane was added over 5 minutes. The reaction mixture was left to warm to room temperature and stirred overnight. The clear solution was cooled to 0°C and quenched with 5% hydrochloric acid and

with brine, dried (MgSO_4) and evaporated to give a clear yellow liquid. Purification by flash chromatography and elution with 1:1 hexane/ethyl acetate containing 10% acetic acid yielded 2.0 g (28%) of benzyl[2,4-

5 di(benzyloxycarbonyl)butyl]phosphinic acid (3, $\text{R}=\text{CH}_2\text{Ph}$, $\text{R}_1=\text{H}$) as a clear light yellow oil. R_f 0.37 (1:1 Hex./EtOAc, 10%AcOH).

^1H NMR (d_6 -DMSO): 7.2 ppm(m, 15H), 5.0 ppm(s, 4H), 3.0 (d, 2H), 2.8 ppm(m, 1H), 2.3 ppm(t, 2H), 1.9 ppm(m, 2H), 1.7 ppm(t, 1H)

10 .

2-[(Benzylhydroxyphosphinyl)methyl]pentanedioic acid

Benzyl[2,4-di(benzyloxycarbonyl)butyl]phosphinic acid (0.5 g, 1.0 mmol) in 20 mL of water containing 120 mg of 10% Pd/C was hydrogenated on a Parr hydrogenator at 40 psi for 6 hours.

15 Filtration through a Celite pad followed by evaporation on high vacuum gave 0.17 g (57%) of 2-[(benzylhydroxyphosphinyl)methyl]pentanedioic acid (4, $\text{R}=\text{CH}_2\text{Ph}$) as a white solid.

^1H NMR (D_2O): 7.1 ppm(m, 5H), 2.9 ppm(d, 2H), 2.4 ppm(m, 1H),

20 2.1 ppm(t, 2H), 1.8 ppm(m, 1H), 1.6 ppm(m, 3H)

Elemental Analysis, Calculated $\text{C}_{13}\text{H}_{17}\text{O}_5\text{P}$: C52.00H5.71: Found: C51.48H5.70

EXAMPLE 4

25 Preparation of 2-

[phenylacetylhydroxyphosphinyl)methyl]pentanedioic acid

Scheme IV $\text{R}=\text{CH}_2\text{CH}_2\text{Ph}$, $\text{R}_1=\text{H}$

Phenethylphosphinic acid

Diethylchlorophosphite (15.6 g, 0.1 mol) in 100 mL of dry diethyl ether was cooled to 5°C under an atmosphere of nitrogen. Phenethylmagnesium chloride (100 mL, 0.1 mol, 1.0 M in THF) was added dropwise over 2 hours while maintaining a temperature between 0-10°C. A thick white slurry formed and stirred at room temperature overnight. The mixture was filtered under a nitrogen atmosphere and the filtrate evaporated under reduced pressure to give a clear and colorless liquid. The liquid was stirred as 15 mL of water was added followed by 0.5 mL of concentrated hydrochloric acid. An exothermic reaction was observed and stirring continued for 15 minutes followed by extraction with ethyl acetate. The organics were combined, washed with brine, dried (MgSO₄) and evaporated. The clear liquid was brought up in sodium hydroxide (40 mL, 2.0 M NaOH), stirred for 1 hour and washed once with diethyl ether. The aqueous layer was acidified to pH 1.0 with concentrated hydrochloric acid and extracted with ethyl acetate. The organics were combined, dried (MgSO₄) and evaporated to give phenethylphosphinic acid (1, R=CH₂CH₂Ph, R₁=H) (9.8 g, 58%) as a clear light yellow oil.

¹H NMR (d₆-DMSO): 7.2 ppm (m, 5H), 6.9 ppm (d, 1H), 2.8 ppm (m, 2H), 1.9 ppm (m, 2H)

2,4-Di(benzylloxycarbonyl)butyl(phenethyl)phosphinic acid

Phenethylphosphinic acid (1.0 g, 5.9 mmol) in 50 mL of dry dichloromethane was cooled to -5°C under a nitrogen

atmosphere. Triethylamine (2.3g, 23 mmol) was added followed by trimethylsilyl chloride 2.2 g, 21 mmol) while the reaction temperature was maintained at 0°C. After 10 minutes dibenzyl 2-methylenepentanedioate (2) in 10 mL of dichloromethane was added over 10 minutes. The reaction mixture was left to warm to room temperature and stirred overnight. The clear solution was cooled to 0°C and quenched with 5% hydrochloric acid followed by removal of the organic layer. The organic layer was washed with brine, dried (MgSO₄) and evaporated to give a clear light golden liquid. Purification by flash chromatography and elution with 1:1 Hexane/EtOAc containing 5% AcOH resulted in 1.2g (41%) of 2,4-di(benzyloxycarbonyl)butyl(phenethyl)phosphinic acid(3,R=CH₂CH₂Ph,R₁=H) as a clear and colorless oil.

¹H NMR (d₆-DMSO): 7.2 ppm(m,15H), 5.0 ppm (s,4H), 3.3 ppm (m,1H), 2.8 ppm(m,4H), 2.3 ppm(m,2), 1.8 ppm(m,4H)

2,4-[(Phenethylhydroxyphosphinyl)methyl]pentanedioic acid

2,4-Di(benzyloxycarbonyl)butyl(phenethyl)phosphinic acid(1.1 g,2.2 mmol) in 20 mL of water containing 120 mg of 10% Pd/C was hydrogenated on a Parr hydrogenator at 40 psi overnight. Filtration through a Celite pad followed by evaporation on high vacuum gave 0.8 g (114%) of 2-[(phenethylhydroxyphosphinyl)methyl]pentanedioic

acid(4,R=CH₂CH₂Ph) as a white solid.

¹H NMR(D₂O): 7.2 ppm (m,5H), 2.7 ppm (m,2H), 2.5 ppm (m,1H), 2.3 ppm (t,2H), 1.9 ppm (m,6H), 1.5 ppm (t,1H)

Elemental Analysis:

Calculated $C_{14}H_{19}O_5P \cdot 0.75H_2O \cdot 0.5 AcOH$: C 50.35 H 6.34

Found: C 50.26 H 5.78

EXAMPLE 5

5 Preparation of 2-[(3-phenylpropylhydroxyphosphinyl)methyl]pentanedioic acid

Scheme IV R=CH₂CH₂CH₂Ph, R₁=H

3-Phenylpropylphosphinic acid

10 Magnesium turnings (2.44 g, 0.10 mol) in 20 mL of dry diethyl ether under an atmosphere of nitrogen was added several iodine crystals. Phenylpropyl bromide (20.0 g, 0.10 mol) in 80 mL of diethyl ether was placed in a dropping funnel. Approximately 10 mL of the bromide solution was added to the magnesium

15 turnings and stirring was initiated. After several minutes the iodine was consumed and additional phenylpropyl bromide was added while maintaining a temperature of 35°C. Once additional was complete (1.5 hours) the mixture was sealed and stored at 5°C.

20 Diethylchlorophosphite (15.7 g, 0.1 mol) in 50 mL of dry diethyl ether was cooled to 5°C under an atmosphere of nitrogen. Phenylpropylmagnesium bromide (100 mL, 0.1 mol, 1.0 M solution of in Et₂O) was added dropwise over 2 hours while maintaining a temperature between 0-10°C. A thick white

25 slurry formed and was stirred an additional 30 minutes. The mixture was filtered under a nitrogen atmosphere and the filtrate evaporated under reduced pressure to give a clear and colorless liquid. To the liquid was added 20 mL of water

followed by 0.5ml of concentrated hydrochloric acid. An exothermic reaction was observed and stirring continued for 20 minutes followed by extraction with ethyl acetate. The organics were combined, washed with brine, dried (MgSO_4) and evaporated. To the clear liquid was added sodium hydroxide (40 mL, 2.0 M NaOH), the resulting solution stirred for 1 hour and then washed with diethyl ether. The aqueous layer was acidified to pH 1.0 with concentrated hydrochloric acid and extracted twice with ethyl acetate. The organics were combined, dried (MgSO_4) and evaporated to give 3-phenylpropylphosphinic acid (1, $\text{R}=\text{CH}_2\text{CH}_2\text{CH}_2\text{Ph}$, $\text{R}_1=\text{H}$) (9.8 g, 53%) as a clear and colorless oil.

^1H NMR (d_6 -DMSO): 7.2 ppm (m, 5H), 6.9 ppm (d, 1H), 2.6 ppm (t, 2H), 1.7 ppm (m, 2H), 1.6 ppm (m, 2H)

2,4-Di(benzylloxycarbonyl)butyl(3-phenylpropyl)phosphinic acid
3-phenylpropylphosphinic acid (1.0 g, 5.4 mmol) in 50 mL of dry dichloromethane was cooled to -5°C under a nitrogen atmosphere. Triethylamine (2.2 g, 22 mmol) was added followed by trimethylsilyl chloride (2.1 g, 19 mmol) while the reaction temperature was maintained at 0°C . After 10 minutes dibenzyl 2-methylenepentanedioate (2) in 10 mL of dichloromethane was added over 10 minutes. The reaction mixture was warmed to room temperature and stirred overnight. The clear solution was cooled to 0°C and quenched with 5% hydrochloric acid followed by removal of the organic layer. The organic layer was washed with brine, dried (MgSO_4) and evaporated to give a clear yellow liquid. Purification by flash chromatography and

elution with 4:1 hexane/ethyl acetate containing 5% acetic acid resulted in 1.5g (56%) of 2,4-di(benzyloxycarbonyl)butyl(3-phenylpropyl)phosphinic acid(3,R=CH₂CH₂CH₂Ph, R₁=H) as a clear light yellow oil. R_f 0.58 (1:1 Hex./EtOAc, 5%AcOH);

¹H NMR (d₆-DMSO): 7.2 ppm (m, 15H), 5.0 ppm (s, 4H), 2.7 ppm (m, 1H), 2.5 ppm (m, 5H), 2.2 ppm (m, 2H), 1.8 ppm (m, 3H), 1.6 ppm (m, 2H)

Elemental Analysis:

Calculated C₂₃H₃₃O₅P. 1.3H₂O: C 65.48 H 6.75

Found: C 65.24 H 6.39

2-[(3-Phenylpropyl)hydroxyphosphinyl)methyl]pentanedioic acid

2,4-Di(benzyloxycarbonyl)butyl(3-phenylpropyl)phosphinic acid(15) (1.4 g, 2.8 mmol) in 20 mL of water containing 150 mg of 10% Pd/C was hydrogenated on a Parr hydrogenator at 40 psi overnight. Filtration through a Celite pad followed by evaporation on high vacuum gave 0.8 g (89%) of 2-[(3-phenylpropyl)hydroxyphosphinyl)methyl]pentanedioic acid(4,R=CH₂CH₂CH₂Ph) as a light yellow viscous oil).

¹H NMR (D₂O): 7.4 ppm (m, 5H), 2.7 ppm (m, 3H), 2.4 ppm (t, 3H), 1.8 ppm (m, 7H);

Elemental Analysis:

Calculated C₁₃H₂₁O₅P 0.75 H₂O, 0.75 AcOH: C 51.23 H 6.64

Found: C 50.85 H 6.02

EXAMPLE 6

Preparation of 2-[(4-methylbenzyl)hydroxyphosphinyl]

methylpentanedioic acid

Scheme V, Compound 5

Hexamethyldisilazane (21.1 mL, 100 mmol) was added to vigorously stirred ammonium phosphinate (8.30 g, 100 mmol), and the resulting suspension was stirred at 105 C for 2 h. A solution of 4-methylbenzyl bromide (5.00 g, 27.0 mmol) was then dropwise added to the suspension at 0°C. The mixture was stirred at rt for 19 h. The reaction mixture was then diluted with dichloromethane (50 mL) and washed with 1 N HCl (50 mL).

The organic layer was separated, dried over Na₂SO₄, and concentrated to give 4.72 g of a white solid. This was dissolved in dichloromethane (50 mL) and benzyl alcohol (3.24 g, 30 mmol) was added to the solution. 1,3-

Dicyclohexylcarbodiimide (DCC) (6.19 g, 30 mmol) was then added to the solution at 0°C, and the suspension was stirred at rt for 14 h. The solvent was removed under reduced pressure and the residue was suspended in EtOAc. The resulting suspension was filtered and the filtrate was concentrated. The residue was purified by silica gel chromatography (hexanes: EtOAc, 4:1 to 1:1) to give 2.40 g of 4-methylbenzyl-O-benzylphosphinic acid (2, R=4-methylbenzyl) as a white solid (34% yield): *R*_f 0.42 (EtOAc); ¹H NMR (DMSO-*d*₆) delta 2.30 (s, 3 H), 3.29 (d, *J* = 16.6 Hz, 2 H), 5.2 (m, 2 H), 7.0 (d, *J* = 543 Hz, 1 H), 7.1-7.2 (m, 4 H), 7.3-7.4 (m, 5 H).

To a solution of 4-methylbenzyl-O-benzylphosphinic acid (2, R=4-methylbenzyl) (2.16 g, 8.3 mmol) in THF (15 mL) was added sodium hydride (0.10g, 60 % dispersion in oil) followed by

dibenzyl 2-methylenepentanedioate at 0 C, and the mixture was stirred at rt for 4 h. The reaction mixture was then diluted with EtOAc (50 mL) and poured into 1N HCl (50 mL). The organic layer was separated, dried over Na₂SO₄, and concentrated. This material was purified by silica gel chromatography (hexanes: EtOAc, 4:1 to 1:1) to give 3.41 g of 2,4-di(benzyloxycarbonyl)butyl(4-methylbenzyl)-o-benzylphosphinic acid (4, R = 4-methylbenzyl) as colorless oil (70% yield): *Rf* 0.61 (EtOAc); ¹H NMR (CDCl₃) delta 1.6-1.8 (m, 1 H), 1.9-2.0 (m, 2 H), 2.1-2.4 (m, 6 H), 2.7-2.9 (m, 1 H), 3.05 (dd, *J* = 9.0, 16.8 Hz, 2 H), 4.8-5.1 (m, 6 H), 7.0-7.1 (m, 4 H), 7.2-7.4 (m, 15 H).

To a solution of 2,4-di(benzyloxycarbonyl)butyl(4-methylbenzyl)-o-benzylphosphinic acid (0.70 g, 1.2 mmol) in ethanol (30 mL) was added Pd/C (5%, 0.10 g) and the suspension was shaken under hydrogen (50 psi) for 18 h. The suspension was then filtered through a pad of Celite and concentrated under reduced pressure. The resulting residue was dissolved in distilled water (5 mL), passed through a column of AG 50W-X8 resin (H⁺ form), and lyophilized to give 0.21 g of 2-[[[4-methylbenzyl]hydroxyphosphinyl]methyl]pentanedioic acid (5, R = 4-methylbenzyl) as a white solid (55% yield): *Rf* 0.62 (*i*-PrOH:H₂O, 7:3); ¹H NMR (D₂O) delta 1.7-1.9 (m, 3 H), 2.0-2.2 (m, 1 H), 2.33 (dt, *J* = 1.7 Hz, 7.4 Hz, 2 H), 2.55-2.70 (m, 1 H), 3.12 (d, *J* = 16.5 Hz, 2 H), 7.0-7.1 (m, 2 H), 7.2-7.3 (m, 2 H). Anal. Calcd for C₁₃H₁₇O₅P*0.30H₂O: C, 52.60; H, 6.18. Found: C, 52.60; H, 6.28.

EXAMPLE 7

Preparation of 2-[[[4-Fluorobenzyl]hydroxyphosphinyl]

methylpentanedioic acid (R = 4-fluorobenzyl):

Scheme V, prepared as described in the above example where R =
5 methylbenzyl:

R_f 0.64 (i-PrOH:H₂O, 7:3); ¹H NMR (D₂O) delta 1.7-1.9 (m, 3 H),
2.0-2.2 (m, 1 H), 2.3-2.4 (m, 2 H), 2.55-2.70 (m, 1 H), 3.12
(d, J = 16.5 Hz, 2 H), 7.0-7.1 (m, 2 H), 7.2-7.3 (m, 2 H).

Anal. Calcd for C₁₃H₁₅FO₅P*0.25H₂O: C, 48.38; H, 5.15. Found: C,
10 48.38; H, 5.15.

EXAMPLE 8

Preparation of 2-[[[4-Methoxybenzyl]hydroxyphosphinyl]

methylpentanedioic acid (R = 4-methoxybenzyl):

15 Scheme V, prepared as described in the above example where R =
methylbenzyl:

R_f 0.56 (i-PrOH:H₂O, 7:3); ¹H NMR (D₂O) delta 1.8-1.9 (m, 3 H),
2.0-2.2 (m, 1 H), 2.3-2.4 (m, 2 H), 2.55-2.70 (m, 1 H), 3.16
(d, J = 16.7 Hz, 2 H), 3.81 (s, 3 H), 6.98 (d, J = 8.7 Hz, 2

20 H), 7.25 (d, J = 8.7 Hz, 2 H). Anal. Calcd for
C₁₄H₁₉O₇P*0.30H₂O: C, 50.09; H, 5.89. Found: C, 49.98; H, 5.80.

EXAMPLE 9

Preparation of 2-[[[2-Fluorobenzyl]hydroxyphosphinyl]

25 methylpentanedioic acid (R = 2-fluorobenzyl):

Scheme V, prepared as described in the above example where R
= methylbenzyl:

R_f 0.67 (i-PrOH:H₂O, 7:3); ¹H NMR (D₂O) delta 1.8-1.9 (m, 3 H),

2.0-2.2 (m, 1 H), 2.3-2.4 (m, 2 H), 2.55-2.70 (m, 1 H), 3.28 (d, $J = 16.6$ Hz, 2 H), 7.1-7.5 (m, 4 H). Anal. Calcd for $C_{13}H_{15}FO_6P \cdot 0.10H_2O$: C, 48.79; H, 5.10. Found: C, 48.84; H, 5.14.

5

EXAMPLE 10

Preparation of 2-[[[Pentafluorobenzyl]hydroxyphosphinyl]methyl]pentanedioic acid (R = pentafluorobenzyl):

Scheme V, prepared as described in the above example where R = methylbenzyl:

10 R_f 0.69 (i-PrOH:H₂O, 7:3); ¹H NMR (D₂O) delta 1.8-2.0 (m, 3 H), 2.1-2.3 (m, 1 H), 2.3-2.5 (m, 2 H), 2.7-2.9 (m, 1 H), 3.29 (d, $J = 15.4$ Hz, 2 H), Anal. Calcd for $C_{13}H_{12}F_5O_5P \cdot 0.45H_2O$: C, 39.20; H, 3.26. Found: C, 39.17; H, 3.28.

15

EXAMPLE 11

Preparation of 2-[(methylhydroxyphosphinyl)methyl]pentanedioic acid

Scheme VI, Compound 9

20 2,4-Di(benzyloxycarbonyl)butylphosphinic acid (6)
Dry phosphinic acid (100 g, 1.52 mol) was dissolved in 100 ml of chloroform and treated with triethylamine (155 g, 1.52 mol). The mixture was evaporated and transferred to a three liter flask, containing 750 mL of chloroform. The solution was
25 stirred by means of a mechanical stirrer and the flask cooled to 0°C. The clear solution was treated with triethylamine (277 g, 2.72 mol) followed by trimethylsilyl chloride (281 g, 2.58 mol). Once addition of trimethylsilyl chloride was

complete dibenzyl 2-methylenepentanedioate (2) in 150 mL of chloroform was added dropwise over 20 minutes. The low temperature bath was removed and the mixture warmed to room temperature. After 6 hours the thick slurry was filtered and the filtrate cooled to 0°C. The filtrate was then quenched with 5% hydrochloric acid and the organic layer removed. The aqueous layer was extracted with chloroform, the organics combined, dried (MgSO₄) and evaporated under reduced pressure to give 55 g of 2,4-di(benzyloxycarbonyl)butylphosphinic acid (6) as a light yellow liquid. The liquid was purified by flash chromatography and eluted using 3:1 hexanes/ethyl acetate containing 5% trifluoroacetic acid to give 40 g (7%) of the desired product. Rf 0.28 (3:1 Hex./EtOAc 5% TFA); ¹H NMR (CDCl₃): 7.3 ppm (m, 10H), 7.2 ppm (d, 1H), 5.12 ppm (s, 2H), 2.9 ppm (m, 1H), 2.4 ppm (t, 2H), 2.2 ppm (m, 1H), 2.0 ppm (m, 3H)

2,4-Di(benzyloxycarbonyl)butylbenzylphosphinic acid (7)

To a solution of 2,4-di-(benzyloxycarbonyl)butyl phosphinic acid (6) (19.3 g, 49.4 mmol) in tetrahydrofuran was added benzyl alcohol (5.3 g, 49.3 mmol) and dimethylamino in tetrahydrofuran was added benzyl alcohol (5.3 g, 49.3 mmol) and dimethylamino pyridine (0.5 g). Dicyclohexylcarbodiimide (DCC, 12g, 58 mmol) was added and a white precipitate formed. After 30 minutes the white suspension was filtered and the filtrate evaporated under reduced pressure. The clear and colorless oil was purified by flash chromatography and eluted with 1:1 Hex./EtOAc to give 2,4-

di(benzyloxycarbonyl)butylbenzylphosphinic acid (7) (11.5 g, 47%) as a clear and colorless oil. Rf. 0.16 (1:1 Hex./EtOAc); ¹H NMR (CDCl₃): 7.3 ppm (m, 15H), 7.2 ppm (d, 1H), 5.0 ppm (m, 6H), 2.9 ppm (m, 1H), 2.2 ppm (m, 3H), 1.9 ppm (m, 3H)

5

2,4-Di(benzyloxycarbonyl)butyl[hydroxy(phenyl)methyl]benzylphosphinic acid (8)

2,4-Di(benzyloxycarbonyl)butylbenzylphosphinic acid (7) in 5 mL of dry THF was added dropwise to a stirring cooled (0°C) mixture of sodium hydride (0.09 g, 2.3 mmol) in 15 mL of THF. After 15 minutes benzaldehyde (0.23 g, 2.2 mmol) was added via syringe while maintaining a temperature of 0°C. After 30 minutes the mixture was quenched with water and extracted with two portions of dichloromethane. The organics were combined and evaporated to give a clear colorless oil. The oil was chromatographed on silica and eluted with a 1:1 Hex./EtOAc solvent system. The desired fractions were collected and evaporated to give 0.4 g (33%) of 2,4-di(benzyloxycarbonyl)butyl[hydroxy(phenyl)methyl]benzylphosphinic acid (6) as a clear and colorless oil. Rf 0.18 (1:1 Hex./EtOAc);

¹H NMR (CDCl₃): 7.3 ppm (m, 20H), 5.2 ppm (m, 1H), 4.9 ppm (m, 6H), 2.8 ppm (dm, 1H), 2.2 ppm (m, 3H), 1.9 ppm (m, 3H)

25 : 2,4-Di(benzyloxycarbonyl)butyl[hydroxy(phenyl)methyl]benzylphosphinic acid (9)

2,4-Di(benzyloxycarbonyl)butyl[hydroxy(phenyl)methyl]benzylphosphinic acid (6) (0.37 g, 0.6 mmol) in 25 mL of

water containing 0.10g of 10% Pd/C was hydrogenated at 40 psi for 6 hours. The mixture was filtered through a pad of Celite and lyophilized to give 2-

5 ([hydroxy(phenyl)methyl]hydroxyphosphinylmethyl)pentanedioic acid (9) (0.14 g, 70%) as a white solid.

^1H NMR (D_2O): 7.4 ppm (m, 5H), 5.0 ppm (d, 1H), 2.7 ppm (m, 1H), 2.4 ppm (m, 2H), 2.2 ppm (m, 1H), 1.9 ppm (m, 3H)

Element Analysis:

Calculated $\text{C}_{11}\text{H}_{17}\text{O}_7\text{P}$ 0.6 H_2O : C 47.74 H 5.61

10 Found: C 47.73 H 5.68

EXAMPLE 12

Preparation of Dibenzyl 2-Methylenepentanedioate.

Scheme III.

15 Benzyl acrylate (500 g, 3 mol) was heated to 100°C under an atmosphere of nitrogen. The heating was stopped and HMPT (10 g, 61 mmol) was added dropwise while maintaining an internal temperature of 135-145°C. Once addition was complete the mixture was cooled to room temperature and a slurry of
20 silica with 5:1 Hex/EtOAc was added. The slurry was then transferred to a column containing a plug of dry silica. The column was then washed with 1:1 Hex/EtOAc and the solvent was collected and evaporated. The clear yellow liquid was
25 distilled under high vacuum (200 μHg) to give an initial fraction of 8 g distilling at 45°C and then the desired product at 180-185°C (212 g, 42 %) as a clear and colorless liquid.

^1H -NMR (CDCl_3)

7.3 ppm (s, 10H); 6.2 ppm (s, 1H); 5.5 ppm (s, 1H); 5.2 ppm

(s, 2H); 5.1 ppm (s, 2H); 2.6 ppm (m, 4H).

EXAMPLE 13

Preparation of Dibenzyl 2-[[Bis(benzvloxv)phosphorvllmethvll]-
5 pentanedioate.

Scheme III

Dibenzyl phosphite (9.5g, 36mmol) in 350ml of
dichloromethane was cooled to 0°C. To this stirring solution
was added trimethyl aluminum (18.2ml, 2.0M solution in hexane,
10 36.4mmol). After 30 minutes 1 (6.0g, 37 mmol) in 90 ml of
dichloromethane was added dropwise over 10 minutes. The clear
and colorless solution was then warmed to room temperature and
left to stir overnight. The mixture was then quenched by the
slow addition of 5% HCl. After stirring an additional 1.5
15 hours the lower organic layer was removed and the aqueous
layer extracted once with 100ml of dichloromethane. The
organics were combined, dried (MgSO₄), and evaporated to give a
clear light golden liquid. The liquid was chromatographed on
silica gel (4cm*30cm) and eluted with a gradient (4:1-1:1)
20 solvent system (Hexane/EtOAc). The fractions containing the
desired product were combined and evaporated to yield 2 (7.1g,
42%) as a clear and colorless liquid. The liquid was then
distilled on a Kughleror apparatus at 0.5mm Hg and 195-200°C.
The distillate was discarded and the remaining light golden
25 oil was chromatographed on silica gel (1:1, Hex./EtOAc) to
give 2.9g of 2 as a clear and colorless oil. TLC R_f 0.5 (1:1,
Hex./EtOAc).

¹H-NMR (CDCl₃)

-114-

7.1-7.4 (m, 20H); 5.05 (s, 2H); 4.8-5.03 (m, 6H); 2.8 (1H);
2.22-2.40 (m, 3H); 1.80-2.02 (m, 3H).

EXAMPLE 14

5 Preparation of 2-(Phosphonomethyl)pentanedioic Acid.

Scheme III

The benzyl pentanedioate (2.9g, 4.9mmol) was added to a mixture of 20ml of methanol containing 0.29g (6mol%) of 10% Pd/C. This mixture was hydrogenated on a Parr hydrogenator at
10 40 psi for 24 hours, filtered and evaporated to give 3 (1.0g, 90%) as a clear slightly golden viscous oil.

¹H-NMR (D₂O)

2.6-2.78 (m, 1H); 2.25-2.40 (m, 2H); 1.75-2.15 (m, 4H).

15

EXAMPLE 15

A patient is diagnosed with adenocarcinoma of the prostate. The patient may then be administered a NAALADase inhibitor, such as set forth in examples 1 through 3, by direct injection into the tumor. After this initial
20 treatment, the patient may be optionally administered the same or different NAALADase inhibitor by intermittent or continuous administration by subdural pump. It would be expected that no further occurrences of the adenocarcinoma would develop.

EXAMPLE 16

25 A patient is diagnosed with adenocarcinoma of the prostate. The patient may then be administered a NAALADase inhibitor, such as set forth in examples 1 through 3, by direct injection into the tumor. After this initial

treatment, the patient may be optionally administered the same or different NAALADase inhibitor by intermittent or continuous administration by implantation of a biocompatible, polymeric matrix delivery system. It would be expected that no further occurrences of the adenocarcinoma would develop.

EXAMPLE 17

A patient is diagnosed with benign prostatic hyperplasia. The patient may then be administered a NAALADase inhibitor, such as set forth in examples 1 through 3, by direct injection into the tumor. After this initial treatment, the patient may be optionally administered the same or different NAALADase inhibitor by intermittent or continuous administration by injection, subdural pump, or polymeric matrix implant. It would be expected that the benign prostatic hyperplastic cells do not develop into carcinoma.

EXAMPLE 18

A patient is diagnosed with adenocarcinoma of the prostate. The adenocarcinoma appears not to have metastasized. The adenocarcinoma would be removed by surgery. After post-operative recovery, the patient would be locally administered NAALADase inhibitor by intermittent or continuous administration by injection, subdural pump or by polymeric matrix implant. It would be expected that no further occurrences of the carcinoma would develop.

EXAMPLE 19

A patient is diagnosed with metastatic adenocarcinoma of the prostate. The adenocarcinoma appears to have metastasized, but surgery still is indicated as an effective

treatment modality. Tumor tissue would be removed by surgery. The patient would be locally administered a NAALADase inhibitor such as described herein from the time, approximately, of the initial diagnosis and would continue after surgery. After post-operative recovery, the patient would be maintained at this level of NAALADase inhibitor by a regimen of periodic local administration. The patient would be monitored carefully for intolerable adverse side-effects of NAALADase inhibitor administration. It would be expected that no further tumors develop. If some of the original, small tumorous masses are detected after surgery, they would be expected to not grow in size.

EXAMPLE 20

A patient is diagnosed with ACTH-producing tumors. The patient may then be administered a NAALADase inhibitor, such as set forth in examples 1 through 3, by direct injection into the tumor. After this initial treatment, the patient may be optionally administered the same or different NAALADase inhibitor by direct injection, subdural pump, or implantation of a biocompatible, polymeric matrix delivery system. It would be expected that tumor growth or tumor cell growth would be prevented or inhibited and that no further occurrences of the ACTH-producing tumor would develop.

EXAMPLE 21

A treatment such as that described in Example 9 wherein the patient is diagnosed with acute lymphocytic leukemia.

EXAMPLE 22

A treatment such as that described in Example 9 wherein

the patient is diagnosed with acute non-lymphocytic leukemia.

EXAMPLE 23

5 A treatment such as that described in Example 9 wherein the patient is diagnosed with metastatic or non-metastatic cancer of the adrenal cortex.

EXAMPLE 24

10 A treatment such as that described in Example 9 wherein the patient is diagnosed with metastatic or non-metastatic bladder cancer.

EXAMPLE 25

15 A treatment such as that described in Example 9 wherein the patient is diagnosed with metastatic or non-metastatic brain cancer.

EXAMPLE 26

A treatment such as that described in Example 9 wherein the patient is diagnosed with metastatic or non-metastatic breast cancer.

EXAMPLE 27

20 A treatment such as that described in Example 9 wherein the patient is diagnosed with metastatic or non-metastatic cervical cancer.

EXAMPLE 28

25 A treatment such as that described in Example 9 wherein the patient is diagnosed with metastatic or non-metastatic chronic lymphocytic leukemia.

EXAMPLE 29

A treatment such as that described in Example 9 wherein

the patient is diagnosed with metastatic or non-metastatic chronic myelocytic leukemia.

EXAMPLE 30

5 A treatment such as that described in Example 9 wherein the patient is diagnosed with metastatic or non-metastatic colorectal cancer.

EXAMPLE 31

10 A treatment such as that described in Example 9 wherein the patient is diagnosed with metastatic or non-metastatic cutaneous T-cell lymphoma.

EXAMPLE 32

A treatment such as that described in Example 9 wherein the patient is diagnosed with metastatic or non-metastatic endometrial cancer.

15

EXAMPLE 33

A treatment such as that described in Example 9 wherein the patient is diagnosed with metastatic or non-metastatic esophageal cancer.

EXAMPLE 34

20 A treatment such as that described in Example 9 wherein the patient is diagnosed with metastatic or non-metastatic Ewing's sarcoma.

EXAMPLE 35

25 A treatment such as that described in Example 9 wherein the patient is diagnosed with metastatic or non-metastatic gallbladder cancer.

EXAMPLE 36

A treatment such as that described in Example 9 wherein

the patient is diagnosed with metastatic or non-metastatic hairy cell leukemia.

EXAMPLE 37

5 A treatment such as that described in Example 9 wherein the patient is diagnosed with metastatic or non-metastatic head and neck cancer.

EXAMPLE 38

10 A treatment such as that described in Example 9 wherein the patient is diagnosed with metastatic or non-metastatic Hodgkin's lymphoma.

EXAMPLE 39

A treatment such as that described in Example 9 wherein the patient is diagnosed with metastatic or non-metastatic Kaposi's sarcoma.

15 EXAMPLE 40

A treatment such as that described in Example 9 wherein the patient is diagnosed with metastatic or non-metastatic kidney cancer.

EXAMPLE 41

20 A treatment such as that described in Example 9 wherein the patient is diagnosed with metastatic or non-metastatic liver cancer.

EXAMPLE 42

25 A treatment such as that described in Example 9 wherein the patient is diagnosed with metastatic or non-metastatic lung cancer (small cell and/or non-small cell).

EXAMPLE 43

A treatment such as that described in Example 9 wherein

the patient is diagnosed with metastatic or non-metastatic malignant peritoneal effusion.

EXAMPLE 44

5 A treatment such as that described in Example 9 wherein the patient is diagnosed with metastatic or non-metastatic malignant pleural effusion.

EXAMPLE 45

10 A treatment such as that described in Example 9 wherein the patient is diagnosed with metastatic or non-metastatic melanoma.

EXAMPLE 46

A treatment such as that described in Example 9 wherein the patient is diagnosed with metastatic or non-metastatic mesothelioma.

15

EXAMPLE 47

A treatment such as that described in Example 9 wherein the patient is diagnosed with metastatic or non-metastatic multiple myeloma.

20

EXAMPLE 48

A treatment such as that described in Example 9 wherein the patient is diagnosed with metastatic or non-metastatic neuroblastoma.

EXAMPLE 49

25 A treatment such as that described in Example 9 wherein the patient is diagnosed with metastatic or non-metastatic non-Hodgkin's lymphoma.

EXAMPLE 50

A treatment such as that described in Example 9 wherein the patient is diagnosed with metastatic or non-metastatic osteosarcoma.

EXAMPLE 51

5 A treatment such as that described in Example 9 wherein the patient is diagnosed with metastatic or non-metastatic ovarian cancer (and/or germ cell ovarian cancer).

EXAMPLE 52

10 A treatment such as that described in Example 9 wherein the patient is diagnosed with metastatic or non-metastatic pancreatic cancer.

EXAMPLE 53

15 A treatment such as that described in Example 9 wherein the patient is diagnosed with metastatic or non-metastatic penis cancer.

EXAMPLE 54

20 A treatment such as that described in Example 9 wherein the patient is diagnosed with metastatic or non-metastatic retinoblastoma.

EXAMPLE 55

25 A treatment such as that described in Example 9 wherein the patient is diagnosed with metastatic or non-metastatic skin cancer.

EXAMPLE 56

A treatment such as that described in Example 9 wherein the patient is diagnosed with metastatic or non-metastatic soft-tissue sarcoma.

EXAMPLE 57

A treatment such as that described in Example 9 wherein the patient is diagnosed with metastatic or non-metastatic squamous cell carcinoma.

5

EXAMPLE 58

A treatment such as that described in Example 9 wherein the patient is diagnosed with metastatic or non-metastatic stomach cancer.

EXAMPLE 59

10

A treatment such as that described in Example 9 wherein the patient is diagnosed with metastatic or non-metastatic testicular cancer.

EXAMPLE 60

15

A treatment such as that described in Example 9 wherein the patient is diagnosed with metastatic or non-metastatic thyroid cancer.

EXAMPLE 61

20

A treatment such as that described in Example 9 wherein the patient is diagnosed with metastatic or non-metastatic trophoblastic neoplasm.

EXAMPLE 62

25

A treatment such as that described in Example 9 wherein the patient is diagnosed with metastatic or non-metastatic uterine cancer.

EXAMPLE 63

A treatment such as that described in Example 9 wherein the patient is diagnosed with metastatic or non-metastatic vaginal cancer.

EXAMPLE 64

A treatment such as that described in Example 9 wherein the patient is diagnosed with metastatic or non-metastatic cancer of the vulva.

5

EXAMPLE 65

A treatment such as that described in Example 9 wherein the patient is diagnosed with metastatic or non-metastatic Wilm's tumor.

10

The invention being thus described, it will be obvious that the same may be varied in many ways. Such variations are not to be regarded as a departure from the spirit and scope of the invention and all such modification are intended to be included within the scope of the following claims.

What is claimed is:

1. A method of treating cancer in an animal which comprises: administering to an animal suffering from a cancer an effective amount of a NAALADase inhibitor.

5 2. The method of claim 1 wherein the NAALADase inhibitor is selected from the group consisting of: a glutamate-derived hydroxyphosphinyl derivative compound; an acidic peptide analog; a conformationally restricted glutamate mimic; and mixtures thereof.

10 3. The method of claim 2 wherein the acidic peptide analog is selected from the group consisting of: Asp-Glu, Glu-Glu, Gly-Glu, gamma-Glu-Glu, and Glu-Glu-Glu.

15 4. The method of claim 2 wherein the conformationally restricted glutamate mimic is selected from the group consisting of beta-NAAG and quisqualic acid.

5. The method of claim 2 wherein the NAALADase inhibitor is a glutamate-derived hydroxyphosphinyl derivative compound.

20 6. The method of claim 1 wherein the NAALADase inhibitor is administered in combination with an additional therapeutic agent selected from the group consisting of: therapeutic hormones, chemotherapeutic hormones, anti-angiogenesis agents, radiolabelled compounds, and mixtures thereof.


25 7. The method of claim 1 wherein the cancer is selected from the group consisting of: ACTH-producing tumors, acute lymphocytic leukemia, acute nonlymphocytic leukemia, cancer of the adrenal cortex, bladder cancer, brain cancer, breast

cancer, cervical cancer, chronic lymphocytic leukemia, chronic myelocytic leukemia, colorectal cancer, cutaneous T-cell lymphoma, endometrial cancer, esophageal cancer, Ewing's sarcoma, gallbladder cancer, hairy cell leukemia, head & neck cancer, Hodgkin's lymphoma, Kaposi's sarcoma, kidney cancer, liver cancer, lung cancer, lung cancer (small and/or non-small cell), malignant peritoneal effusion, malignant pleural effusion, melanoma, mesothelioma, multiple myeloma, neuroblastoma, non-Hodgkin's lymphoma, osteosarcoma, ovary cancer, ovary (germ cell) cancer, pancreatic cancer, penis cancer, prostate cancer, retinoblastoma, skin cancer, soft tissue sarcoma, squamous cell carcinomas, stomach cancer, testicular cancer, thyroid cancer, trophoblastic neoplasms, cancer of the uterus, vaginal cancer, cancer of the vulva, and Wilm's tumor.

8. The method of claim 1 wherein the cancer is prostatic adenocarcinoma.

9. The method of claim 1 wherein the cancer is selected from the group consisting of: brain cancer, cancer of the adrenal cortex, kidney cancer, and testicular cancer.

10. The method of claim 1 wherein the NAALADase inhibitor comprises a compound having the following formula:



I

25 wherein

R₁ is hydrogen, C₁-C₈ straight or branched chain alkyl, C₂-C₈ straight or branched chain alkenyl group, C₃-C₈ cycloalkyl, C₅-C₈ cycloalkenyl, or Ar₁;

X is CH₂, O, or NR₁, where R₁ is defined above; and

5 R₂ is C₁-C₈ straight or branched chain alkyl, C₂-C₈ straight or branched chain alkenyl group, C₃-C₈ cycloalkyl, C₅-C₈ cycloalkenyl, or Ar₁, wherein said alkyl, alkenyl, cycloalkyl, cycloalkenyl or aryl group may be optionally substituted with carboxylic acid,

10 wherein said alkyl, alkenyl, cycloalkyl, cycloalkenyl or aryl groups may be optionally substituted with C₁-C₈ cycloalkyl, C₃ or C₅ cycloalkyl, C₅-C₈ cycloalkenyl, C₁-C₄ alkyl, C₁-C₄ alkenyl, halo, hydroxy, carboxy, nitro, trifluoromethyl, C₁-C₈ straight or branched chain alkyl or alkenyl, C₁-C₄ alkoxy, C₁-C₄ alkenyloxy, phenoxy, benzyloxy, amino, or Ar₁, and where Ar₁ is selected from the group consisting of 1-naphthyl, 2-naphthyl, 2-indolyl, 3-indolyl, 4-indolyl, 2-furyl, 3-furyl, tetrahydrofuran-2-yl, 2-thienyl, 3-thienyl, 4-thienyl, 2-, 3-, or 4-pyridyl, or phenyl, having one to five substituents which are independently selected from the group consisting of hydrogen, halo, hydroxy, carboxy, nitro, trifluoromethyl, C₁-C₈ straight or branched alkyl or alkenyl, C₁-C₄ alkoxy or C₁-C₄ alkenyloxy, phenoxy, benzyloxy, and amino; or pharmaceutically acceptable salts, hydrates, or mixtures thereof.

11. The method of claim 10 wherein R₁ and R₂ are straight or branched aliphatic groups or carbocyclic groups and X is CH₂.

12. The method of claim 11 wherein the NAALADase inhibitor is selected from the group consisting of:

2-[[methylhydroxyphosphinyl]methyl]pentanedioic acid;

2-[[ethylhydroxyphosphinyl]methyl]pentanedioic acid;

5 2-[[propylhydroxyphosphinyl]methyl]pentanedioic acid;

2-[[butylhydroxyphosphinyl]methyl]pentanedioic acid;

2-[[cyclohexylhydroxyphosphinyl]methyl]pentanedioic acid;

2-[[[(cyclohexyl)methylhydroxyphosphinyl]methyl]pentanedioic acid;

10 2-[[phenylhydroxyphosphinyl]methyl]pentanedioic acid;

2-[[benzylhydroxyphosphinyl]methyl]pentanedioic acid;

2-[[phenylethylhydroxyphosphinyl]methyl]pentanedioic acid;

2-[[phenylpropylhydroxyphosphinyl]methyl]pentanedioic acid;

2-[[phenylbutylhydroxyphosphinyl]methyl]pentanedioic acid;

15 2-[[[(4-methylbenzyl)hydroxyphosphinyl]methyl]pentanedioic acid;

2-[[[(4-fluorobenzyl)hydroxyphosphinyl]methyl]pentanedioic acid;

2-[[[(2-fluorobenzyl)hydroxyphosphinyl]methyl]pentanedioic acid;

20 2-[[[(pentafluorobenzyl)hydroxyphosphinyl]methyl]pentanedioic acid;

2-[[[(methoxybenzyl)hydroxyphosphinyl]methyl]pentanedioic acid;

2-[[[(2, 3, 4-trimethoxyphenyl)hydroxyphosphinyl]methyl]

25 pentanedioic acid;

2-[[[(1-naphthyl)hydroxyphosphinyl]methyl]pentanedioic acid;

2-[[[(2-naphthyl)hydroxyphosphinyl]methyl]pentanedioic acid;

2-[[[(1-naphthyl)methylhydroxyphosphinyl]methyl]pentanedioic

acid;

2-[[[(2-naphthyl)methylhydroxyphosphinyl]methyl]pentanedioic
acid;

5 2-[[[(1-naphthyl)ethylhydroxyphosphinyl]methyl]pentanedioic
acid;

2-[[[(2-naphthyl)ethylhydroxyphosphinyl]methyl]pentanedioic
acid;

2-[[[(1-naphthyl)propylhydroxyphosphinyl]methyl]pentanedioic
acid;

10 2-[[[(2-naphthyl)propylhydroxyphosphinyl]methyl]pentanedioic
acid;

2-[[[(1-naphthyl)butylhydroxyphosphinyl]methyl]pentanedioic
acid;

15 2-[[[(2-naphthyl)butylhydroxyphosphinyl]methyl]pentanedioic
acid;

2-[[[(phenylprop-2-enyl)hydroxyphosphinyl]methyl]pentanedioic
acid;

2-[[[(2-fluorobenzyl)hydroxyphosphinyl]methyl]pentanedioic
acid;

20 2-[[[(hydroxy)phenylmethyl]hydroxyphosphinyl]methyl]
pentanedioic acid;

2-[[[(3-methylbenzyl)hydroxyphosphinyl]methyl]pentanedioic
acid;

25 2-[[[(4-fluorophenyl)hydroxyphosphinyl]methyl]pentanedioic
acid;

2-(phosphonomethyl)pentanedioic acid;

2-[[[(3-trifluoromethylbenzyl)hydroxyphosphinyl]methyl]
pentanedioic acid;

- 3-(methylhydroxyphosphinyl)-2-phenylpropanoic acid;
3-(ethylhydroxyphosphinyl)-2-phenylpropanoic acid;
3-(propylhydroxyphosphinyl)-2-phenylpropanoic acid;
3-(butylhydroxyphosphinyl)-2-phenylpropanoic acid;
5 3-(cyclohexylhydroxyphosphinyl)-2-phenylpropanoic acid;
3-((cyclohexyl)methylhydroxyphosphinyl)-2-phenylpropanoic acid;
3-(phenylhydroxyphosphinyl)-2-phenylpropanoic acid;
3-(benzylhydroxyphosphinyl)-2-phenylpropanoic acid;
10 3-(phenylethylhydroxyphosphinyl)-2-phenylpropanoic acid;
3-(phenylpropylhydroxyphosphinyl)-2-phenylpropanoic acid;
3-(phenylbutylhydroxyphosphinyl)-2-phenylpropanoic acid;
3-((2, 3, 4-trimethoxyphenyl)-3-hydroxyphosphinyl)-2-phenylpropanoic acid;
15 3-((1-naphthyl)hydroxyphosphinyl)-2-phenylpropanoic acid;
3-((2-naphthyl)hydroxyphosphinyl)-2-phenylpropanoic acid;
3-((1-naphthyl)methylhydroxyphosphinyl)-2-phenylpropanoic acid;
3-((2-naphthyl)methylhydroxyphosphinyl)-2-phenylpropanoic acid;
20 3-((1-naphthyl)ethylhydroxyphosphinyl)-2-phenylpropanoic acid;
3-((2-naphthyl)ethylhydroxyphosphinyl)-2-phenylpropanoic acid;
3-((1-naphthyl)propylhydroxyphosphinyl)-2-phenylpropanoic acid;
3-((2-naphthyl)propylhydroxyphosphinyl)-2-phenylpropanoic acid;
25 3-((1-naphthyl)butylhydroxyphosphinyl)-2-phenylpropanoic acid;
3-((2-naphthyl)butylhydroxyphosphinyl)-2-phenylpropanoic acid;

- 3-(phenylprop-2-enylhydroxyphosphinyl)-2-phenylpropanoic acid;
2-[(benzylhydroxyphosphinyl)methyl]pentanedioic acid;
2-[(methylhydroxyphosphinyl)methyl]hexanedioic acid;
2-[(benzylhydroxyphosphinyl)methyl]hexanedioic acid;
5 2-[(methylhydroxyphosphinyl)methyl]heptanedioic acid;
2-[(benzylhydroxyphosphinyl)methyl]heptanedioic acid;
2-[(methylhydroxyphosphinyl)methyl]octanedioic acid;
2-[(benzylhydroxyphosphinyl)methyl]octanedioic acid;
2-[(methylhydroxyphosphinyl)methyl]nonanedioic acid;
10 2-[(benzylhydroxyphosphinyl)methyl]nonanedioic acid;
2-[(methylhydroxyphosphinyl)methyl]decanedioic acid;
2-[(benzylhydroxyphosphinyl)methyl]decanedioic acid;
3-(benzylhydroxyphosphinyl)-2-methylpropanoic acid;
3-(benzylhydroxyphosphinyl)-2-ethylpropanoic acid;
15 3-(benzylhydroxyphosphinyl)-2-propylpropanoic acid;
3-(benzylhydroxyphosphinyl)-2-butylpropanoic acid;
3-(benzylhydroxyphosphinyl)-2-cyclohexylpropanoic acid;
3-(benzylhydroxyphosphinyl)-2-(cyclohexyl)methylpropanoic
acid;
20 3-(benzylhydroxyphosphinyl)-2-phenylpropanoic acid;
3-(benzylhydroxyphosphinyl)-2-benzylpropanoic acid;
3-(benzylhydroxyphosphinyl)-2-phenylethylpropanoic acid;
3-(benzylhydroxyphosphinyl)-2-phenylpropylpropanoic acid;
3-(benzylhydroxyphosphinyl)-2-phenylbutylpropanoic acid;
25 3-(benzylhydroxyphosphinyl)-2-(2, 3, 4-trimethoxyphenyl)
propanoic acid;
3-(benzylhydroxyphosphinyl)-2-(1-naphthyl)propanoic acid;
3-(benzylhydroxyphosphinyl)-2-(2-naphthyl)propanoic acid;

3-(benzylhydroxyphosphinyl)-2-(1-naphthyl)methylpropanoic acid;

3-(benzylhydroxyphosphinyl)-2-(2-naphthyl)methylpropanoic acid;

5 3-(benzylhydroxyphosphinyl)-2-(1-naphthyl)ethylpropanoic acid;

3-(benzylhydroxyphosphinyl)-2-(2-naphthyl)ethylpropanoic acid;

3-(benzylhydroxyphosphinyl)-2-(1-naphthyl)propylpropanoic acid;

10 3-(benzylhydroxyphosphinyl)-2-(2-naphthyl)propylpropanoic acid;

3-(benzylhydroxyphosphinyl)-2-(1-naphthyl)butylpropanoic acid;

3-(benzylhydroxyphosphinyl)-2-(2-naphthyl)butylpropanoic acid;

3-(benzylhydroxyphosphinyl)-2-phenylprop-2-enylpropanoic acid;

15 2-[[(2-pyridyl)methylhydroxyphosphinyl]methyl]pentanedioic acid;

2-[[(3-pyridyl)methylhydroxyphosphinyl]methyl]pentanedioic acid;

2-[[(4-pyridyl)methylhydroxyphosphinyl]methyl]pentanedioic acid;

20 2-[[(3-pyridyl)ethylhydroxyphosphinyl]methyl]pentanedioic acid;

2-[[(3-pyridyl)propylhydroxyphosphinyl]methyl]pentanedioic acid;

25 2-[[(tetrahydrofuranyl)methylhydroxyphosphinyl]methyl]pentanedioic acid;

2-[[(tetrahydrofuranyl)ethylhydroxyphosphinyl]methyl]pentanedioic acid;

2-[[(tetrahydrofuranyl)propylhydroxyphosphinyl]methyl]

pentanedioic acid;

2-[[[(2-indolyl)methylhydroxyphosphinyl]methyl]pentanedioic acid;

5 2-[[[(3-indolyl)methylhydroxyphosphinyl]methyl]pentanedioic acid;

2-[[[(4-indolyl)methylhydroxyphosphinyl]methyl]pentanedioic acid;

2-[[[(3-indolyl)ethylhydroxyphosphinyl]methyl]pentanedioic acid;

10 2-[[[(3-indolyl)propylhydroxyphosphinyl]methyl]pentanedioic acid;

2-[[[(2-thienyl)methylhydroxyphosphinyl]methyl]pentanedioic acid;

15 2-[[[(3-thienyl)methylhydroxyphosphinyl]methyl]pentanedioic acid;

2-[[[(4-thienyl)methylhydroxyphosphinyl]methyl]pentanedioic acid;

2-[[[(3-thienyl)ethylhydroxyphosphinyl]methyl]pentanedioic acid;

20 2-[[[(3-thienyl)propylhydroxyphosphinyl]methyl]pentanedioic acid;

3-[(2-pyridyl)methylhydroxyphosphinyl]-2-phenylpropanoic acid;

3-[(3-pyridyl)methylhydroxyphosphinyl]-2-phenylpropanoic acid;

3-[(4-pyridyl)methylhydroxyphosphinyl]-2-phenylpropanoic acid;

25 3-[(3-pyridyl)ethylhydroxyphosphinyl]-2-phenylpropanoic acid;

3-[(3-pyridyl)propylhydroxyphosphinyl]-2-phenylpropanoic acid;

3-[(tetrahydrofuranyl)methylhydroxyphosphinyl]-2-phenyl propanoic acid;

- 3-[(tetrahydrofuranyl)ethylhydroxyphosphinyl]-2-phenyl
propanoic acid;
- 3-[(tetrahydrofuranyl)propylhydroxyphosphinyl]-2-phenyl
propanoic acid;
- 5 3-[(2-indolyl)methylhydroxyphosphinyl]-2-phenylpropanoic acid;
3-[(3-indolyl)methylhydroxyphosphinyl]-2-phenylpropanoic acid;
3-[(4-indolyl)methylhydroxyphosphinyl]-2-phenylpropanoic acid;
3-[(3-indolyl)ethylhydroxyphosphinyl]-2-phenylpropanoic acid;
3-[(3-indolyl)propylhydroxyphosphinyl]-2-phenylpropanoic acid;
- 10 3-[(2-thienyl)methylhydroxyphosphinyl]-2-phenylpropanoic acid;
3-[(3-thienyl)methylhydroxyphosphinyl]-2-phenylpropanoic acid;
3-[(4-thienyl)methylhydroxyphosphinyl]-2-phenylpropanoic acid;
3-[(3-thienyl)ethylhydroxyphosphinyl]-2-phenylpropanoic acid;
3-[(3-thienyl)propylhydroxyphosphinyl]-2-phenylpropanoic acid;
- 15 3-(benzylhydroxyphosphinyl)-2-(2-pyridyl)methylpropanoic acid;
3-(benzylhydroxyphosphinyl)-2-(3-pyridyl)methylpropanoic acid;
3-(benzylhydroxyphosphinyl)-2-(4-pyridyl)methylpropanoic acid;
3-(benzylhydroxyphosphinyl)-2-(3-pyridyl)ethylpropanoic acid;
3-(benzylhydroxyphosphinyl)-2-(3-pyridyl)propylpropanoic acid;
- 20 3-(benzylhydroxyphosphinyl)-2-(tetrahydrofuranyl)methyl
propanoic acid;
3-(benzylhydroxyphosphinyl)-2-(tetrahydrofuranyl)ethyl
propanoic acid;
3-(benzylhydroxyphosphinyl)-2-(tetrahydrofuranyl)propyl
- 25 propanoic acid;
3-(benzylhydroxyphosphinyl)-2-(2-indolyl)methylpropanoic acid;
3-(benzylhydroxyphosphinyl)-2-(3-indolyl)methylpropanoic acid;
3-(benzylhydroxyphosphinyl)-2-(4-indolyl)methylpropanoic acid;

3-(benzylhydroxyphosphinyl)-2-(3-indolyl)ethylpropanoic acid;
3-(benzylhydroxyphosphinyl)-2-(3-indolyl)propylpropanoic acid;
3-(benzylhydroxyphosphinyl)-2-(2-thienyl)methylpropanoic acid;
3-(benzylhydroxyphosphinyl)-2-(3-thienyl)methylpropanoic acid;
5 3-(benzylhydroxyphosphinyl)-2-(4-thienyl)methylpropanoic acid;
3-(benzylhydroxyphosphinyl)-2-(3-thienyl)ethylpropanoic acid;
3-(benzylhydroxyphosphinyl)-2-(3-thienyl)propylpropanoic acid;
and a pharmaceutically acceptable salt, hydrate, or a mixture thereof.

10 13. The method of claim 11 wherein the NAALADase inhibitor is selected from the group consisting of:
2-[(benzylhydroxyphosphinyl)methyl]pentanedioic acid;
2-[(phenylhydroxyphosphinyl)methyl]pentanedioic acid;
2-[[[(hydroxy)phenylmethyl]hydroxyphosphinyl]methyl]
15 pentanedioic acid;
2-[(butylhydroxyphosphinyl)methyl]pentanedioic acid;
2-[[[(3-methylbenzyl)hydroxyphosphinyl]methyl]pentanedioic acid;
2-[(3-phenylpropylhydroxyphosphinyl)methyl]pentanedioic acid;
20 2-[[[(4-fluorophenyl)hydroxyphosphinyl]methyl]pentanedioic acid;
2-[(methylhydroxyphosphinyl)methyl]pentanedioic acid;
2-[(phenylethylhydroxyphosphinyl)methyl]pentanedioic acid;
2-[[[(4-methylbenzyl)hydroxyphosphinyl]methyl]pentanedioic
25 acid;
2-[[[(4-fluorobenzyl)hydroxyphosphinyl]methyl]pentanedioic acid;
2-[[[(4-methoxybenzyl)hydroxyphosphinyl]methyl]pentanedioic

acid;

2-(phosphonomethyl)pentanedioic acid;

2-[[[(3-trifluoromethylbenzyl)hydroxyphosphinyl]methyl]

pentanedioic acid;

5 2-[[[(2-fluorobenzyl)hydroxyphosphinyl]methyl]pentanedioic acid;

2-[[[(pentafluorobenzyl)hydroxyphosphinyl]methyl]pentanedioic acid; and a pharmaceutically acceptable salt, hydrate, or a mixture thereof.

10 14. The method of claim 10 wherein R₁ and R₂ are straight or branched aliphatic groups or carbocyclic groups and X is oxygen.

15 15. The method of claim 14 wherein the NAALADase inhibitor is selected from the group consisting of:

2-[[methylhydroxyphosphinyl]oxy]pentanedioic acid;

2-[[ethylhydroxyphosphinyl]oxy]pentanedioic acid;

2-[[propylhydroxyphosphinyl]oxy]pentanedioic acid;

2-[[butylhydroxyphosphinyl]oxy]pentanedioic acid;

2-[[cyclohexylhydroxyphosphinyl]oxy]pentanedioic acid;

20 2-[[[(cyclohexyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;

2-[[phenylhydroxyphosphinyl]oxy]pentanedioic acid;

2-[[benzylhydroxyphosphinyl]oxy]pentanedioic acid;

2-[[phenylethylhydroxyphosphinyl]oxy]pentanedioic acid;

2-[[phenylpropylhydroxyphosphinyl]oxy]pentanedioic acid;

25 2-[[phenylbutylhydroxyphosphinyl]oxy]pentanedioic acid;

2-[[[(4-methylbenzyl)hydroxyphosphinyl]oxy]pentanedioic acid;

2-[[[(4-fluorobenzyl)hydroxyphosphinyl]oxy]pentanedioic acid;

2-[[[(2-fluorobenzyl)hydroxyphosphinyl]oxy]pentanedioic acid;

2-[[[(pentafluorobenzyl)hydroxyphosphinyl]oxy] pentanedioic acid;

2-[[[(methoxybenzyl)hydroxyphosphinyl]oxy]pentanedioic acid;

2-[[[(2, 3, 4-trimethoxyphenyl)hydroxyphosphinyl]oxy]

5 pentanedioic acid;

2-[[[(1-naphthyl)hydroxyphosphinyl]oxy]pentanedioic acid;

2-[[[(2-naphthyl)hydroxyphosphinyl]oxy]pentanedioic acid;

2-[[[(1-naphthyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;

2-[[[(2-naphthyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;

10 2-[[[(1-naphthyl)ethylhydroxyphosphinyl]oxy]pentanedioic acid;

2-[[[(2-naphthyl)ethylhydroxyphosphinyl]oxy]pentanedioic acid;

2-[[[(1-naphthyl)propylhydroxyphosphinyl]oxy]pentanedioic acid;

2-[[[(2-naphthyl)propylhydroxyphosphinyl]oxy]pentanedioic acid;

2-[[[(1-naphthyl)butylhydroxyphosphinyl]oxy]pentanedioic acid;

15 2-[[[(2-naphthyl)butylhydroxyphosphinyl]oxy]pentanedioic acid;

2-[[[(phenylprop-2-enyl)hydroxyphosphinyl]oxy]pentanedioic acid;

2-[[[benzylhydroxyphosphinyl]oxy]pentanedioic acid;

20 2-[[[(hydroxy)phenylmethyl]hydroxyphosphinyl]oxy]pentanedioic acid;

2-[[[(3-methylbenzyl)hydroxyphosphinyl]oxy]pentanedioic acid;

2-[[[(4-fluorophenyl)hydroxyphosphinyl]oxy]pentanedioic acid;

2-[[[(2-fluorobenzyl)hydroxyphosphinyl]oxy]pentanedioic acid;

2-(phosphono)oxy]pentanedioic acid;

25 2-[[[(3-trifluoromethylbenzyl)hydroxyphosphinyl]oxy]pentanedioic acid;

2-[[[methylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;

2-[[[ethylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;

- 2-[[propylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
2-[[butylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
2-[[cyclohexylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
2-[[[(cyclohexyl)methylhydroxyphosphinyl]oxy]-2-phenylethanoic
5 acid;
2-[[phenylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
2-[[phenylethylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
2-[[phenylpropylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
10 2-[[phenylbutylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
2-[[[(2, 3, 4-trimethoxyphenyl)-3-hydroxyphosphinyl]oxy]-2-
phenylethanoic acid;
2-[[[(1-naphthyl)hydroxyphosphinyl]oxy]-2-phenylethanoic acid;
2-[[[(2-naphthyl)hydroxyphosphinyl]oxy]-2-phenylethanoic acid;
15 2-[[[(1-naphthyl)methylhydroxyphosphinyl]oxy]-2-phenylethanoic
acid;
2-[[[(2-naphthyl)methylhydroxyphosphinyl]oxy]-2-phenylethanoic
acid;
2-[[[(1-naphthyl)ethylhydroxyphosphinyl]oxy]-2-phenylethanoic
20 acid;
2-[[[(2-naphthyl)ethylhydroxyphosphinyl]oxy]-2-phenylethanoic
acid;
2-[[[(1-naphthyl)propylhydroxyphosphinyl]oxy]-2-phenylethanoic
acid;
25 2-[[[(2-naphthyl)propylhydroxyphosphinyl]oxy]-2-phenylethanoic
acid;
2-[[[(1-naphthyl)butylhydroxyphosphinyl]oxy]-2-phenylethanoic
acid;

2-[[(2-naphthyl)butylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;

2-[[phenylprop-2-enylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;

5 2-[(methylhydroxyphosphinyl)oxy]hexanedioic acid;

2-[(benzylhydroxyphosphinyl)oxy]hexanedioic acid;

2-[(methylhydroxyphosphinyl)oxy]heptanedioic acid;

2-[(benzylhydroxyphosphinyl)oxy]heptanedioic acid;

2-[(methylhydroxyphosphinyl)oxy]octanedioic acid;

10 2-[(benzylhydroxyphosphinyl)oxy]octanedioic acid;

2-[(methylhydroxyphosphinyl)oxy]nonanedioic acid;

2-[(benzylhydroxyphosphinyl)oxy]nonanedioic acid;

2-[(methylhydroxyphosphinyl)oxy]decanedioic acid;

2-[(benzylhydroxyphosphinyl)oxy]decanedioic acid;

15 2-[[benzylhydroxyphosphinyl]oxy]-2-methylethanoic acid;

2-[[benzylhydroxyphosphinyl]oxy]-2-ethylethanoic acid;

2-[[benzylhydroxyphosphinyl]oxy]-2-propylethanoic acid;

2-[[benzylhydroxyphosphinyl]oxy]-2-butylethanoic acid;

2-[[benzylhydroxyphosphinyl]oxy]-2-cyclohexylethanoic acid;

20 2-[[benzylhydroxyphosphinyl]oxy]-2-(cyclohexyl)methylethanoic acid;

2-[[benzylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;

2-[[benzylhydroxyphosphinyl]oxy]-2-benzylethanoic acid;

2-[[benzylhydroxyphosphinyl]oxy]-2-phenylethylethanoic acid;

25 2-[[benzylhydroxyphosphinyl]oxy]-2-phenylpropylethanoic acid;

2-[[benzylhydroxyphosphinyl]oxy]-2-phenylbutylethanoic acid;

2-[[benzylhydroxyphosphinyl]oxy]-2-(2, 3, 4-trimethoxyphenyl)ethanoic acid;

- 2-[[benzylhydroxyphosphinyl]oxy]-2-(1-naphthyl)ethanoic acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-(2-naphthyl)ethanoic acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-(1-naphthyl)methylethanoic
acid;
5 2-[[benzylhydroxyphosphinyl]oxy]-2-(2-naphthyl)methylethanoic
acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-(1-naphthyl)ethylethanoic
acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-(2-naphthyl)ethylethanoic
10 acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-(1-naphthyl)propylethanoic
acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-(2-naphthyl)propylethanoic
acid;
15 2-[[benzylhydroxyphosphinyl]oxy]-2-(1-naphthyl)butylethanoic
acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-(2-naphthyl)butylethanoic
acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-phenylprop-2-enylethanoic
20 acid;
2-[[[(2-pyridyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(3-pyridyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(4-pyridyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(3-pyridyl)ethylhydroxyphosphinyl]oxy]pentanedioic acid;
25 2-[[[(3-pyridyl)propylhydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(tetrahydrofuranyl)methylhydroxyphosphinyl]oxy]
pentanedioic acid;
2-[[[(tetrahydrofuranyl)ethylhydroxyphosphinyl]oxy]pentanedioic

acid;

2-[[[(tetrahydrofuranyl)propylhydroxyphosphinyl]oxy]

pentanedioic acid;

2-[[[(2-indolyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;

5 2-[[[(3-indolyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;

2-[[[(4-indolyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;

2-[[[(3-indolyl)ethylhydroxyphosphinyl]oxy]pentanedioic acid;

2-[[[(3-indolyl)propylhydroxyphosphinyl]oxy]pentanedioic acid;

2-[[[(2-thienyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;

10 2-[[[(3-thienyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;

2-[[[(4-thienyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;

2-[[[(3-thienyl)ethylhydroxyphosphinyl]oxy]pentanedioic acid;

2-[[[(3-thienyl)propylhydroxyphosphinyl]oxy]pentanedioic acid;

2-[[[(2-pyridyl)methylhydroxyphosphinyl]oxy]-2-phenylethanoic

15 acid;

2-[[[(3-pyridyl)methylhydroxyphosphinyl]oxy]-2-phenylethanoic
acid;

2-[[[(4-pyridyl)methylhydroxyphosphinyl]oxy]-2-phenylethanoic
acid;

20 2-[[[(3-pyridyl)ethylhydroxyphosphinyl]oxy]-2-phenylethanoic
acid;

2-[[[(3-pyridyl)propylhydroxyphosphinyl]oxy]-2-phenylethanoic
acid;

2-[[[(tetrahydrofuranyl)methylhydroxyphosphinyl]oxy]-2-phenyl
25 ethanoic acid;

2-[[[(tetrahydrofuranyl)ethylhydroxyphosphinyl]oxy]-2-phenyl
ethanoic acid;

2-[[[(tetrahydrofuranyl)propylhydroxyphosphinyl]oxy]-2-phenyl

ethanoic acid;

2-[[[(2-indolyl)methylhydroxyphosphinyl]oxy]-2-phenyl ethanoic acid;

2-[[[(3-indolyl)methylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;

2-[[[(4-indolyl)methylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;

2-[[[(3-indolyl)ethylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;

2-[[[(3-indolyl)propylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;

2-[[[(2-thienyl)methylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;

2-[[[(3-thienyl)methylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;

2-[[[(4-thienyl)methylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;

2-[[[(3-thienyl)ethylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;

2-[[[(3-thienyl)propylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;

2-[[benzylhydroxyphosphinyl]oxy]-2-(2-pyridyl)methylethanoic acid;

2-[[benzylhydroxyphosphinyl]oxy]-2-(3-pyridyl)methylethanoic acid;

2-[[benzylhydroxyphosphinyl]oxy]-2-(4-pyridyl)methylethanoic acid;

2-[[benzylhydroxyphosphinyl]oxy]-2-(3-pyridyl)ethylethanoic

acid;

2-[[benzylhydroxyphosphinyl]oxy]-2-(3-pyridyl)propylethanoic
acid;

2-[[benzylhydroxyphosphinyl]oxy]-2-(tetrahydrofuran-2-yl)

5 methylethanoic acid;

2-[[benzylhydroxyphosphinyl]oxy]-2-(tetrahydrofuran-2-yl)
ethylethanoic acid;

2-[[benzylhydroxyphosphinyl]oxy]-2-(tetrahydrofuran-2-yl)
propylethanoic acid;

10 2-[[benzylhydroxyphosphinyl]oxy]-2-(2-indolyl)methylethanoic
acid;

2-[[benzylhydroxyphosphinyl]oxy]-2-(3-indolyl)methylethanoic
acid;

2-[[benzylhydroxyphosphinyl]oxy]-2-(4-indolyl)methylethanoic
15 acid;

2-[[benzylhydroxyphosphinyl]oxy]-2-(3-indolyl)ethylethanoic
acid;

2-[[benzylhydroxyphosphinyl]oxy]-2-(3-indolyl)propylethanoic
acid;

20 2-[[benzylhydroxyphosphinyl]oxy]-2-(2-thienyl)methylethanoic
acid;

2-[[benzylhydroxyphosphinyl]oxy]-2-(3-thienyl)methylethanoic
acid;

2-[[benzylhydroxyphosphinyl]oxy]-2-(4-thienyl)methylethanoic
25 acid;

2-[[benzylhydroxyphosphinyl]oxy]-2-(3-thienyl)ethylethanoic
acid;

2-[[benzylhydroxyphosphinyl]oxy]-2-(3-thienyl)propylethanoic

acid; and a pharmaceutically acceptable salt, hydrate, or a mixture thereof.

16. The method of claim 14 wherein the NAALADase inhibitor is selected from the group consisting of:

- 5 2-[(benzylhydroxyphosphinyl)oxy]pentanedioic acid;
2-[(phenylhydroxyphosphinyl)oxy]pentanedioic acid;
2-[[[(hydroxy)phenylmethyl]hydroxyphosphinyl]oxy]pentanedioic acid;
2-[(butylhydroxyphosphinyl)oxy]pentanedioic acid;
10 2-[[[(3-methylbenzyl)hydroxyphosphinyl]oxy]pentanedioic acid;
2-[(3-phenylpropylhydroxyphosphinyl)oxy]pentanedioic acid;
2-[[[(4-fluorophenyl)hydroxyphosphinyl]oxy]pentanedioic acid;
2-[(methylhydroxyphosphinyl)oxy]pentanedioic acid;
2-[(phenylethylhydroxyphosphinyl)oxy]pentanedioic acid;
15 2-[[[(4-methylbenzyl)hydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(4-fluorobenzyl)hydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(4-methoxybenzyl)hydroxyphosphinyl]oxy]pentanedioic acid;
2-[(phosphono)oxy]pentanedioic acid;
2-[[[(3-trifluoromethylbenzyl)hydroxyphosphinyl]oxy]
20 pentanedioic acid;
2-[[[(2-fluorobenzyl)hydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(pentafluorobenzyl)hydroxyphosphinyl]oxy]pentanedioic acid; and a pharmaceutically acceptable salt, hydrate, or a mixture thereof.

- 25 17. The method of claim 10 wherein R₁ and R₂ are straight or branched aliphatic groups or carbocyclic groups and X is NR₃.

18. The method of claim 17 wherein the NAALADase

inhibitor is selected from the group consisting of:

2-[[methylhydroxyphosphinyl]amino]pentanedioic acid;

2-[[ethylhydroxyphosphinyl]amino]pentanedioic acid;

2-[[propylhydroxyphosphinyl]amino]pentanedioic acid;

5 2-[[butylhydroxyphosphinyl]amino]pentanedioic acid;

2-[[cyclohexylhydroxyphosphinyl]amino]pentanedioic acid;

2-[[[(cyclohexyl)methylhydroxyphosphinyl]amino]pentanedioic acid;

2-[[phenylhydroxyphosphinyl]amino]pentanedioic acid;

10 2-[[benzylhydroxyphosphinyl]amino]pentanedioic acid;

2-[[phenylethylhydroxyphosphinyl]amino]pentanedioic acid;

2-[[phenylpropylhydroxyphosphinyl]amino]pentanedioic acid;

2-[[phenylbutylhydroxyphosphinyl]amino]pentanedioic acid;

2-[[[(4-methylbenzyl)hydroxyphosphinyl]amino]pentanedioic acid;

15 2-[[[(4-fluorobenzyl)hydroxyphosphinyl]amino]pentanedioic acid;

2-[[[(2-fluorobenzyl)hydroxyphosphinyl]amino]pentanedioic acid;

2-[[[(pentafluorobenzyl)hydroxyphosphinyl]amino]pentanedioic acid;

2-[[[(methoxybenzyl)hydroxyphosphinyl]amino]pentanedioic acid;

20 2-[[[(2, 3, 4-trimethoxyphenyl)hydroxyphosphinyl]amino]pentanedioic acid;

2-[[[(1-naphthyl)hydroxyphosphinyl]amino]pentanedioic acid;

2-[[[(2-naphthyl)hydroxyphosphinyl]amino]pentanedioic acid;

2-[[[(1-naphthyl)methylhydroxyphosphinyl]amino]pentanedioic acid;

25 2-[[[(2-naphthyl)methylhydroxyphosphinyl]amino]pentanedioic acid;

2-[[[(1-naphthyl)ethylhydroxyphosphinyl]amino]pentanedioic

acid;
2-[[[(2-naphthyl)ethylhydroxyphosphinyl]amino]pentanedioic
acid;
2-[[[(1-naphthyl)propylhydroxyphosphinyl]amino]pentanedioic
5 acid;
2-[[[(2-naphthyl)propylhydroxyphosphinyl]amino]pentanedioic
acid;
2-[[[(1-naphthyl)butylhydroxyphosphinyl]amino]pentanedioic
acid;
10 2-[[[(2-naphthyl)butylhydroxyphosphinyl]amino]pentanedioic
acid;
2-[[[(phenylprop-2-enyl)hydroxyphosphinyl]amino]pentanedioic
acid;
2-[[benzylhydroxyphosphinyl]amino]pentanedioic acid;
15 2-[[[(2-fluorobenzyl)hydroxyphosphinyl]amino]-2-pentanedioic
acid;
2-[[[(hydroxy)phenylmethyl]hydroxyphosphinyl]amino]
pentanedioic acid;
2-[[[(3-methylbenzyl)hydroxyphosphinyl]amino]pentanedioic acid;
20 2-[[[(4-fluorophenyl)hydroxyphosphinyl]amino]pentanedioic acid;
2-[(phosphono)amino]pentanedioic acid;
2-[[[(3-trifluoromethylbenzyl)hydroxyphosphinyl]amino]
pentanedioic acid;
2-[[methylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
25 2-[[ethylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
2-[[propylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
2-[[butylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
2-[[cyclohexylhydroxyphosphinyl]amino]-2-phenylethanoic acid;

2-[[[(cyclohexyl)methylhydroxyphosphinyl]amino]-2-phenyl
ethanoic acid;

2-[[[phenylhydroxyphosphinyl]amino]-2-phenylethanoic acid;

2-[[[benzylhydroxyphosphinyl]amino]-2-phenylethanoic acid;

5 2-[[[phenylethylhydroxyphosphinyl]amino]-2-phenylethanoic acid;

2-[[[phenylpropylhydroxyphosphinyl]amino]-2-phenylethanoic
acid;

2-[[[phenylbutylhydroxyphosphinyl]amino]-2-phenylethanoic acid;

10 2-[[[(2, 3, 4-trimethoxyphenyl)-3-hydroxyphosphinyl]amino]-2-
phenylethanoic acid;

2-[[[(1-naphthyl)hydroxyphosphinyl]amino]-2-phenylethanoic
acid;

2-[[[(2-naphthyl)hydroxyphosphinyl]amino]-2-phenylethanoic
acid;

15 2-[[[(1-naphthyl)methylhydroxyphosphinyl]amino]-2-phenyl
ethanoic acid;

2-[[[(2-naphthyl)methylhydroxyphosphinyl]amino]-2-phenyl
ethanoic acid;

2-[[[(1-naphthyl)ethylhydroxyphosphinyl]amino]-2-phenylethanoic
20 acid;

2-[[[(2-naphthyl)ethylhydroxyphosphinyl]amino]-2-phenylethanoic
acid;

2-[[[(1-naphthyl)propylhydroxyphosphinyl]amino]-2-phenyl
ethanoic acid;

25 2-[[[(2-naphthyl)propylhydroxyphosphinyl]amino]-2-phenyl
ethanoic acid;

2-[[[(1-naphthyl)butylhydroxyphosphinyl]amino]-2-phenylethanoic
acid;

2-[[[(2-naphthyl)butylhydroxyphosphinyl]amino]-2-phenylethanoic acid;

2-[[[phenylprop-2-enylhydroxyphosphinyl]amino]-2-phenylethanoic acid;

5 2-[[[benzylhydroxyphosphinyl]amino]-2-methylethanoic acid;

2-[[[benzylhydroxyphosphinyl]amino]-2-ethylethanoic acid;

2-[[[benzylhydroxyphosphinyl]amino]-2-propylethanoic acid;

2-[[[benzylhydroxyphosphinyl]amino]-2-butylethanoic acid;

2-[[[benzylhydroxyphosphinyl]amino]-2-cyclohexylethanoic acid;

10 2-[[[benzylhydroxyphosphinyl]amino]-2-(cyclohexyl)methyl ethanoic acid;

2-[[[benzylhydroxyphosphinyl]amino]-2-phenylethanoic acid;

2-[[[benzylhydroxyphosphinyl]amino]-2-benzylethanoic acid;

2-[[[benzylhydroxyphosphinyl]amino]-2-phenylethylethanoic acid;

15 2-[[[benzylhydroxyphosphinyl]amino]-2-phenylpropylethanoic acid;

2-[[[benzylhydroxyphosphinyl]amino]-2-phenylbutylethanoic acid;

2-[[[benzylhydroxyphosphinyl]amino]-2-(2, 3, 4-trimethoxyphenyl)ethanoic acid;

20 2-[[[benzylhydroxyphosphinyl]amino]-2-(1-naphthyl)ethanoic acid;

2-[[[benzylhydroxyphosphinyl]amino]-2-(2-naphthyl)ethanoic acid;

2-[[[benzylhydroxyphosphinyl]amino]-2-(1-naphthyl)methyl ethanoic acid;

25 2-[[[benzylhydroxyphosphinyl]amino]-2-(2-naphthyl)methyl ethanoic acid;

2-[[[benzylhydroxyphosphinyl]amino]-2-(1-naphthyl)ethylethanoic

acid;

2-[[benzylhydroxyphosphinyl]amino]-2-(2-naphthyl)ethylethanoic
acid;

2-[[benzylhydroxyphosphinyl]amino]-2-(1-naphthyl)propyl

5 ethanoic acid;

2-[[benzylhydroxyphosphinyl]amino]-2-(2-naphthyl)propyl
ethanoic acid;

2-[[benzylhydroxyphosphinyl]amino]-2-(1-naphthyl)butylethanoic
acid;

10 2-[[benzylhydroxyphosphinyl]amino]-2-(2-naphthyl)butylethanoic
acid;

2-[[benzylhydroxyphosphinyl]amino]-2-phenolprop-2-enylethanoic
acid;

2-[(methylhydroxyphosphinyl)amino]hexanedioic acid;

15 2-[(benzylhydroxyphosphinyl)amino]hexanedioic acid;

2-[(methylhydroxyphosphinyl)amino]heptanedioic acid;

2-[(benzylhydroxyphosphinyl)amino]heptanedioic acid;

2-[(methylhydroxyphosphinyl)amino]octanedioic acid;

2-[(benzylhydroxyphosphinyl)amino]octanedioic acid;

20 2-[(methylhydroxyphosphinyl)amino]nonanedioic acid;

2-[(benzylhydroxyphosphinyl)amino]nonanedioic acid;

2-[(methylhydroxyphosphinyl)amino]decanedioic acid;

2-[(benzylhydroxyphosphinyl)amino]decanedioic acid;

3-[(2-pyridyl)methylhydroxyphosphinyl]amino]pentanedioic
25 acid;

3-[(3-pyridyl)methylhydroxyphosphinyl]amino]pentanedioic
acid;

3-[(4-pyridyl)methylhydroxyphosphinyl]amino]pentanedioic

acid;

3-[[[(3-pyridyl)ethylhydroxyphosphinyl]amino]pentanedioic acid;

3-[[[(3-pyridyl)propylhydroxyphosphinyl]amino]pentanedioic
acid;

5 3-[[[(tetrahydrofuranyl)methylhydroxyphosphinyl]amino]
pentanedioic acid;

3-[[[(tetrahydrofuranyl)ethylhydroxyphosphinyl]amino]
pentanedioic acid;

10 3-[[[(tetrahydrofuranyl)propylhydroxyphosphinyl]amino]
pentanedioic acid;

3-[[[(2-indolyl)methylhydroxyphosphinyl]amino]pentanedioic
acid;

3-[[[(3-indolyl)methylhydroxyphosphinyl]amino]pentanedioic
acid;

15 3-[[[(4-indolyl)methylhydroxyphosphinyl]amino]pentanedioic
acid;

3-[[[(3-indolyl)ethylhydroxyphosphinyl]amino]pentanedioic acid;

3-[[[(3-indolyl)propylhydroxyphosphinyl]amino]pentanedioic
acid;

20 3-[[[(2-thienyl)methylhydroxyphosphinyl]amino]pentanedioic
acid;

3-[[[(3-thienyl)methylhydroxyphosphinyl]amino]pentanedioic
acid;

25 3-[[[(4-thienyl)methylhydroxyphosphinyl]amino]pentanedioic
acid;

3-[[[(3-thienyl)ethylhydroxyphosphinyl]amino]pentanedioic acid;

3-[[[(3-thienyl)propylhydroxyphosphinyl]amino]pentanedioic
acid;

2-[[(2-pyridyl)methylhydroxyphosphinyl]amino]-2-phenylethanoic acid;

2-[[(3-pyridyl)methylhydroxyphosphinyl]amino]-2-phenylethanoic acid;

5 2-[[(4-pyridyl)methylhydroxyphosphinyl]amino]-2-phenylethanoic acid;

2-[[(3-pyridyl)ethylhydroxyphosphinyl]amino]-2-phenylethanoic acid;

10 2-[[(3-pyridyl)propylhydroxyphosphinyl]amino]-2-phenylethanoic acid;

2-[[(tetrahydrofuranyl)methylhydroxyphosphinyl]amino]-2-phenyl ethanoic acid;

2-[[(tetrahydrofuranyl)ethylhydroxyphosphinyl]amino]-2-phenyl ethanoic acid;

15 2-[[(tetrahydrofuranyl)propylhydroxyphosphinyl]amino]-2-phenyl ethanoic acid;

2-[[(2-indolyl)methylhydroxyphosphinyl]amino]-2-phenylethanoic acid;

20 2-[[(3-indolyl)methylhydroxyphosphinyl]amino]-2-phenylethanoic acid;

2-[[(4-indolyl)methylhydroxyphosphinyl]amino]-2-phenylethanoic acid;

2-[[(3-indolyl)ethylhydroxyphosphinyl]amino]-2-phenylethanoic acid;

25 2-[[(3-indolyl)propylhydroxyphosphinyl]amino]-2-phenylethanoic acid;

2-[[(2-thienyl)methylhydroxyphosphinyl]amino]-2-phenylethanoic acid;

- 2-[[(3-thienyl) methylhydroxyphosphinyl] amino] -2-phenylethanoic acid;
- 2-[[(4-thienyl) methylhydroxyphosphinyl] amino] -2-phenylethanoic acid;
- 5 2-[[(3-thienyl) ethylhydroxyphosphinyl] amino] -2-phenylethanoic acid;
- 2-[[(3-thienyl) propylhydroxyphosphinyl] amino] -2-phenylethanoic acid;
- 10 2-[[benzylhydroxyphosphinyl] amino] -2-(2-pyridyl) methylethanoic acid;
- 2-[[benzylhydroxyphosphinyl] amino] -2-(3-pyridyl) methylethanoic acid;
- 2-[[benzylhydroxyphosphinyl] amino] -2-(4-pyridyl) methylethanoic acid;
- 15 2-[[benzylhydroxyphosphinyl] amino] -2-(3-pyridyl) ethylethanoic acid;
- 2-[[benzylhydroxyphosphinyl] amino] -2-(3-pyridyl) propylethanoic acid;
- 20 2-[[benzylhydroxyphosphinyl] amino] -2-(tetrahydrofuran-2-yl) methyl ethanoic acid;
- 2-[[benzylhydroxyphosphinyl] amino] -2-(tetrahydrofuran-2-yl) ethyl ethanoic acid;
- 2-[[benzylhydroxyphosphinyl] amino] -2-(tetrahydrofuran-2-yl) propyl ethanoic acid;
- 25 2-[[benzylhydroxyphosphinyl] amino] -2-(2-indolyl) methylethanoic acid;
- 2-[[benzylhydroxyphosphinyl] amino] -2-(3-indolyl) methylethanoic acid;

- 2-[[benzylhydroxyphosphinyl]amino]-2-(4-indolyl)methylethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-(3-indolyl)ethylethanoic acid;
- 5 2-[[benzylhydroxyphosphinyl]amino]-2-(3-indolyl)propylethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-(2-thienyl)methylethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-(3-thienyl)methylethanoic acid;
- 10 2-[[benzylhydroxyphosphinyl]amino]-2-(4-thienyl)methylethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-(3-thienyl)ethylethanoic acid;
- 15 2-[[benzylhydroxyphosphinyl]amino]-2-(3-thienyl)propylethanoic acid; and a pharmaceutically acceptable salt, hydrate, or a mixture thereof.

19. The method of claim 17 wherein the NAALADase inhibitor is selected from the group consisting of:
- 20 2-[(benzylhydroxyphosphinyl)amino]pentanedioic acid;
- 2-[(phenylhydroxyphosphinyl)amino]pentanedioic acid;
- 2-[[[(hydroxy)phenylmethyl]hydroxyphosphinyl]amino]pentanedioic acid;
- 2-[(butylhydroxyphosphinyl)amino]pentanedioic acid;
- 25 2-[[[(3-methylbenzyl)hydroxyphosphinyl]amino]pentanedioic acid;
- 2-[(3-phenylpropylhydroxyphosphinyl)amino]pentanedioic acid;
- 2-[[[(4-fluorophenyl)hydroxyphosphinyl]amino]pentanedioic acid;
- 2-[(methylhydroxyphosphinyl)amino]pentanedioic acid;

2-[(phenylethylhydroxyphosphinyl)amino]pentanedioic acid;
2-[[[(4-methylbenzyl)hydroxyphosphinyl]amino]pentanedioic acid;
2-[[[(4-fluorobenzyl)hydroxyphosphinyl]amino]pentanedioic acid;
2-[[[(4-methoxybenzyl)hydroxyphosphinyl]amino]pentanedioic
5 acid;
2-[(phosphono)amino]pentanedioic acid;
2-[[[(3-trifluoromethylbenzyl)hydroxyphosphinyl]amino]
pentanedioic acid;
2-[[[(2-fluorobenzyl)hydroxyphosphinyl]amino]pentanedioic acid;
10 2-[[[(pentafluorobenzyl)hydroxyphosphinyl]amino]pentanedioic
acid; and a pharmaceutically acceptable salt, hydrate, or a
mixture thereof.

20. The method of claim 10 wherein R_1 or R_2 is
heterocyclic and X is CH_2 .

15 21. The method of claim 20 wherein the NAALADase
inhibitor is selected from the group consisting of:
2-[[[(2-pyridyl)hydroxyphosphinyl]methyl]pentanedioic acid;
2-[[[(3-pyridyl)hydroxyphosphinyl]methyl]pentanedioic acid;
2-[[[(4-pyridyl)hydroxyphosphinyl]methyl]pentanedioic acid;
20 2-[[[(tetrahydrofuranyl)hydroxyphosphinyl]methyl]pentanedioic
acid;
2-[[[(2-indolyl)hydroxyphosphinyl]methyl]pentanedioic acid;
2-[[[(3-indolyl)hydroxyphosphinyl]methyl]pentanedioic acid;
2-[[[(4-indolyl)hydroxyphosphinyl]methyl]pentanedioic acid;
25 2-[[[(2-thienyl)hydroxyphosphinyl]methyl]pentanedioic acid;
2-[[[(3-thienyl)hydroxyphosphinyl]methyl]pentanedioic acid;
2-[[[(4-thienyl)hydroxyphosphinyl]methyl]pentanedioic acid;
3-[(2-pyridyl)hydroxyphosphinyl]-2-phenylpropanoic acid;

3-[(3-pyridyl)hydroxyphosphinyl]-2-phenylpropanoic acid;
3-[(4-pyridyl)hydroxyphosphinyl]-2-phenylpropanoic acid;
3-[(tetrahydrofuranyl)hydroxyphosphinyl]-2-phenylpropanoic
acid;
5 3-[(2-indolyl)hydroxyphosphinyl]-2-phenylpropanoic acid;
3-[(3-indolyl)hydroxyphosphinyl]-2-phenylpropanoic acid;
3-[(4-indolyl)hydroxyphosphinyl]-2-phenylpropanoic acid;
3-[(2-thienyl)hydroxyphosphinyl]-2-phenylpropanoic acid;
3-[(3-thienyl)hydroxyphosphinyl]-2-phenylpropanoic acid;
10 3-[(4-thienyl)hydroxyphosphinyl]-2-phenylpropanoic acid;
3-(benzylhydroxyphosphinyl)-2-(2-pyridyl)propanoic acid;
3-(benzylhydroxyphosphinyl)-2-(3-pyridyl)propanoic acid;
3-(benzylhydroxyphosphinyl)-2-(4-pyridyl)propanoic acid;
3-(benzylhydroxyphosphinyl)-2-(tetrahydrofuranyl)propanoic
15 acid;
3-(benzylhydroxyphosphinyl)-2-(2-indolyl)propanoic acid;
3-(benzylhydroxyphosphinyl)-2-(3-indolyl)propanoic acid;
3-(benzylhydroxyphosphinyl)-2-(4-indolyl)propanoic acid;
3-(benzylhydroxyphosphinyl)-2-(2-thienyl)propanoic acid;
20 3-(benzylhydroxyphosphinyl)-2-(3-thienyl)propanoic acid;
3-(benzylhydroxyphosphinyl)-2-(4-thienyl)propanoic acid; and
a pharmaceutically acceptable salt, hydrate, or a mixture
thereof.

22. The method of claim 10 wherein R_1 or R_2 is
25 heterocyclic and X is oxygen.

23. The method of claim 22 wherein the NAALADase
inhibitor is selected from the group consisting of:
2-[[[(2-pyridyl)hydroxyphosphinyl]oxy]pentanedioic acid;

- 2-[[[(3-pyridyl)hydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(4-pyridyl)hydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(tetrahydrofuranyl)hydroxyphosphinyl]oxy]pentanedioic acid;
5 2-[[[(2-indolyl)hydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(3-indolyl)hydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(4-indolyl)hydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(2-thienyl)hydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(3-thienyl)hydroxyphosphinyl]oxy]pentanedioic acid;
10 2-[[[(4-thienyl)hydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(2-pyridyl)hydroxyphosphinyl]oxy]-2-phenylethanoic acid;
2-[[[(3-pyridyl)hydroxyphosphinyl]oxy]-2-phenylethanoic acid;
2-[[[(4-pyridyl)hydroxyphosphinyl]oxy]-2-phenylethanoic acid;
2-[[[(tetrahydrofuranyl)hydroxyphosphinyl]oxy]-2-phenylethanoic acid;
15 acid;
2-[[[(2-indolyl)hydroxyphosphinyl]oxy]-2-phenylethanoic acid;
2-[[[(3-indolyl)hydroxyphosphinyl]oxy]-2-phenylethanoic acid;
2-[[[(4-indolyl)hydroxyphosphinyl]oxy]-2-phenylethanoic acid;
2-[[[(2-thienyl)hydroxyphosphinyl]oxy]-2-phenylethanoic acid;
20 2-[[[(3-thienyl)hydroxyphosphinyl]oxy]-2-phenylethanoic acid;
2-[[[(4-thienyl)hydroxyphosphinyl]oxy]-2-phenylethanoic acid;
2-[[[benzylhydroxyphosphinyl]oxy]-2-(2-pyridyl)ethanoic acid;
2-[[[benzylhydroxyphosphinyl]oxy]-2-(3-pyridyl)ethanoic acid;
2-[[[benzylhydroxyphosphinyl]oxy]-2-(4-pyridyl)ethanoic acid;
25 2-[[[benzylhydroxyphosphinyl]oxy]-2-(tetrahydrofuranyl)ethanoic acid;
2-[[[benzylhydroxyphosphinyl]oxy]-2-(2-indolyl)ethanoic acid;
2-[[[benzylhydroxyphosphinyl]oxy]-2-(3-indolyl)ethanoic acid;

2-[[benzylhydroxyphosphinyl]oxy]-2-(4-indolyl)ethanoic acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-(2-thienyl)ethanoic acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-(3-thienyl)ethanoic acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-(4-thienyl)ethanoic acid;
5 and a pharmaceutically acceptable salt, hydrate, or a mixture thereof.

24. The method of claim 10 wherein R_1 or R_2 is heterocyclic and X is NR_1 .

25. The method of claim 24 wherein the NAALADase
10 inhibitor is a compound selected from the group consisting of:
2-[[[(2-pyridyl)hydroxyphosphinyl]amino]pentanedioic acid;
2-[[[(3-pyridyl)hydroxyphosphinyl]amino]pentanedioic acid;
2-[[[(4-pyridyl)hydroxyphosphinyl]amino]pentanedioic acid;
2-[[[(tetrahydrofuranyl)hydroxyphosphinyl]amino]pentanedioic
15 acid;
2-[[[(2-indolyl)hydroxyphosphinyl]amino]pentanedioic acid;
2-[[[(3-indolyl)hydroxyphosphinyl]amino]pentanedioic acid;
2-[[[(4-indolyl)hydroxyphosphinyl]amino]pentanedioic acid;
2-[[[(2-thienyl)hydroxyphosphinyl]amino]pentanedioic acid;
20 2-[[[(3-thienyl)hydroxyphosphinyl]amino]pentanedioic acid;
2-[[[(4-thienyl)hydroxyphosphinyl]amino]pentanedioic acid;
2-[[[(2-pyridyl)hydroxyphosphinyl]amino]-2-phenylethanoic acid;
2-[[[(3-pyridyl)hydroxyphosphinyl]amino]-2-phenylethanoic acid;
2-[[[(4-pyridyl)hydroxyphosphinyl]amino]-2-phenylethanoic acid;
25 2-[[[(tetrahydrofuranyl)hydroxyphosphinyl]amino]-2-phenyl
ethanoic acid;
2-[[[(2-indolyl)hydroxyphosphinyl]amino]-2-phenylethanoic acid;
2-[[[(3-indolyl)hydroxyphosphinyl]amino]-2-phenylethanoic acid;

-157-

2-[[(4-indolyl)hydroxyphosphinyl]amino]-2-phenylethanoic acid;
2-[[(2-thienyl)hydroxyphosphinyl]amino]-2-phenylethanoic acid;
2-[[(3-thienyl)hydroxyphosphinyl]amino]-2-phenylethanoic acid;
2-[[(4-thienyl)hydroxyphosphinyl]amino]-2-phenylethanoic acid;
5 2-[[benzylhydroxyphosphinyl]amino]-2-(2-pyridyl)ethanoic acid;
2-[[benzylhydroxyphosphinyl]amino]-2-(3-pyridyl)ethanoic acid;
2-[[benzylhydroxyphosphinyl]amino]-2-(4-pyridyl)ethanoic acid;
2-[[benzylhydroxyphosphinyl]amino]-2-(tetrahydrofuranyl)
ethanoic acid;
10 2-[[benzylhydroxyphosphinyl]amino]-2-(2-indolyl)ethanoic acid;
2-[[benzylhydroxyphosphinyl]amino]-2-(3-indolyl)ethanoic acid;
2-[[benzylhydroxyphosphinyl]amino]-2-(4-indolyl)ethanoic acid;
2-[[benzylhydroxyphosphinyl]amino]-2-(2-thienyl)ethanoic acid;
2-[[benzylhydroxyphosphinyl]amino]-2-(3-thienyl)ethanoic acid;
15 2-[[benzylhydroxyphosphinyl]amino]-2-(4-thienyl)ethanoic acid;
and a pharmaceutically acceptable salt, hydrate, or a mixture
thereof.

26. A method of inhibiting tumor cell growth in an
animal which comprises: administering to an animal suffering
20 from a tumor cell growth an effective amount of a NAALADase
inhibitor.

27. The method of claim 26 where the tumor is prostate
adenocarcinoma.

28. The method of claim 26 wherein the NAALADase
25 inhibitor is selected from the group consisting of: a
glutamate-derived hydroxyphosphinyl derivative compound; an
acidic peptide analog; a conformationally restricted glutamate
mimic; and mixtures thereof.

29. The method of claim 28 wherein the acidic peptide analog is selected from the group consisting of: Asp-Glu, Glu-Glu, Gly-Glu, gamma-Glu-Glu, and Glu-Glu-Glu.

30. The method of claim 28 wherein the conformationally restricted glutamate mimic is selected from the group consisting of: beta:NAAG and quisqualic acid.

31. The method of claim 28 wherein the NAALADase inhibitor is a glutamate-derived hydroxyphosphinyl derivative compound.

32. The method of claim 26 wherein the NAALADase inhibitor is administered in combination with an additional therapeutic agent selected from the group consisting of: therapeutic hormones, chemotherapeutic hormones, anti-angiogenesis agents, radiolabelled compounds, and mixtures thereof.


33. The method of claim 26 wherein the tumor is selected from the group consisting of: ACTH-producing tumor, acute lymphocytic leukemia, acute nonlymphocytic leukemia, cancer of the adrenal cortex, bladder cancer, brain cancer, breast cancer, cervical cancer, chronic lymphocytic leukemia, chronic myelocytic leukemia, colorectal cancer, cutaneous T-cell lymphoma, endometrial cancer, esophageal cancer, Ewing's sarcoma, gallbladder cancer, hairy cell leukemia, head & neck cancer, Hodgkin's lymphoma, Kaposi's sarcoma, kidney cancer, liver cancer, lung cancer, lung cancer (small and/or non-small cell), malignant peritoneal effusion, malignant pleural effusion, melanoma, mesothelioma, multiple myeloma, neuroblastoma, non-Hodgkin's lymphoma, osteosarcoma, ovary

cancer, ovary (germ cell) cancer, pancreatic cancer, penis cancer, prostate adenocarcinoma, retinoblastoma, skin cancer, soft tissue sarcoma, squamous cell carcinomas, stomach cancer, testicular cancer, thyroid cancer, trophoblastic neoplasms, cancer of the uterus, vaginal cancer, cancer of the vulva, and Wilm's tumor.

34. The method of claim 26 wherein the tumor is prostatic adenocarcinoma.

35. The method of claim 26 wherein the tumor is selected from the group consisting of: brain cancer, cancer of the adrenal cortex, kidney cancer, and testicular cancer.

36. The method of claim 26 wherein the NAALADase inhibitor comprises a compound having the following formula:



15

I

wherein

- R_1 is hydrogen, C_1 - C_9 straight or branched chain alkyl, C_2 - C_9 straight or branched chain alkenyl group, C_3 - C_8 cycloalkyl, C_5 - C_7 cycloalkenyl, or Ar_1 ;
- X is CH_2 , O , or NR_1 , where R_1 is defined above; and
- R_2 is C_1 - C_9 straight or branched chain alkyl, C_2 - C_9 straight or branched chain alkenyl group, C_3 - C_8 cycloalkyl, C_5 - C_7 cycloalkenyl, or Ar_1 , wherein said alkyl, alkenyl, cycloalkyl, cycloalkenyl or aryl group may be optionally substituted with carboxylic

25

acid,

wherein said alkyl, alkenyl, cycloalkyl, cycloalkenyl or aryl groups may be optionally substituted with C₁-C₈ cycloalkyl, C₁ or C₂ cycloalkyl, C₃-C₇ cycloalkenyl, C₁-C₄ alkyl, C₁-C₄ alkenyl, halo, hydroxy, carboxy, nitro, trifluoromethyl, C₁-C₅ straight or branched chain alkyl or alkenyl, C₁-C₄ alkoxy, C₁-C₄ alkenyloxy, phenoxy, benzyloxy, amino, or Ar₁, and where Ar₁ is selected from the group consisting of 1-naphthyl, 2-naphthyl, 2-indolyl, 3-indolyl, 4-indolyl, 2-furyl, 3-furyl, tetrahydrofuranyl, 2-thienyl, 3-thienyl, 4-thienyl, 2-, 3-, or 4-pyridyl, or phenyl, having one to five substituents which are independently selected from the group consisting of hydrogen, halo, hydroxy, carboxy, nitro, trifluoromethyl, C₁-C₅ straight or branched alkyl or alkenyl, C₁-C₄ alkoxy or C₁-C₄ alkenyloxy, phenoxy, benzyloxy, and amino; or pharmaceutically acceptable salts, hydrates, or mixtures thereof.

37. The method of claim 36 wherein R₁ and R₂ are straight or branched aliphatic groups or carbocyclic groups and X is CH₂.

38. The method of claim 37 wherein the NAALADase inhibitor is selected from the group consisting of:

2-[[methylhydroxyphosphinyl]methyl]pentanedioic acid;
2-[[ethylhydroxyphosphinyl]methyl]pentanedioic acid;
2-[[propylhydroxyphosphinyl]methyl]pentanedioic acid;
2-[[butylhydroxyphosphinyl]methyl]pentanedioic acid;
2-[[cyclohexylhydroxyphosphinyl]methyl]pentanedioic acid;
2-[[[(cyclohexyl)methylhydroxyphosphinyl]methyl]pentanedioic acid;

- 2-[[phenylhydroxyphosphinyl]methyl]pentanedioic acid;
2-[[benzylhydroxyphosphinyl]methyl]pentanedioic acid;
2-[[phenylethylhydroxyphosphinyl]methyl]pentanedioic acid;
2-[[phenylpropylhydroxyphosphinyl]methyl]pentanedioic acid;
5 2-[[phenylbutylhydroxyphosphinyl]methyl]pentanedioic acid;
2-[[[(4-methylbenzyl)hydroxyphosphinyl]methyl]pentanedioic
acid;
2-[[[(4-fluorobenzyl)hydroxyphosphinyl]methyl]pentanedioic
acid;
10 2-[[[(2-fluorobenzyl)hydroxyphosphinyl]methyl]pentanedioic
acid;
2-[[[(pentafluorobenzyl)hydroxyphosphinyl]methyl]pentanedioic
acid;
2-[[[(methoxybenzyl)hydroxyphosphinyl]methyl]pentanedioic acid;
15 2-[[[(2, 3, 4-trimethoxyphenyl)hydroxyphosphinyl]methyl]
pentanedioic acid;
2-[[[(1-naphthyl)hydroxyphosphinyl]methyl]pentanedioic acid;
2-[[[(2-naphthyl)hydroxyphosphinyl]methyl]pentanedioic acid;
2-[[[(1-naphthyl)methylhydroxyphosphinyl]methyl]pentanedioic
20 acid;
2-[[[(2-naphthyl)methylhydroxyphosphinyl]methyl]pentanedioic
acid;
2-[[[(1-naphthyl)ethylhydroxyphosphinyl]methyl]pentanedioic
acid;
25 2-[[[(2-naphthyl)ethylhydroxyphosphinyl]methyl]pentanedioic
acid;
2-[[[(1-naphthyl)propylhydroxyphosphinyl]methyl]pentanedioic
acid;

- 2-[[[(2-naphthyl)propylhydroxyphosphinyl]methyl]pentanedioic acid;
- 2-[[[(1-naphthyl)butylhydroxyphosphinyl]methyl]pentanedioic acid;
- 5 2-[[[(2-naphthyl)butylhydroxyphosphinyl]methyl]pentanedioic acid;
- 2-[[[(phenylprop-2-enyl)hydroxyphosphinyl]methyl]pentanedioic acid;
- 2-[[[(2-fluorobenzyl)hydroxyphosphinyl]methyl]pentanedioic acid;
- 10 2-[[[(4-fluorophenyl)hydroxyphosphinyl]methyl]pentanedioic acid;
- 2-[[[(hydroxy)phenylmethyl]hydroxyphosphinyl]methyl]pentanedioic acid;
- 15 2-[[[(3-methylbenzyl)hydroxyphosphinyl]methyl]pentanedioic acid;
- 2-(phosphonomethyl)pentanedioic acid;
- 2-[[[(3-trifluoromethylbenzyl)hydroxyphosphinyl]methyl]pentanedioic acid;
- 20 3-(methylhydroxyphosphinyl)-2-phenylpropanoic acid;
- 3-(ethylhydroxyphosphinyl)-2-phenylpropanoic acid;
- 3-(propylhydroxyphosphinyl)-2-phenylpropanoic acid;
- 3-(butylhydroxyphosphinyl)-2-phenylpropanoic acid;
- 3-(cyclohexylhydroxyphosphinyl)-2-phenylpropanoic acid;
- 25 3-((cyclohexyl)methylhydroxyphosphinyl)-2-phenylpropanoic acid;
- 3-(phenylhydroxyphosphinyl)-2-phenylpropanoic acid;
- 3-(benzylhydroxyphosphinyl)-2-phenylpropanoic acid;

-163-

- 3-(phenylethylhydroxyphosphinyl)-2-phenylpropanoic acid;
3-(phenylpropylhydroxyphosphinyl)-2-phenylpropanoic acid;
3-(phenylbutylhydroxyphosphinyl)-2-phenylpropanoic acid;
3-((2, 3, 4-trimethoxyphenyl)-3-hydroxyphosphinyl)-2-phenylpropanoic acid;
5 3-((1-naphthyl)hydroxyphosphinyl)-2-phenylpropanoic acid;
3-((2-naphthyl)hydroxyphosphinyl)-2-phenylpropanoic acid;
3-((1-naphthyl)methylhydroxyphosphinyl)-2-phenylpropanoic acid;
10 3-((2-naphthyl)methylhydroxyphosphinyl)-2-phenylpropanoic acid;
3-((1-naphthyl)ethylhydroxyphosphinyl)-2-phenylpropanoic acid;
3-((2-naphthyl)ethylhydroxyphosphinyl)-2-phenylpropanoic acid;
3-((1-naphthyl)propylhydroxyphosphinyl)-2-phenylpropanoic acid;
15 3-((2-naphthyl)propylhydroxyphosphinyl)-2-phenylpropanoic acid;
3-((1-naphthyl)butylhydroxyphosphinyl)-2-phenylpropanoic acid;
3-((2-naphthyl)butylhydroxyphosphinyl)-2-phenylpropanoic acid;
20 3-(phenylprop-2-enylhydroxyphosphinyl)-2-phenylpropanoic acid;
2-[(benzylhydroxyphosphinyl)methyl]pentanedioic acid;
2-[(methylhydroxyphosphinyl)methyl]hexanedioic acid;
2-[(benzylhydroxyphosphinyl)methyl]hexanedioic acid;
2-[(methylhydroxyphosphinyl)methyl]heptanedioic acid;
25 2-[(benzylhydroxyphosphinyl)methyl]heptanedioic acid;
2-[(methylhydroxyphosphinyl)methyl]octanedioic acid;
2-[(benzylhydroxyphosphinyl)methyl]octanedioic acid;
2-[(methylhydroxyphosphinyl)methyl]nonanedioic acid;

- 2-[(benzylhydroxyphosphinyl)methyl]nonanedioic acid;
2-[(methylhydroxyphosphinyl)methyl]decanedioic acid;
2-[(benzylhydroxyphosphinyl)methyl]decanedioic acid;
3-(benzylhydroxyphosphinyl)-2-methylpropanoic acid;
5 3-(benzylhydroxyphosphinyl)-2-ethylpropanoic acid;
3-(benzylhydroxyphosphinyl)-2-propylpropanoic acid;
3-(benzylhydroxyphosphinyl)-2-butylpropanoic acid;
3-(benzylhydroxyphosphinyl)-2-cyclohexylpropanoic acid;
3-(benzylhydroxyphosphinyl)-2-(cyclohexyl)methylpropanoic
10 acid;
3-(benzylhydroxyphosphinyl)-2-phenylpropanoic acid;
3-(benzylhydroxyphosphinyl)-2-benzylpropanoic acid;
3-(benzylhydroxyphosphinyl)-2-phenylethylpropanoic acid;
3-(benzylhydroxyphosphinyl)-2-phenylpropylpropanoic acid;
15 3-(benzylhydroxyphosphinyl)-2-phenylbutylpropanoic acid;
3-(benzylhydroxyphosphinyl)-2-(2, 3, 4-trimethoxyphenyl)
propanoic acid;
3-(benzylhydroxyphosphinyl)-2-(1-naphthyl)propanoic acid;
3-(benzylhydroxyphosphinyl)-2-(2-naphthyl)propanoic acid;
20 3-(benzylhydroxyphosphinyl)-2-(1-naphthyl)methylpropanoic
acid;
3-(benzylhydroxyphosphinyl)-2-(2-naphthyl)methylpropanoic
acid;
3-(benzylhydroxyphosphinyl)-2-(1-naphthyl)ethylpropanoic acid;
25 3-(benzylhydroxyphosphinyl)-2-(2-naphthyl)ethylpropanoic acid;
3-(benzylhydroxyphosphinyl)-2-(1-naphthyl)propylpropanoic
acid;
3-(benzylhydroxyphosphinyl)-2-(2-naphthyl)propylpropanoic

- acid;
- 3-(benzylhydroxyphosphinyl)-2-(1-naphthyl)butylpropanoic acid;
- 3-(benzylhydroxyphosphinyl)-2-(2-naphthyl)butylpropanoic acid;
- 3-(benzylhydroxyphosphinyl)-2-phenylprop-2-enylpropanoic acid;
- 5 2-[[(2-pyridyl)methylhydroxyphosphinyl]methyl]pentanedioic acid;
- 2-[[(3-pyridyl)methylhydroxyphosphinyl]methyl]pentanedioic acid;
- 2-[[(4-pyridyl)methylhydroxyphosphinyl]methyl]pentanedioic acid;
- 10 2-[[(3-pyridyl)ethylhydroxyphosphinyl]methyl]pentanedioic acid;
- 2-[[(3-pyridyl)propylhydroxyphosphinyl]methyl]pentanedioic acid;
- 15 2-[[(tetrahydrofuranyl)methylhydroxyphosphinyl]methyl]pentanedioic acid;
- 2-[[(tetrahydrofuranyl)ethylhydroxyphosphinyl]methyl]pentanedioic acid;
- 2-[[(tetrahydrofuranyl)propylhydroxyphosphinyl]methyl]pentanedioic acid;
- 20 2-[[(2-indolyl)methylhydroxyphosphinyl]methyl]pentanedioic acid;
- 2-[[(3-indolyl)methylhydroxyphosphinyl]methyl]pentanedioic acid;
- 25 2-[[(4-indolyl)methylhydroxyphosphinyl]methyl]pentanedioic acid;
- 2-[[(3-indolyl)ethylhydroxyphosphinyl]methyl]pentanedioic acid;

-166-

- 2-[[[(3-indolyl)propylhydroxyphosphinyl)methyl]pentanedioic acid;
- 2-[[[(2-thienyl)methylhydroxyphosphinyl)methyl]pentanedioic acid;
- 5 2-[[[(3-thienyl)methylhydroxyphosphinyl)methyl]pentanedioic acid;
- 2-[[[(4-thienyl)methylhydroxyphosphinyl)methyl]pentanedioic acid;
- 2-[[[(3-thienyl)ethylhydroxyphosphinyl)methyl]pentanedioic acid;
- 10 2-[[[(3-thienyl)propylhydroxyphosphinyl)methyl]pentanedioic acid;
- 3-[[[(2-pyridyl)methylhydroxyphosphinyl]-2-phenylpropanoic acid;
- 15 3-[[[(3-pyridyl)methylhydroxyphosphinyl]-2-phenylpropanoic acid;
- 3-[[[(4-pyridyl)methylhydroxyphosphinyl]-2-phenylpropanoic acid;
- 3-[[[(3-pyridyl)ethylhydroxyphosphinyl]-2-phenylpropanoic acid;
- 20 3-[[[(3-pyridyl)propylhydroxyphosphinyl]-2-phenylpropanoic acid;
- 3-[[[(tetrahydrofuranyl)methylhydroxyphosphinyl]-2-phenylpropanoic acid;
- 3-[[[(tetrahydrofuranyl)ethylhydroxyphosphinyl]-2-phenylpropanoic acid;
- 25 3-[[[(tetrahydrofuranyl)propylhydroxyphosphinyl]-2-phenylpropanoic acid;
- 3-[[[(2-indolyl)methylhydroxyphosphinyl]-2-phenylpropanoic

- acid;
3-[[(3-indolyl) methylhydroxyphosphinyl] -2-phenylpropanoic
acid;
3-[[(4-indolyl) methylhydroxyphosphinyl] -2-phenylpropanoic
5 acid;
3-[[(3-indolyl) ethylhydroxyphosphinyl] -2-phenylpropanoic acid;
3-[[(3-indolyl) propylhydroxyphosphinyl] -2-phenylpropanoic
acid;
3-[[(2-thienyl) methylhydroxyphosphinyl] -2-phenylpropanoic
10 acid;
3-[[(3-thienyl) methylhydroxyphosphinyl] -2-phenylpropanoic
acid;
3-[[(4-thienyl) methylhydroxyphosphinyl] -2-phenylpropanoic
acid;
15 3-[[(3-thienyl) ethylhydroxyphosphinyl] -2-phenylpropanoic acid;
3-[[(3-thienyl) propylhydroxyphosphinyl] -2-phenylpropanoic
acid;
3- (benzylhydroxyphosphinyl) -2- (2-pyridyl) methylpropanoic acid;
3- (benzylhydroxyphosphinyl) -2- (3-pyridyl) methylpropanoic acid;
20 3- (benzylhydroxyphosphinyl) -2- (4-pyridyl) methylpropanoic acid;
3- (benzylhydroxyphosphinyl) -2- (3-pyridyl) ethylpropanoic acid;
3- (benzylhydroxyphosphinyl) -2- (3-pyridyl) propylpropanoic acid;
3- (benzylhydroxyphosphinyl) -2- (tetrahydrofuranyl) methyl
propanoic acid;
25 3- (benzylhydroxyphosphinyl) -2- (tetrahydrofuranyl) ethyl
propanoic acid;
3- (benzylhydroxyphosphinyl) -2- (tetrahydrofuranyl) propyl
propanoic acid;

3-(benzylhydroxyphosphinyl)-2-(2-indolyl)methylpropanoic acid;
3-(benzylhydroxyphosphinyl)-2-(3-indolyl)methylpropanoic acid;
3-(benzylhydroxyphosphinyl)-2-(4-indolyl)methylpropanoic acid;
3-(benzylhydroxyphosphinyl)-2-(3-indolyl)ethylpropanoic acid;
5 3-(benzylhydroxyphosphinyl)-2-(3-indolyl)propylpropanoic acid;
3-(benzylhydroxyphosphinyl)-2-(2-thienyl)methylpropanoic acid;
3-(benzylhydroxyphosphinyl)-2-(3-thienyl)methylpropanoic acid;
3-(benzylhydroxyphosphinyl)-2-(4-thienyl)methylpropanoic acid;
3-(benzylhydroxyphosphinyl)-2-(3-thienyl)ethylpropanoic acid;
10 3-(benzylhydroxyphosphinyl)-2-(3-thienyl)propylpropanoic acid;
and a pharmaceutically acceptable salt, hydrate, or a mixture thereof.

39. The method of claim 37 wherein the NAALADase inhibitor is selected from the group consisting of:

15 2-[(benzylhydroxyphosphinyl)methyl]pentanedioic acid;
2-[(phenylhydroxyphosphinyl)methyl]pentanedioic acid;
2-[[[(hydroxy)phenylmethyl]hydroxyphosphinyl]methyl]
pentanedioic acid;
2-[(butylhydroxyphosphinyl)methyl]pentanedioic acid;
20 2-[[[(3-methylbenzyl)hydroxyphosphinyl]methyl]pentanedioic
acid;
2-[(3-phenylpropylhydroxyphosphinyl)methyl]pentanedioic acid;
2-[[[(4-fluorophenyl)hydroxyphosphinyl]methyl]pentanedioic
acid;
25 2-[(methylhydroxyphosphinyl)methyl]pentanedioic acid;
2-[(phenylethylhydroxyphosphinyl)methyl]pentanedioic acid;
2-[[[(4-methylbenzyl)hydroxyphosphinyl]methyl]pentanedioic
acid;

2-[[[(4-fluorobenzyl)hydroxyphosphinyl)methyl]pentanedioic acid;

2-[[[(4-methoxybenzyl)hydroxyphosphinyl)methyl]pentanedioic acid;

5 2-(phosphonomethyl)pentanedioic acid;

2-[[[(3-trifluoromethylbenzyl)hydroxyphosphinyl)methyl]pentanedioic acid;

2-[[[(2-fluorobenzyl)hydroxyphosphinyl)methyl]pentanedioic acid;

10 2-[[[(pentafluorobenzyl)hydroxyphosphinyl)methyl]pentanedioic acid; and a pharmaceutically acceptable salt, hydrate, or a mixture thereof.

40. The method of claim 36 wherein R_1 and R_2 are straight or branched or carbocyclic groups and X is oxygen.

15 41. The method of claim 40 wherein the NAALADase inhibitor is selected from the group consisting of:

2-[[methylhydroxyphosphinyl]oxy]pentanedioic acid;

2-[[ethylhydroxyphosphinyl]oxy]pentanedioic acid;

2-[[propylhydroxyphosphinyl]oxy]pentanedioic acid;

20 2-[[butylhydroxyphosphinyl]oxy]pentanedioic acid;

2-[[cyclohexylhydroxyphosphinyl]oxy]pentanedioic acid;

2-[[[(cyclohexyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;

2-[[phenylhydroxyphosphinyl]oxy]pentanedioic acid;

2-[[benzylhydroxyphosphinyl]oxy]pentanedioic acid;

25 2-[[phenylethylhydroxyphosphinyl]oxy]pentanedioic acid;

2-[[phenylpropylhydroxyphosphinyl]oxy]pentanedioic acid;

2-[[phenylbutylhydroxyphosphinyl]oxy]pentanedioic acid;

2-[[[(4-methylbenzyl)hydroxyphosphinyl]oxy]pentanedioic acid;

- 2-[[[(4-fluorobenzyl)hydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(2-fluorobenzyl)hydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(pentafluorobenzyl)hydroxyphosphinyl]oxy]pentanedioic acid;
5 2-[[[(methoxybenzyl)hydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(2, 3, 4-trimethoxyphenyl)hydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(1-naphthyl)hydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(2-naphthyl)hydroxyphosphinyl]oxy]pentanedioic acid;
10 2-[[[(1-naphthyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(2-naphthyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(1-naphthyl)ethylhydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(2-naphthyl)ethylhydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(1-naphthyl)propylhydroxyphosphinyl]oxy]pentanedioic acid;
15 2-[[[(2-naphthyl)propylhydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(1-naphthyl)butylhydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(2-naphthyl)butylhydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(phenylprop-2-enyl)hydroxyphosphinyl]oxy]pentanedioic acid;
20 2-[[benzylhydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(2-fluorobenzyl)hydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(4-fluorophenyl)hydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(hydroxy)phenylmethyl]hydroxyphosphinyl]methyl]pentanedioic acid;
25 2-[[[(3-methylbenzyl)hydroxyphosphinyl]methyl]pentanedioic acid;
2-[(phosphono)oxy]pentanedioic acid;
2-[[[(3-trifluoromethylbenzyl)hydroxyphosphinyl]oxy]

- pentanedioic acid;
- 2-[[methylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[ethylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[propylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 5 2-[[butylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[cyclohexylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[[(cyclohexyl)methylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[phenylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 10 2-[[benzylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[phenylethylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[phenylpropylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[phenylbutylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[[(2, 3, 4-trimethoxyphenyl)-3-hydroxyphosphinyl]oxy]-2-
- 15 phenylethanoic acid;
- 2-[[[(1-naphthyl)hydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[[(2-naphthyl)hydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[[(1-naphthyl)methylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 20 2-[[[(2-naphthyl)methylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[[(1-naphthyl)ethylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[[(2-naphthyl)ethylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 25 2-[[[(1-naphthyl)propylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[[(2-naphthyl)propylhydroxyphosphinyl]oxy]-2-phenylethanoic

- acid;
- 2-[[(1-naphthyl)butylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[(2-naphthyl)butylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 5 2-[[phenylprop-2-enylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[(methylhydroxyphosphinyl)oxy]hexanedioic acid;
- 2-[(benzylhydroxyphosphinyl)oxy]hexanedioic acid;
- 10 2-[(methylhydroxyphosphinyl)oxy]heptanedioic acid;
- 2-[(benzylhydroxyphosphinyl)oxy]heptanedioic acid;
- 2-[(methylhydroxyphosphinyl)oxy]octanedioic acid;
- 2-[(benzylhydroxyphosphinyl)oxy]octanedioic acid;
- 2-[(methylhydroxyphosphinyl)oxy]nonanedioic acid;
- 15 2-[(benzylhydroxyphosphinyl)oxy]nonanedioic acid;
- 2-[(methylhydroxyphosphinyl)oxy]decanedioic acid;
- 2-[(benzylhydroxyphosphinyl)oxy]decanedioic acid;
- 2-[[benzylhydroxyphosphinyl]oxy]-2-methylethanoic acid;
- 2-[[benzylhydroxyphosphinyl]oxy]-2-ethylethanoic acid;
- 20 2-[[benzylhydroxyphosphinyl]oxy]-2-propylethanoic acid;
- 2-[[benzylhydroxyphosphinyl]oxy]-2-butylethanoic acid;
- 2-[[benzylhydroxyphosphinyl]oxy]-2-cyclohexylethanoic acid;
- 2-[[benzylhydroxyphosphinyl]oxy]-2-(cyclohexyl)methylethanoic acid;
- 25 2-[[benzylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[benzylhydroxyphosphinyl]oxy]-2-benzylethanoic acid;
- 2-[[benzylhydroxyphosphinyl]oxy]-2-phenylethylethanoic acid;
- 2-[[benzylhydroxyphosphinyl]oxy]-2-phenylpropylethanoic acid;

- 2-[[benzylhydroxyphosphinyl]oxy]-2-phenylbutylethanoic acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-(2, 3, 4-trimethoxyphenyl)
ethanoic acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-(1-naphthyl)ethanoic acid;
5 2-[[benzylhydroxyphosphinyl]oxy]-2-(2-naphthyl)ethanoic acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-(1-naphthyl)methylethanoic
acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-(2-naphthyl)methylethanoic
acid;
10 2-[[benzylhydroxyphosphinyl]oxy]-2-(1-naphthyl)ethylethanoic
acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-(2-naphthyl)ethylethanoic
acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-(1-naphthyl)propylethanoic
15 acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-(2-naphthyl)propylethanoic
acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-(1-naphthyl)butylethanoic
acid;
20 2-[[benzylhydroxyphosphinyl]oxy]-2-(2-naphthyl)butylethanoic
acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-phenylprop-2-enylethanoic
acid;
2-[[[(2-pyridyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;
25 2-[[[(3-pyridyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(4-pyridyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(3-pyridyl)ethylhydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(3-pyridyl)propylhydroxyphosphinyl]oxy]pentanedioic acid;

- 2-[[[(tetrahydrofuranyl)methylhydroxyphosphinyl]oxy]
pentanedioic acid;
- 2-[[[(tetrahydrofuranyl)ethylhydroxyphosphinyl]oxy]pentanedioic
acid;
- 5 2-[[[(tetrahydrofuranyl)propylhydroxyphosphinyl]oxy]
pentanedioic acid;
- 2-[[[(2-indolyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;
- 2-[[[(3-indolyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;
- 2-[[[(4-indolyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;
- 10 2-[[[(3-indolyl)ethylhydroxyphosphinyl]oxy]pentanedioic acid;
- 2-[[[(3-indolyl)propylhydroxyphosphinyl]oxy]pentanedioic acid;
- 2-[[[(2-thienyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;
- 2-[[[(3-thienyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;
- 2-[[[(4-thienyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;
- 15 2-[[[(3-thienyl)ethylhydroxyphosphinyl]oxy]pentanedioic acid;
- 2-[[[(3-thienyl)propylhydroxyphosphinyl]oxy]pentanedioic acid;
- 2-[[[(2-pyridyl)methylhydroxyphosphinyl]oxy]-2-phenylethanoic
acid;
- 2-[[[(3-pyridyl)methylhydroxyphosphinyl]oxy]-2-phenylethanoic
20 acid;
- 2-[[[(4-pyridyl)methylhydroxyphosphinyl]oxy]-2-phenylethanoic
acid;
- 2-[[[(3-pyridyl)ethylhydroxyphosphinyl]oxy]-2-phenylethanoic
acid;
- 25 2-[[[(3-pyridyl)propylhydroxyphosphinyl]oxy]-2-phenylethanoic
acid;
- 2-[[[(tetrahydrofuranyl)methylhydroxyphosphinyl]oxy]-2-phenyl
ethanoic acid;

-175-

- 2-[[[(tetrahydrofuranyl)ethylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[[(tetrahydrofuranyl)propylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 5 2-[[[(2-indolyl)methylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[[(3-indolyl)methylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[[(4-indolyl)methylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 10 2-[[[(3-indolyl)ethylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[[(3-indolyl)propylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 15 2-[[[(2-thienyl)methylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[[(3-thienyl)methylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[[(4-thienyl)methylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 20 2-[[[(3-thienyl)ethylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[[(3-thienyl)propylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 25 2-[[benzylhydroxyphosphinyl]oxy]-2-(2-pyridyl)methylethanoic acid;
- 2-[[benzylhydroxyphosphinyl]oxy]-2-(3-pyridyl)methylethanoic acid;

2-[[benzylhydroxyphosphinyl]oxy]-2-(4-pyridyl)methylethanoic acid;

2-[[benzylhydroxyphosphinyl]oxy]-2-(3-pyridyl)ethylethanoic acid;

5 2-[[benzylhydroxyphosphinyl]oxy]-2-(3-pyridyl)propylethanoic acid;

2-[[benzylhydroxyphosphinyl]oxy]-2-(tetrahydrofuran-2-yl)methylethanoic acid;

10 2-[[benzylhydroxyphosphinyl]oxy]-2-(tetrahydrofuran-2-yl)ethylethanoic acid;

2-[[benzylhydroxyphosphinyl]oxy]-2-(tetrahydrofuran-2-yl)propylethanoic acid;

2-[[benzylhydroxyphosphinyl]oxy]-2-(2-indolyl)methylethanoic acid;

15 2-[[benzylhydroxyphosphinyl]oxy]-2-(3-indolyl)methylethanoic acid;

2-[[benzylhydroxyphosphinyl]oxy]-2-(4-indolyl)methylethanoic acid;

20 2-[[benzylhydroxyphosphinyl]oxy]-2-(3-indolyl)ethylethanoic acid;

2-[[benzylhydroxyphosphinyl]oxy]-2-(3-indolyl)propylethanoic acid;

2-[[benzylhydroxyphosphinyl]oxy]-2-(2-thienyl)methylethanoic acid;

25 2-[[benzylhydroxyphosphinyl]oxy]-2-(3-thienyl)methylethanoic acid;

2-[[benzylhydroxyphosphinyl]oxy]-2-(4-thienyl)methylethanoic acid;

2-[[benzylhydroxyphosphinyl]oxy]-2-(3-thienyl)ethylethanoic acid;

2-[[benzylhydroxyphosphinyl]oxy]-2-(3-thienyl)propylethanoic acid; and a pharmaceutically acceptable salt, hydrate, or a mixture thereof.

5

42. The method of claim 40 wherein the NAALADase inhibitor is selected from the group consisting of:

2-[(benzylhydroxyphosphinyl)oxy]pentanedioic acid;

2-[(phenylhydroxyphosphinyl)oxy]pentanedioic acid;

10

2-[[[(hydroxy)phenylmethyl]hydroxyphosphinyl]oxy]pentanedioic acid;

2-[(butylhydroxyphosphinyl)oxy]pentanedioic acid;

2-[[[(3-methylbenzyl)hydroxyphosphinyl]oxy]pentanedioic acid;

2-[(3-phenylpropylhydroxyphosphinyl)oxy]pentanedioic acid;

15

2-[[[(4-fluorophenyl)hydroxyphosphinyl]oxy]pentanedioic acid;

2-[(methylhydroxyphosphinyl)oxy]pentanedioic acid;

2-[(phenylethylhydroxyphosphinyl)oxy]pentanedioic acid;

2-[[[(4-methylbenzyl)hydroxyphosphinyl]oxy]pentanedioic acid;

2-[[[(4-fluorobenzyl)hydroxyphosphinyl]oxy]pentanedioic acid;

20

2-[[[(4-methoxybenzyl)hydroxyphosphinyl]oxy]pentanedioic acid;

2-[(phosphono)oxy]pentanedioic acid;

2-[[[(3-trifluoromethylbenzyl)hydroxyphosphinyl]oxy]pentanedioic acid;

2-[[[(2-fluorobenzyl)hydroxyphosphinyl]oxy]pentanedioic acid;

25

2-[[[(pentafluorobenzyl)hydroxyphosphinyl]oxy]pentanedioic acid; and a pharmaceutically acceptable salt, hydrate, or a mixture thereof.

43. The method of claim 36 wherein R₁ and R₂ are straight

or branched or carbocyclic and X is NR_1 .

44. The method of claim 43 wherein the NAALADase inhibitor is selected from the group consisting of:

- 2-[[methylhydroxyphosphinyl]amino]pentanedioic acid;
- 5 2-[[ethylhydroxyphosphinyl]amino]pentanedioic acid;
- 2-[[propylhydroxyphosphinyl]amino]pentanedioic acid;
- 2-[[butylhydroxyphosphinyl]amino]pentanedioic acid;
- 2-[[cyclohexylhydroxyphosphinyl]amino]pentanedioic acid;
- 2-[[[(cyclohexyl)methylhydroxyphosphinyl]amino]pentanedioic
- 10 acid;
- 2-[[phenylhydroxyphosphinyl]amino]pentanedioic acid;
- 2-[[benzylhydroxyphosphinyl]amino]pentanedioic acid;
- 2-[[phenylethylhydroxyphosphinyl]amino]pentanedioic acid;
- 2-[[phenylpropylhydroxyphosphinyl]amino]pentanedioic acid;
- 15 2-[[phenylbutylhydroxyphosphinyl]amino]pentanedioic acid;
- 2-[[[(4-methylbenzyl)hydroxyphosphinyl]amino]pentanedioic acid;
- 2-[[[(4-fluorobenzyl)hydroxyphosphinyl]amino]pentanedioic acid;
- 2-[[[(2-fluorobenzyl)hydroxyphosphinyl]amino]pentanedioic acid;
- 2-[[[(pentafluorobenzyl)hydroxyphosphinyl]amino]pentanedioic
- 20 acid;
- 2-[[[(methoxybenzyl)hydroxyphosphinyl]amino]pentanedioic acid;
- 2-[[[(2, 3, 4-trimethoxyphenyl)hydroxyphosphinyl]amino]
- pentanedioic acid;
- 2-[[[(1-naphthyl)hydroxyphosphinyl]amino]pentanedioic acid;
- 25 2-[[[(2-naphthyl)hydroxyphosphinyl]amino]pentanedioic acid;
- 2-[[[(1-naphthyl)methylhydroxyphosphinyl]amino]pentanedioic
- acid;
- 2-[[[(2-naphthyl)methylhydroxyphosphinyl]amino]pentanedioic

- acid;
- 2-[[[(1-naphthyl)ethylhydroxyphosphinyl]amino]pentanedioic acid;
- 2-[[[(2-naphthyl)ethylhydroxyphosphinyl]amino]pentanedioic acid;
- 5 acid;
- 2-[[[(1-naphthyl)propylhydroxyphosphinyl]amino]pentanedioic acid;
- 2-[[[(2-naphthyl)propylhydroxyphosphinyl]amino]pentanedioic acid;
- 10 2-[[[(1-naphthyl)butylhydroxyphosphinyl]amino]pentanedioic acid;
- 2-[[[(2-naphthyl)butylhydroxyphosphinyl]amino]pentanedioic acid;
- 2-[[[(phenylprop-2-enyl)hydroxyphosphinyl]amino]pentanedioic acid;
- 15 acid;
- 2-[[benzylhydroxyphosphinyl]amino]pentanedioic acid;
- 2-[[[(2-fluorobenzyl)hydroxyphosphinyl]amino]pentanedioic acid;
- 2-[[[(hydroxy)phenylmethyl]hydroxyphosphinyl]amino]pentanedioic acid;
- 20 2-[[[(3-methylbenzyl)hydroxyphosphinyl]amino]pentanedioic acid;
- 2-[[[(4-fluorophenyl)hydroxyphosphinyl]amino]pentanedioic acid;
- 2-[(phosphono)amino]pentanedioic acid;
- 2-[[[(3-trifluoromethylbenzyl)hydroxyphosphinyl]amino]pentanedioic acid;
- 25 2-[[methylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
- 2-[[ethylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
- 2-[[propylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
- 2-[[butylhydroxyphosphinyl]amino]-2-phenylethanoic acid;

- 2-[[cyclohexylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
2-[[[(cyclohexyl)methylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
2-[[phenylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
5 2-[[benzylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
2-[[phenylethylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
2-[[phenylpropylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
2-[[phenylbutylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
10 2-[[[(2, 3, 4-trimethoxyphenyl)-3-hydroxyphosphinyl]amino]-2-phenylethanoic acid;
2-[[[(1-naphthyl)hydroxyphosphinyl]amino]-2-phenylethanoic acid;
2-[[[(2-naphthyl)hydroxyphosphinyl]amino]-2-phenylethanoic acid;
15 2-[[[(1-naphthyl)methylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
2-[[[(2-naphthyl)methylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
20 2-[[[(1-naphthyl)ethylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
2-[[[(2-naphthyl)ethylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
2-[[[(1-naphthyl)propylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
25 2-[[[(2-naphthyl)propylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
2-[[[(1-naphthyl)butylhydroxyphosphinyl]amino]-2-phenylethanoic

- acid;
- 2-[[(2-naphthyl)butylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
- 2-[[phenylprop-2-enylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
- 5 2-[[benzylhydroxyphosphinyl]amino]-2-methylethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-ethylethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-propylethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-butylethanoic acid;
- 10 2-[[benzylhydroxyphosphinyl]amino]-2-cyclohexylethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-(cyclohexyl)methyl ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-benzylethanoic acid;
- 15 2-[[benzylhydroxyphosphinyl]amino]-2-phenylethylethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-phenylpropylethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-phenylbutylethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-(2, 3, 4-
- 20 trimethoxyphenyl)ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-(1-naphthyl)ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-(2-naphthyl)ethanoic acid;
- 25 2-[[benzylhydroxyphosphinyl]amino]-2-(1-naphthyl)methyl ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-(2-naphthyl)methyl ethanoic acid;

- 2-[[benzylhydroxyphosphinyl]amino]-2-(1-naphthyl)ethylethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-(2-naphthyl)ethylethanoic acid;
- 5 2-[[benzylhydroxyphosphinyl]amino]-2-(1-naphthyl)propyl ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-(2-naphthyl)propyl ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-(1-naphthyl)butylethanoic acid;
- 10 2-[[benzylhydroxyphosphinyl]amino]-2-(2-naphthyl)butylethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-phenolprop-2-enylethanoic acid;
- 15 2-[(methylhydroxyphosphinyl)amino]hexanedioic acid;
- 2-[(benzylhydroxyphosphinyl)amino]hexanedioic acid;
- 2-[(methylhydroxyphosphinyl)amino]heptanedioic acid;
- 2-[(benzylhydroxyphosphinyl)amino]heptanedioic acid;
- 2-[(methylhydroxyphosphinyl)amino]octanedioic acid;
- 20 2-[(benzylhydroxyphosphinyl)amino]octanedioic acid;
- 2-[(methylhydroxyphosphinyl)amino]nonanedioic acid;
- 2-[(benzylhydroxyphosphinyl)amino]nonanedioic acid;
- 2-[(methylhydroxyphosphinyl)amino]decanedioic acid;
- 2-[(benzylhydroxyphosphinyl)amino]decanedioic acid;
- 25 2-[[[(2-pyridyl)methylhydroxyphosphinyl]amino]pentanedioic acid;
- 2-[[[(3-pyridyl)methylhydroxyphosphinyl]amino]pentanedioic acid;

- 2-[[[(4-pyridyl)methylhydroxyphosphinyl]amino]pentanedioic acid;
- 2-[[[(3-pyridyl)ethylhydroxyphosphinyl]amino]pentanedioic acid;
- 2-[[[(3-pyridyl)propylhydroxyphosphinyl]amino]pentanedioic acid;
- 5 2-[[[(tetrahydrofuranyl)methylhydroxyphosphinyl]amino]pentanedioic acid;
- 2-[[[(tetrahydrofuranyl)ethylhydroxyphosphinyl]amino]pentanedioic acid;
- 10 2-[[[(tetrahydrofuranyl)propylhydroxyphosphinyl]amino]pentanedioic acid;
- 2-[[[(2-indolyl)methylhydroxyphosphinyl]amino]pentanedioic acid;
- 2-[[[(3-indolyl)methylhydroxyphosphinyl]amino]pentanedioic acid;
- 15 2-[[[(4-indolyl)methylhydroxyphosphinyl]amino]pentanedioic acid;
- 2-[[[(3-indolyl)ethylhydroxyphosphinyl]amino]pentanedioic acid;
- 2-[[[(3-indolyl)propylhydroxyphosphinyl]amino]pentanedioic acid;
- 20 2-[[[(2-thienyl)methylhydroxyphosphinyl]amino]pentanedioic acid;
- 2-[[[(3-thienyl)methylhydroxyphosphinyl]amino]pentanedioic acid;
- 25 2-[[[(4-thienyl)methylhydroxyphosphinyl]amino]pentanedioic acid;
- 2-[[[(3-thienyl)ethylhydroxyphosphinyl]amino]pentanedioic acid;
- 2-[[[(3-thienyl)propylhydroxyphosphinyl]amino]pentanedioic

acid;

2-[[[(2-pyridyl)methylhydroxyphosphinyl]amino]-2-phenylethanoic
acid;

5 2-[[[(3-pyridyl)methylhydroxyphosphinyl]amino]-2-phenylethanoic
acid;

2-[[[(4-pyridyl)methylhydroxyphosphinyl]amino]-2-phenylethanoic
acid;

2-[[[(3-pyridyl)ethylhydroxyphosphinyl]amino]-2-phenylethanoic
acid;

10 2-[[[(3-pyridyl)propylhydroxyphosphinyl]amino]-2-phenylethanoic
acid;

2-[[[(tetrahydrofuranyl)methylhydroxyphosphinyl]amino]-2-phenyl
ethanoic acid;

15 2-[[[(tetrahydrofuranyl)ethylhydroxyphosphinyl]amino]-2-phenyl
ethanoic acid;

2-[[[(tetrahydrofuranyl)propylhydroxyphosphinyl]amino]-2-phenyl
ethanoic acid;

2-[[[(2-indolyl)methylhydroxyphosphinyl]amino]-2-phenylethanoic
acid;

20 2-[[[(3-indolyl)methylhydroxyphosphinyl]amino]-2-phenylethanoic
acid;

2-[[[(4-indolyl)methylhydroxyphosphinyl]amino]-2-phenylethanoic
acid;

25 2-[[[(3-indolyl)ethylhydroxyphosphinyl]amino]-2-phenylethanoic
acid;

2-[[[(3-indolyl)propylhydroxyphosphinyl]amino]-2-phenylethanoic
acid;

2-[[[(2-thienyl)methylhydroxyphosphinyl]amino]-2-phenylethanoic

- acid;
- 2-[[[(3-thienyl)methylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
- 2-[[[(4-thienyl)methylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
- 5 2-[[[(3-thienyl)ethylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
- 2-[[[(3-thienyl)propylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
- 10 2-[[benzylhydroxyphosphinyl]amino]-2-(2-pyridyl)methylethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-(3-pyridyl)methylethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-(4-pyridyl)methylethanoic acid;
- 15 2-[[benzylhydroxyphosphinyl]amino]-2-(3-pyridyl)ethylethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-(3-pyridyl)propylethanoic acid;
- 20 2-[[benzylhydroxyphosphinyl]amino]-2-(tetrahydrofuran-2-yl)methylethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-(tetrahydrofuran-2-yl)ethylethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-(tetrahydrofuran-2-yl)propylethanoic acid;
- 25 2-[[benzylhydroxyphosphinyl]amino]-2-(2-indolyl)methylethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-(3-indolyl)methylethanoic acid;

acid;

2-[[benzylhydroxyphosphinyl]amino]-2-(4-indolyl)methylethanoic acid;

2-[[benzylhydroxyphosphinyl]amino]-2-(3-indolyl)ethylethanoic acid;

2-[[benzylhydroxyphosphinyl]amino]-2-(3-indolyl)propylethanoic acid;

2-[[benzylhydroxyphosphinyl]amino]-2-(2-thienyl)methylethanoic acid;

2-[[benzylhydroxyphosphinyl]amino]-2-(3-thienyl)methylethanoic acid;

2-[[benzylhydroxyphosphinyl]amino]-2-(4-thienyl)methylethanoic acid;

2-[[benzylhydroxyphosphinyl]amino]-2-(3-thienyl)ethylethanoic acid;

2-[[benzylhydroxyphosphinyl]amino]-2-(3-thienyl)propylethanoic acid; and a pharmaceutically acceptable salt, hydrate, or a mixture thereof.

45. The method of claim 43 wherein the NAALADase inhibitor is selected from the group consisting of:

2-[(benzylhydroxyphosphinyl)amino]pentanedioic acid;

2-[(phenylhydroxyphosphinyl)amino]pentanedioic acid;

2-[[[(hydroxy)phenylmethyl]hydroxyphosphinyl]amino]pentanedioic acid;

2-[(butylhydroxyphosphinyl)amino]pentanedioic acid;

2-[[[(3-methylbenzyl)hydroxyphosphinyl]amino]pentanedioic acid;

2-[(3-phenylpropylhydroxyphosphinyl)amino]pentanedioic acid;

2-[[[(4-fluorophenyl)hydroxyphosphinyl]amino]pentanedioic acid;

2-[(methylhydroxyphosphinyl)amino]pentanedioic acid;
2-[(phenylethylhydroxyphosphinyl)amino]pentanedioic acid;
2-[[[(4-methylbenzyl)hydroxyphosphinyl]amino]pentanedioic acid;
2-[[[(4-fluorobenzyl)hydroxyphosphinyl]amino]pentanedioic acid;
5 2-[[[(4-methoxybenzyl)hydroxyphosphinyl]amino]pentanedioic
acid;
2-[(phosphono)amino]pentanedioic acid;
2-[[[(3-trifluoromethylbenzyl)hydroxyphosphinyl]amino]
pentanedioic acid;
10 2-[[[(2-fluorobenzyl)hydroxyphosphinyl]amino]pentanedioic acid;
2-[[[(pentafluorobenzyl)hydroxyphosphinyl]amino]pentanedioic
acid; and a pharmaceutically acceptable salt, hydrate, or a
mixture thereof.

46. The method of claim 36 wherein R_1 or R_2 is
15 heterocyclic and X is CH_2 .

47. The method of claim 46 wherein the NAALADase
inhibitor is selected from the group consisting of:
2-[[[(2-pyridyl)hydroxyphosphinyl]methyl]pentanedioic acid;
2-[[[(3-pyridyl)hydroxyphosphinyl]methyl]pentanedioic acid;
20 2-[[[(4-pyridyl)hydroxyphosphinyl]methyl]pentanedioic acid;
2-[[[(tetrahydrofuranyl)hydroxyphosphinyl]methyl]pentanedioic
acid;
2-[[[(2-indolyl)hydroxyphosphinyl]methyl]pentanedioic acid;
2-[[[(3-indolyl)hydroxyphosphinyl]methyl]pentanedioic acid;
25 2-[[[(4-indolyl)hydroxyphosphinyl]methyl]pentanedioic acid;
2-[[[(2-thienyl)hydroxyphosphinyl]methyl]pentanedioic acid;
2-[[[(3-thienyl)hydroxyphosphinyl]methyl]pentanedioic acid;
2-[[[(4-thienyl)hydroxyphosphinyl]methyl]pentanedioic acid;

3-[(2-pyridyl)hydroxyphosphinyl]-2-phenylpropanoic acid;
3-[(3-pyridyl)hydroxyphosphinyl]-2-phenylpropanoic acid;
3-[(4-pyridyl)hydroxyphosphinyl]-2-phenylpropanoic acid;
3-[(tetrahydrofuranyl)hydroxyphosphinyl]-2-phenylpropanoic
5 acid;
3-[(2-indolyl)hydroxyphosphinyl]-2-phenylpropanoic acid;
3-[(3-indolyl)hydroxyphosphinyl]-2-phenylpropanoic acid;
3-[(4-indolyl)hydroxyphosphinyl]-2-phenylpropanoic acid;
3-[(2-thienyl)hydroxyphosphinyl]-2-phenylpropanoic acid;
10 3-[(3-thienyl)hydroxyphosphinyl]-2-phenylpropanoic acid;
3-[(4-thienyl)hydroxyphosphinyl]-2-phenylpropanoic acid;
3-(benzylhydroxyphosphinyl)-2-(2-pyridyl)propanoic acid;
3-(benzylhydroxyphosphinyl)-2-(3-pyridyl)propanoic acid;
3-(benzylhydroxyphosphinyl)-2-(4-pyridyl)propanoic acid;
15 3-(benzylhydroxyphosphinyl)-2-(tetrahydrofuranyl)propanoic
acid;
3-(benzylhydroxyphosphinyl)-2-(2-indolyl)propanoic acid;
3-(benzylhydroxyphosphinyl)-2-(3-indolyl)propanoic acid;
3-(benzylhydroxyphosphinyl)-2-(4-indolyl)propanoic acid;
20 3-(benzylhydroxyphosphinyl)-2-(2-thienyl)propanoic acid;
3-(benzylhydroxyphosphinyl)-2-(3-thienyl)propanoic acid;
3-(benzylhydroxyphosphinyl)-2-(4-thienyl)propanoic acid; and
a pharmaceutically acceptable salt, hydrate, or a mixture
thereof.

25 48. The method of claim 36 wherein R₁ or R₂ is
heterocyclic and X is oxygen.

49. The method of claim 48 wherein the NAALADase
inhibitor is selected from the group consisting of:

- 2-[[[(2-pyridyl)hydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(3-pyridyl)hydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(4-pyridyl)hydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(tetrahydrofuranyl)hydroxyphosphinyl]oxy]pentanedioic
5 acid;
2-[[[(2-indolyl)hydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(3-indolyl)hydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(4-indolyl)hydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(2-thienyl)hydroxyphosphinyl]oxy]pentanedioic acid;
10 2-[[[(3-thienyl)hydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(4-thienyl)hydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(2-pyridyl)hydroxyphosphinyl]oxy]-2-phenylethanoic acid;
2-[[[(3-pyridyl)hydroxyphosphinyl]oxy]-2-phenylethanoic acid;
2-[[[(4-pyridyl)hydroxyphosphinyl]oxy]-2-phenylethanoic acid;
15 2-[[[(tetrahydrofuranyl)hydroxyphosphinyl]oxy]-2-phenylethanoic
acid;
2-[[[(2-indolyl)hydroxyphosphinyl]oxy]-2-phenylethanoic acid;
2-[[[(3-indolyl)hydroxyphosphinyl]oxy]-2-phenylethanoic acid;
2-[[[(4-indolyl)hydroxyphosphinyl]oxy]-2-phenylethanoic acid;
20 2-[[[(2-thienyl)hydroxyphosphinyl]oxy]-2-phenylethanoic acid;
2-[[[(3-thienyl)hydroxyphosphinyl]oxy]-2-phenylethanoic acid;
2-[[[(4-thienyl)hydroxyphosphinyl]oxy]-2-phenylethanoic acid;
2-[[[benzylhydroxyphosphinyl]oxy]-2-(2-pyridyl)ethanoic acid;
2-[[[benzylhydroxyphosphinyl]oxy]-2-(3-pyridyl)ethanoic acid;
25 2-[[[benzylhydroxyphosphinyl]oxy]-2-(4-pyridyl)ethanoic acid;
2-[[[benzylhydroxyphosphinyl]oxy]-2-(tetrahydrofuranyl)ethanoic
acid;
2-[[[benzylhydroxyphosphinyl]oxy]-2-(2-indolyl)ethanoic acid;

2-[[benzylhydroxyphosphinyl]oxy]-2-(3-indolyl)ethanoic acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-(4-indolyl)ethanoic acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-(2-thienyl)ethanoic acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-(3-thienyl)ethanoic acid;
5 2-[[benzylhydroxyphosphinyl]oxy]-2-(4-thienyl)ethanoic acid;
and a pharmaceutically acceptable salt, hydrate, or a mixture thereof.

50. The method of claim 36 wherein R_1 or R_2 is heterocyclic and X is NR_1 .

10 51. The method of claim 50 wherein the NAALADase inhibitor is a compound selected from the group consisting of:
2-[[[(2-pyridyl)hydroxyphosphinyl]amino]pentanedioic acid;
2-[[[(3-pyridyl)hydroxyphosphinyl]amino]pentanedioic acid;
2-[[[(4-pyridyl)hydroxyphosphinyl]amino]pentanedioic acid;
15 2-[[[(tetrahydrofuranyl)hydroxyphosphinyl]amino]pentanedioic acid;
2-[[[(2-indolyl)hydroxyphosphinyl]amino]pentanedioic acid;
2-[[[(3-indolyl)hydroxyphosphinyl]amino]pentanedioic acid;
2-[[[(4-indolyl)hydroxyphosphinyl]amino]pentanedioic acid;
20 2-[[[(2-thienyl)hydroxyphosphinyl]amino]pentanedioic acid;
2-[[[(3-thienyl)hydroxyphosphinyl]amino]pentanedioic acid;
2-[[[(4-thienyl)hydroxyphosphinyl]amino]pentanedioic acid;
2-[[[(2-pyridyl)hydroxyphosphinyl]amino]-2-phenylethanoic acid;
2-[[[(3-pyridyl)hydroxyphosphinyl]amino]-2-phenylethanoic acid;
25 2-[[[(4-pyridyl)hydroxyphosphinyl]amino]-2-phenylethanoic acid;
2-[[[(tetrahydrofuranyl)hydroxyphosphinyl]amino]-2-phenyl ethanoic acid;
2-[[[(2-indolyl)hydroxyphosphinyl]amino]-2-phenylethanoic acid;

2-[[(3-indolyl)hydroxyphosphinyl]amino]-2-phenylethanoic acid;
2-[[(4-indolyl)hydroxyphosphinyl]amino]-2-phenylethanoic acid;
2-[[(2-thienyl)hydroxyphosphinyl]amino]-2-phenylethanoic acid;
2-[[(3-thienyl)hydroxyphosphinyl]amino]-2-phenylethanoic acid;
5 2-[[(4-thienyl)hydroxyphosphinyl]amino]-2-phenylethanoic acid;
2-[[benzylhydroxyphosphinyl]amino]-2-(2-pyridyl)ethanoic acid;
2-[[benzylhydroxyphosphinyl]amino]-2-(3-pyridyl)ethanoic acid;
2-[[benzylhydroxyphosphinyl]amino]-2-(4-pyridyl)ethanoic acid;
2-[[benzylhydroxyphosphinyl]amino]-2-(tetrahydrofuranyl)
10 ethanoic acid;
2-[[benzylhydroxyphosphinyl]amino]-2-(2-indolyl)ethanoic acid;
2-[[benzylhydroxyphosphinyl]amino]-2-(3-indolyl)ethanoic acid;
2-[[benzylhydroxyphosphinyl]amino]-2-(4-indolyl)ethanoic acid;
2-[[benzylhydroxyphosphinyl]amino]-2-(2-thienyl)ethanoic acid;
15 2-[[benzylhydroxyphosphinyl]amino]-2-(3-thienyl)ethanoic acid;
2-[[benzylhydroxyphosphinyl]amino]-2-(4-thienyl)ethanoic acid;
and a pharmaceutically acceptable salt, hydrate, or a mixture thereof.

52. A method of inhibiting NAALADase enzyme activity in
20 an animal which comprises: administering to an animal
suffering from a disorder related to NAALADase enzyme activity
an effective amount of a NAALADase inhibitor.

53. The method of claim 52 wherein the NAALADase
inhibitor is selected from the group consisting of: a
25 glutamate-derived hydroxyphosphinyl derivative compound; an
acidic peptide analog; a conformationally restricted glutamate
mimic; and mixtures thereof.

54. The method of claim 53 wherein the acidic peptide

analog is selected from the group consisting of: Asp-Glu, Glu-Glu, Gly-Glu, gamma-Glu-Glu, and Glu-Glu-Glu.


55. The method of claim 53 wherein the conformationally restricted glutamate mimic is selected from the group consisting of: beta:NAAG and quisqualic acid.

56. The method of claim 53 wherein the NAALADase inhibitor is a glutamate-derived hydroxyphosphinyl derivative compound.

57. The method of claim 52 wherein the NAALADase inhibitor is administered in combination with an additional therapeutic agent selected from the group consisting of: therapeutic hormones, chemotherapeutic hormones, anti-angiogenesis agents, radiolabelled compounds, and mixtures thereof.

58. The method of claim 52 wherein the disorder related to NAALADase enzyme activity is prostate disease wherein the prostate disease is selected from the group consisting of prostate cancer and benign prostatic hyperplasia.

59. The method of claim 52 wherein the NAALADase inhibitor comprises a compound having the following formula:



I

wherein

R₁ is hydrogen, C₁-C₈ straight or branched chain alkyl, C₂-C₈ straight or branched chain alkenyl group, C₃-C₈

cycloalkyl, C₅-C₇ cycloalkenyl, or Ar₁;

X is CH₂, O, or NR₁, where R₁ is defined above; and

R₂ is C₁-C₈ straight or branched chain alkyl, C₂-C₈ straight or branched chain alkenyl group, C₁-C₈ cycloalkyl, C₅-C₇ cycloalkenyl, or Ar₁, wherein said alkyl, alkenyl, cycloalkyl, cycloalkenyl or aryl group may be optionally substituted with carboxylic acid,

wherein said alkyl, alkenyl, cycloalkyl, cycloalkenyl or aryl groups may be optionally substituted with C₁-C₈ cycloalkyl, C₁ or C₅ cycloalkyl, C₅-C₇ cycloalkenyl, C₁-C₄ alkyl, C₁-C₄ alkenyl, halo, hydroxy, carboxy, nitro, trifluoromethyl, C₁-C₆ straight or branched chain alkyl or alkenyl, C₁-C₄ alkoxy, C₁-C₄ alkenyloxy, phenoxy, benzyloxy, amino, or Ar₁, and where Ar₁ is selected from the group consisting of 1-naphthyl, 2-naphthyl, 2-indolyl, 3-indolyl, 4-indolyl, 2-furyl, 3-furyl, tetrahydrofuranyl, 2-thienyl, 3-thienyl, 4-thienyl, 2-, 3-, or 4-pyridyl, or phenyl, having one to five substituents which are independently selected from the group consisting of hydrogen, halo, hydroxy, carboxy, nitro, trifluoromethyl, C₁-C₈ straight or branched alkyl or alkenyl, C₁-C₄ alkoxy or C₁-C₄ alkenyloxy, phenoxy, benzyloxy, and amino; or pharmaceutically acceptable salts, hydrates, or mixtures thereof.

60. The method of claim 59 wherein R₁ and R₂ are straight or branched aliphatic groups or carbocyclic groups and X is CH₂.

61. The method of claim 60 wherein the NAALADase inhibitor is selected from the group consisting of:

- 2-[[methylhydroxyphosphinyl]methyl]pentanedioic acid;
2-[[ethylhydroxyphosphinyl]methyl]pentanedioic acid;
2-[[propylhydroxyphosphinyl]methyl]pentanedioic acid;
2-[[butylhydroxyphosphinyl]methyl]pentanedioic acid;
5 2-[[cyclohexylhydroxyphosphinyl]methyl]pentanedioic acid;
2-[[((cyclohexyl)methylhydroxyphosphinyl)methyl]pentanedioic acid;
2-[[phenylhydroxyphosphinyl]methyl]pentanedioic acid;
2-[[benzylhydroxyphosphinyl]methyl]pentanedioic acid;
10 2-[[phenylethylhydroxyphosphinyl]methyl]pentanedioic acid;
2-[[phenylpropylhydroxyphosphinyl]methyl]pentanedioic acid;
2-[[phenylbutylhydroxyphosphinyl]methyl]pentanedioic acid;
2-[[((4-methylbenzyl)hydroxyphosphinyl)methyl]pentanedioic acid;
15 2-[[((4-fluorobenzyl)hydroxyphosphinyl)methyl]pentanedioic acid;
2-[[((2-fluorobenzyl)hydroxyphosphinyl)methyl]pentanedioic acid;
2-[[((pentafluorobenzyl)hydroxyphosphinyl)methyl]pentanedioic acid;
20 2-[[((methoxybenzyl)hydroxyphosphinyl)methyl]pentanedioic acid;
2-[[((2, 3, 4-trimethoxyphenyl)hydroxyphosphinyl)methyl]pentanedioic acid;
2-[[((1-naphthyl)hydroxyphosphinyl)methyl]pentanedioic acid;
25 2-[[((2-naphthyl)hydroxyphosphinyl)methyl]pentanedioic acid;
2-[[((1-naphthyl)methylhydroxyphosphinyl)methyl]pentanedioic acid;
2-[[((2-naphthyl)methylhydroxyphosphinyl)methyl]pentanedioic

- acid;
- 2-[[[(1-naphthyl)ethylhydroxyphosphinyl]methyl]pentanedioic acid;
- 2-[[[(2-naphthyl)ethylhydroxyphosphinyl]methyl]pentanedioic acid;
- 5 acid;
- 2-[[[(1-naphthyl)propylhydroxyphosphinyl]methyl]pentanedioic acid;
- 2-[[[(2-naphthyl)propylhydroxyphosphinyl]methyl]pentanedioic acid;
- 10 2-[[[(1-naphthyl)butylhydroxyphosphinyl]methyl]pentanedioic acid;
- 2-[[[(2-naphthyl)butylhydroxyphosphinyl]methyl]pentanedioic acid;
- 2-[[[(phenylprop-2-enyl)hydroxyphosphinyl]methyl]pentanedioic acid;
- 15 acid;
- 2-[[[(2-fluorobenzyl)hydroxyphosphinyl]methyl]pentanedioic acid;
- 2-[[[(4-fluorophenyl)hydroxyphosphinyl]methyl]pentanedioic acid;
- 20 2-[[[(hydroxy)phenylmethyl]hydroxyphosphinyl]methyl]pentanedioic acid;
- 2-[[[(3-methylbenzyl)hydroxyphosphinyl]methyl]pentanedioic acid;
- 2- (phosphonomethyl)pentanedioic acid;
- 25 2-[[[(3-trifluoromethylbenzyl)hydroxyphosphinyl]methyl]pentanedioic acid;
- 3- (methylhydroxyphosphinyl) -2-phenylpropanoic acid;
- 3- (ethylhydroxyphosphinyl) -2-phenylpropanoic acid;

- 3-(propylhydroxyphosphinyl)-2-phenylpropanoic acid;
3-(butylhydroxyphosphinyl)-2-phenylpropanoic acid;
3-(cyclohexylhydroxyphosphinyl)-2-phenylpropanoic acid;
3-((cyclohexyl)methylhydroxyphosphinyl)-2-phenylpropanoic
5 acid;
3-(phenylhydroxyphosphinyl)-2-phenylpropanoic acid;
3-(benzylhydroxyphosphinyl)-2-phenylpropanoic acid;
3-(phenylethylhydroxyphosphinyl)-2-phenylpropanoic acid;
3-(phenylpropylhydroxyphosphinyl)-2-phenylpropanoic acid;
10 3-(phenylbutylhydroxyphosphinyl)-2-phenylpropanoic acid;
3-((2, 3, 4-trimethoxyphenyl)-3-hydroxyphosphinyl)-2-phenylpropanoic acid;
3-((1-naphthyl)hydroxyphosphinyl)-2-phenylpropanoic acid;
3-((2-naphthyl)hydroxyphosphinyl)-2-phenylpropanoic acid;
15 3-((1-naphthyl)methylhydroxyphosphinyl)-2-phenylpropanoic acid;
3-((2-naphthyl)methylhydroxyphosphinyl)-2-phenylpropanoic acid;
3-((1-naphthyl)ethylhydroxyphosphinyl)-2-phenylpropanoic acid;
20 3-((2-naphthyl)ethylhydroxyphosphinyl)-2-phenylpropanoic acid;
3-((1-naphthyl)propylhydroxyphosphinyl)-2-phenylpropanoic acid;
3-((2-naphthyl)propylhydroxyphosphinyl)-2-phenylpropanoic acid;
25 3-((1-naphthyl)butylhydroxyphosphinyl)-2-phenylpropanoic acid;
3-((2-naphthyl)butylhydroxyphosphinyl)-2-phenylpropanoic acid;
3-(phenylprop-2-enylhydroxyphosphinyl)-2-phenylpropanoic acid;
2-[(benzylhydroxyphosphinyl)methyl]pentanedioic acid;

- 2-[(methylhydroxyphosphinyl)methyl]hexanedioic acid;
2-[(benzylhydroxyphosphinyl)methyl]hexanedioic acid;
2-[(methylhydroxyphosphinyl)methyl]heptanedioic acid;
2-[(benzylhydroxyphosphinyl)methyl]heptanedioic acid;
5 2-[(methylhydroxyphosphinyl)methyl]octanedioic acid;
2-[(benzylhydroxyphosphinyl)methyl]octanedioic acid;
2-[(methylhydroxyphosphinyl)methyl]nonanedioic acid;
2-[(benzylhydroxyphosphinyl)methyl]nonanedioic acid;
2-[(methylhydroxyphosphinyl)methyl]decanedioic acid;
10 2-[(benzylhydroxyphosphinyl)methyl]decanedioic acid;
3-(benzylhydroxyphosphinyl)-2-methylpropanoic acid;
3-(benzylhydroxyphosphinyl)-2-ethylpropanoic acid;
3-(benzylhydroxyphosphinyl)-2-propylpropanoic acid;
3-(benzylhydroxyphosphinyl)-2-butylpropanoic acid;
15 3-(benzylhydroxyphosphinyl)-2-cyclohexylpropanoic acid;
3-(benzylhydroxyphosphinyl)-2-(cyclohexyl)methylpropanoic acid;
3-(benzylhydroxyphosphinyl)-2-phenylpropanoic acid;
3-(benzylhydroxyphosphinyl)-2-benzylpropanoic acid;
20 3-(benzylhydroxyphosphinyl)-2-phenylethylpropanoic acid;
3-(benzylhydroxyphosphinyl)-2-phenylpropylpropanoic acid;
3-(benzylhydroxyphosphinyl)-2-phenylbutylpropanoic acid;
3-(benzylhydroxyphosphinyl)-2-(2, 3, 4-trimethoxyphenyl)propanoic acid;
25 3-(benzylhydroxyphosphinyl)-2-(1-naphthyl)propanoic acid;
3-(benzylhydroxyphosphinyl)-2-(2-naphthyl)propanoic acid;
3-(benzylhydroxyphosphinyl)-2-(1-naphthyl)methylpropanoic acid;

- 3-(benzylhydroxyphosphinyl)-2-(2-naphthyl)methylpropanoic acid;
- 3-(benzylhydroxyphosphinyl)-2-(1-naphthyl)ethylpropanoic acid;
- 3-(benzylhydroxyphosphinyl)-2-(2-naphthyl)ethylpropanoic acid;
- 5 3-(benzylhydroxyphosphinyl)-2-(1-naphthyl)propylpropanoic acid;
- 3-(benzylhydroxyphosphinyl)-2-(2-naphthyl)propylpropanoic acid;
- 3-(benzylhydroxyphosphinyl)-2-(1-naphthyl)butylpropanoic acid;
- 10 3-(benzylhydroxyphosphinyl)-2-(2-naphthyl)butylpropanoic acid;
- 3-(benzylhydroxyphosphinyl)-2-phenylprop-2-enylpropanoic acid;
- 2-[[(2-pyridyl)methylhydroxyphosphinyl]methyl]pentanedioic acid;
- 2-[[(3-pyridyl)methylhydroxyphosphinyl]methyl]pentanedioic acid;
- 15 2-[[(4-pyridyl)methylhydroxyphosphinyl]methyl]pentanedioic acid;
- 2-[[(3-pyridyl)ethylhydroxyphosphinyl]methyl]pentanedioic acid;
- 20 2-[[(3-pyridyl)propylhydroxyphosphinyl]methyl]pentanedioic acid;
- 2-[[(tetrahydrofuranyl)methylhydroxyphosphinyl]methyl]pentanedioic acid;
- 2-[[(tetrahydrofuranyl)ethylhydroxyphosphinyl]methyl]pentanedioic acid;
- 25 2-[[(tetrahydrofuranyl)propylhydroxyphosphinyl]methyl]pentanedioic acid;
- 2-[[(2-indolyl)methylhydroxyphosphinyl]methyl]pentanedioic

- acid;
- 2-[[[(3-indolyl)methylhydroxyphosphinyl]methyl]pentanedioic acid;
- 2-[[[(4-indolyl)methylhydroxyphosphinyl]methyl]pentanedioic acid;
- 5 2-[[[(3-indolyl)ethylhydroxyphosphinyl]methyl]pentanedioic acid;
- 2-[[[(3-indolyl)propylhydroxyphosphinyl]methyl]pentanedioic acid;
- 10 2-[[[(2-thienyl)methylhydroxyphosphinyl]methyl]pentanedioic acid;
- 2-[[[(3-thienyl)methylhydroxyphosphinyl]methyl]pentanedioic acid;
- 2-[[[(4-thienyl)methylhydroxyphosphinyl]methyl]pentanedioic acid;
- 15 2-[[[(3-thienyl)ethylhydroxyphosphinyl]methyl]pentanedioic acid;
- 2-[[[(3-thienyl)propylhydroxyphosphinyl]methyl]pentanedioic acid;
- 20 3-[(2-pyridyl)methylhydroxyphosphinyl]-2-phenylpropanoic acid;
- 3-[(3-pyridyl)methylhydroxyphosphinyl]-2-phenylpropanoic acid;
- 3-[(4-pyridyl)methylhydroxyphosphinyl]-2-phenylpropanoic acid;
- 3-[(3-pyridyl)ethylhydroxyphosphinyl]-2-phenylpropanoic acid;
- 3-[(3-pyridyl)propylhydroxyphosphinyl]-2-phenylpropanoic acid;
- 25 3-[(tetrahydrofuranyl)methylhydroxyphosphinyl]-2-phenyl propanoic acid;
- 3-[(tetrahydrofuranyl)ethylhydroxyphosphinyl]-2-phenyl propanoic acid;

- 3-[(tetrahydrofuranyl)propylhydroxyphosphinyl]-2-phenyl
propanoic acid;
- 3-[(2-indolyl)methylhydroxyphosphinyl]-2-phenylpropanoic acid;
- 3-[(3-indolyl)methylhydroxyphosphinyl]-2-phenylpropanoic acid;
- 5 3-[(4-indolyl)methylhydroxyphosphinyl]-2-phenylpropanoic acid;
- 3-[(3-indolyl)ethylhydroxyphosphinyl]-2-phenylpropanoic acid;
- 3-[(3-indolyl)propylhydroxyphosphinyl]-2-phenylpropanoic acid;
- 3-[(2-thienyl)methylhydroxyphosphinyl]-2-phenylpropanoic acid;
- 3-[(3-thienyl)methylhydroxyphosphinyl]-2-phenylpropanoic acid;
- 10 3-[(4-thienyl)methylhydroxyphosphinyl]-2-phenylpropanoic acid;
- 3-[(3-thienyl)ethylhydroxyphosphinyl]-2-phenylpropanoic acid;
- 3-[(3-thienyl)propylhydroxyphosphinyl]-2-phenylpropanoic acid;
- 3-(benzylhydroxyphosphinyl)-2-(2-pyridyl)methylpropanoic acid;
- 3-(benzylhydroxyphosphinyl)-2-(3-pyridyl)methylpropanoic acid;
- 15 3-(benzylhydroxyphosphinyl)-2-(4-pyridyl)methylpropanoic acid;
- 3-(benzylhydroxyphosphinyl)-2-(3-pyridyl)ethylpropanoic acid;
- 3-(benzylhydroxyphosphinyl)-2-(3-pyridyl)propylpropanoic acid;
- 3-(benzylhydroxyphosphinyl)-2-(tetrahydrofuranyl)methyl
propanoic acid;
- 20 3-(benzylhydroxyphosphinyl)-2-(tetrahydrofuranyl)ethyl
propanoic acid;
- 3-(benzylhydroxyphosphinyl)-2-(tetrahydrofuranyl)propyl
propanoic acid;
- 3-(benzylhydroxyphosphinyl)-2-(2-indolyl)methylpropanoic acid;
- 25 3-(benzylhydroxyphosphinyl)-2-(3-indolyl)methylpropanoic acid;
- 3-(benzylhydroxyphosphinyl)-2-(4-indolyl)methylpropanoic acid;
- 3-(benzylhydroxyphosphinyl)-2-(3-indolyl)ethylpropanoic acid;
- 3-(benzylhydroxyphosphinyl)-2-(3-indolyl)propylpropanoic acid;

3-(benzylhydroxyphosphinyl)-2-(2-thienyl)methylpropanoic acid;
3-(benzylhydroxyphosphinyl)-2-(3-thienyl)methylpropanoic acid;
3-(benzylhydroxyphosphinyl)-2-(4-thienyl)methylpropanoic acid;
3-(benzylhydroxyphosphinyl)-2-(3-thienyl)ethylpropanoic acid;
5 3-(benzylhydroxyphosphinyl)-2-(3-thienyl)propylpropanoic acid;
and a pharmaceutically acceptable salt, hydrate, or a mixture thereof.

62. The method of claim 60 wherein the NAALADase inhibitor is selected from the group consisting of:

10 2-[(benzylhydroxyphosphinyl)methyl]pentanedioic acid;
2-[(phenylhydroxyphosphinyl)methyl]pentanedioic acid;
2-[[[(hydroxy)phenylmethyl]hydroxyphosphinyl]methyl]
pentanedioic acid;
2-[(butylhydroxyphosphinyl)methyl]pentanedioic acid;
15 2-[[[(3-methylbenzyl)hydroxyphosphinyl]methyl]pentanedioic
acid;
2-[(3-phenylpropylhydroxyphosphinyl)methyl]pentanedioic acid;
2-[[[(4-fluorophenyl)hydroxyphosphinyl]methyl]pentanedioic
acid;
20 2-[(methylhydroxyphosphinyl)methyl]pentanedioic acid;
2-[(phenylethylhydroxyphosphinyl)methyl]pentanedioic acid;
2-[[[(4-methylbenzyl)hydroxyphosphinyl]methyl]pentanedioic
acid;
2-[[[(4-fluorobenzyl)hydroxyphosphinyl]methyl]pentanedioic
25 acid;
2-[[[(4-methoxybenzyl)hydroxyphosphinyl]methyl]pentanedioic
acid;
2-(phosphonomethyl)pentanedioic acid;

2-[[[(3-trifluoromethylbenzyl)hydroxyphosphinyl)methyl]pentanedioic acid;

2-[[[(2-fluorobenzyl)hydroxyphosphinyl)methyl]pentanedioic acid;

5 2-[[[(pentafluorobenzyl)hydroxyphosphinyl)methyl]pentanedioic acid; and a pharmaceutically acceptable salt, hydrate, or a mixture thereof.

10 63. The method of claim 59 wherein R₁ and R₂ are straight or branched aliphatic groups or carbocyclic groups and X is oxygen.

64. The method of claim 63 wherein the NAALADase inhibitor is selected from the group consisting of:

2-[[methylhydroxyphosphinyl]oxy]pentanedioic acid;

2-[[ethylhydroxyphosphinyl]oxy]pentanedioic acid;

15 2-[[propylhydroxyphosphinyl]oxy]pentanedioic acid;

2-[[butylhydroxyphosphinyl]oxy]pentanedioic acid;

2-[[cyclohexylhydroxyphosphinyl]oxy]pentanedioic acid;

2-[[[(cyclohexyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;

2-[[phenylhydroxyphosphinyl]oxy]pentanedioic acid;

20 2-[[benzylhydroxyphosphinyl]oxy]pentanedioic acid;

2-[[phenylethylhydroxyphosphinyl]oxy]pentanedioic acid;

2-[[phenylpropylhydroxyphosphinyl]oxy]pentanedioic acid;

2-[[phenylbutylhydroxyphosphinyl]oxy]pentanedioic acid;

2-[[[(4-methylbenzyl)hydroxyphosphinyl]oxy]pentanedioic acid;

25 2-[[[(4-fluorobenzyl)hydroxyphosphinyl]oxy]pentanedioic acid;

2-[[[(2-fluorobenzyl)hydroxyphosphinyl]oxy]pentanedioic acid;

2-[[[(pentafluorobenzyl)hydroxyphosphinyl]oxy]pentanedioic acid;

- 2-[[[(methoxybenzyl)hydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(2, 3, 4-trimethoxyphenyl)hydroxyphosphinyl]oxy]
pentanedioic acid;
2-[[[(1-naphthyl)hydroxyphosphinyl]oxy]pentanedioic acid;
5 2-[[[(2-naphthyl)hydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(1-naphthyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(2-naphthyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(1-naphthyl)ethylhydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(2-naphthyl)ethylhydroxyphosphinyl]oxy]pentanedioic acid;
10 2-[[[(1-naphthyl)propylhydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(2-naphthyl)propylhydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(1-naphthyl)butylhydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(2-naphthyl)butylhydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(phenylprop-2-enyl)hydroxyphosphinyl]oxy]pentanedioic
15 acid;
2-[[benzylhydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(4-fluorophenyl)hydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(hydroxy)phenylmethyl]hydroxyphosphinyl]oxy]pentanedioic
acid;
20 2-[[[(3-methylbenzyl)hydroxyphosphinyl]oxy]pentanedioic acid;
2-[(phosphono)oxy]pentanedioic acid;
2-[[[(3-trifluoromethylbenzyl)hydroxyphosphinyl]oxy]
pentanedioic acid;
2-[[methylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
25 2-[[ethylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
2-[[propylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
2-[[butylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
2-[[cyclohexylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;

- 2-[[[(cyclohexyl)methylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[[phenylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[[benzylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 5 2-[[[phenylethylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[[phenylpropylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[[phenylbutylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[[(2, 3, 4-trimethoxyphenyl)-3-hydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 10 2-[[[(2-naphthyl)hydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[[(3-naphthyl)hydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[[(4-naphthyl)hydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[[(2-naphthyl)methylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 15 2-[[[(3-naphthyl)methylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[[(4-naphthyl)methylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[[(3-naphthyl)ethylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 20 2-[[[(3-naphthyl)propylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[[(3-naphthyl)butylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 25 2-[[[phenylprop-2-enylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[(methylhydroxyphosphinyl)oxy]hexanedioic acid;
- 2-[(benzylhydroxyphosphinyl)oxy]hexanedioic acid;

- 2-[(methylhydroxyphosphinyl)oxy]heptanedioic acid;
2-[(benzylhydroxyphosphinyl)oxy]heptanedioic acid;
2-[(methylhydroxyphosphinyl)oxy]octanedioic acid;
2-[(benzylhydroxyphosphinyl)oxy]octanedioic acid;
5 2-[(methylhydroxyphosphinyl)oxy]nonanedioic acid;
2-[(benzylhydroxyphosphinyl)oxy]nonanedioic acid;
2-[(methylhydroxyphosphinyl)oxy]decanedioic acid;
2-[(benzylhydroxyphosphinyl)oxy]decanedioic acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-methylethanoic acid;
10 2-[[benzylhydroxyphosphinyl]oxy]-2-ethylethanoic acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-propylethanoic acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-butylethanoic acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-cyclohexylethanoic acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-(cyclohexyl)methylethanoic
15 acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-benzylethanoic acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-phenylethylethanoic acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-phenylpropylethanoic acid;
20 2-[[benzylhydroxyphosphinyl]oxy]-2-phenylbutylethanoic acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-(2, 3, 4-trimethoxyphenyl)
ethanoic acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-(1-naphthyl)ethanoic acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-(2-naphthyl)ethanoic acid;
25 2-[[benzylhydroxyphosphinyl]oxy]-2-(1-naphthyl)methylethanoic
acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-(2-naphthyl)methylethanoic
acid;

- 2-[[benzylhydroxyphosphinyl]oxy]-2-(1-naphthyl)ethylethanoic acid;
- 2-[[benzylhydroxyphosphinyl]oxy]-2-(2-naphthyl)ethylethanoic acid;
- 5 2-[[benzylhydroxyphosphinyl]oxy]-2-(1-naphthyl)propylethanoic acid;
- 2-[[benzylhydroxyphosphinyl]oxy]-2-(2-naphthyl)propylethanoic acid;
- 2-[[benzylhydroxyphosphinyl]oxy]-2-(1-naphthyl)butylethanoic acid;
- 10 2-[[benzylhydroxyphosphinyl]oxy]-2-(2-naphthyl)butylethanoic acid;
- 2-[[benzylhydroxyphosphinyl]oxy]-2-phenylprop-2-enylethanoic acid;
- 15 2-[[[(2-pyridyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;
- 2-[[[(3-pyridyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;
- 2-[[[(4-pyridyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;
- 2-[[[(3-pyridyl)ethylhydroxyphosphinyl]oxy]pentanedioic acid;
- 2-[[[(3-pyridyl)propylhydroxyphosphinyl]oxy]pentanedioic acid;
- 20 2-[[[(tetrahydrofuranyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;
- 2-[[[(tetrahydrofuranyl)ethylhydroxyphosphinyl]oxy]pentanedioic acid;
- 2-[[[(tetrahydrofuranyl)propylhydroxyphosphinyl]oxy]pentanedioic acid;
- 25 2-[[[(2-indolyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;
- 2-[[[(3-indolyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;
- 2-[[[(4-indolyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;

- 2-[[[(3-indolyl)ethylhydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(3-indolyl)propylhydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(2-thienyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(3-thienyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;
5 2-[[[(4-thienyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(3-thienyl)ethylhydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(3-thienyl)propylhydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(2-pyridyl)methylhydroxyphosphinyl]oxy]-2-phenylethanoic
acid;
10 2-[[[(3-pyridyl)methylhydroxyphosphinyl]oxy]-2-phenylethanoic
acid;
2-[[[(4-pyridyl)methylhydroxyphosphinyl]oxy]-2-phenylethanoic
acid;
2-[[[(3-pyridyl)ethylhydroxyphosphinyl]oxy]-2-phenylethanoic
15 acid;
2-[[[(3-pyridyl)propylhydroxyphosphinyl]oxy]-2-phenylethanoic
acid;
2-[[[(tetrahydrofuranyl)methylhydroxyphosphinyl]oxy]-2-
phenylethanoic acid;
20 2-[[[(tetrahydrofuranyl)ethylhydroxyphosphinyl]oxy]-2-
phenylethanoic acid;
2-[[[(tetrahydrofuranyl)propylhydroxyphosphinyl]oxy]-2-
phenylethanoic acid;
2-[[[(2-indolyl)methylhydroxyphosphinyl]oxy]-2-phenylethanoic
25 acid;
2-[[[(3-indolyl)methylhydroxyphosphinyl]oxy]-2-phenylethanoic
acid;
2-[[[(4-indolyl)methylhydroxyphosphinyl]oxy]-2-phenylethanoic

acid;

2-[[[(3-indolyl)ethylhydroxyphosphinyl]oxy]-2-phenylethanoic
acid;

2-[[[(3-indolyl)propylhydroxyphosphinyl]oxy]-2-phenylethanoic
5 acid;

2-[[[(2-thienyl)methylhydroxyphosphinyl]oxy]-2-phenylethanoic
acid;

2-[[[(3-thienyl)methylhydroxyphosphinyl]oxy]-2-phenylethanoic
acid;

10 2-[[[(4-thienyl)methylhydroxyphosphinyl]oxy]-2-phenylethanoic
acid;

2-[[[(3-thienyl)ethylhydroxyphosphinyl]oxy]-2-phenylethanoic
acid;

2-[[[(3-thienyl)propylhydroxyphosphinyl]oxy]-2-phenylethanoic
15 acid;

2-[[benzylhydroxyphosphinyl]oxy]-2-(2-pyridyl)methylethanoic
acid;

2-[[benzylhydroxyphosphinyl]oxy]-2-(3-pyridyl)methylethanoic
acid;

20 2-[[benzylhydroxyphosphinyl]oxy]-2-(4-pyridyl)methylethanoic
acid;

2-[[benzylhydroxyphosphinyl]oxy]-2-(3-pyridyl)ethylethanoic
acid;

2-[[benzylhydroxyphosphinyl]oxy]-2-(3-pyridyl)propylethanoic
25 acid;

2-[[benzylhydroxyphosphinyl]oxy]-2-(tetrahydrofuran-2-yl)
methylethanoic acid;

2-[[benzylhydroxyphosphinyl]oxy]-2-(tetrahydrofuran-2-yl)

- ethylethanoic acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-(tetrahydrofuran-2-yl)
propylethanoic acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-(2-indolyl)methylethanoic
5 acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-(3-indolyl)methylethanoic
acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-(4-indolyl)methylethanoic
acid;
10 2-[[benzylhydroxyphosphinyl]oxy]-2-(3-indolyl)ethylethanoic
acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-(3-indolyl)propylethanoic
acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-(2-thienyl)methylethanoic
15 acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-(3-thienyl)methylethanoic
acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-(4-thienyl)methylethanoic
acid;
20 2-[[benzylhydroxyphosphinyl]oxy]-2-(3-thienyl)ethylethanoic
acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-(3-thienyl)propylethanoic
acid; and a pharmaceutically acceptable salt, hydrate, or a
mixture thereof.
- 25 65. The method of claim 63 wherein the NAALADase
inhibitor is selected from the group consisting of:
2-[(benzylhydroxyphosphinyl)oxy]pentanedioic acid;
2-[(phenylhydroxyphosphinyl)oxy]pentanedioic acid;

- 2-[[[(hydroxy)phenylmethyl]hydroxyphosphinyl]oxy]pentanedioic acid;
- 2-[[butylhydroxyphosphinyl]oxy]pentanedioic acid;
- 2-[[[(3-methylbenzyl)hydroxyphosphinyl]oxy]pentanedioic acid;
- 5 2-[[[(3-phenylpropyl)hydroxyphosphinyl]oxy]pentanedioic acid;
- 2-[[[(4-fluorophenyl)hydroxyphosphinyl]oxy]pentanedioic acid;
- 2-[[methylhydroxyphosphinyl]oxy]pentanedioic acid;
- 2-[[phenylethylhydroxyphosphinyl]oxy]pentanedioic acid;
- 2-[[[(4-methylbenzyl)hydroxyphosphinyl]oxy]pentanedioic acid;
- 10 2-[[[(4-fluorobenzyl)hydroxyphosphinyl]oxy]pentanedioic acid;
- 2-[[[(4-methoxybenzyl)hydroxyphosphinyl]oxy]pentanedioic acid;
- 2-[[[(2-fluorobenzyl)hydroxyphosphinyl]oxy]pentanedioic acid;
- 2-[[[pentafluorobenzyl)hydroxyphosphinyl]oxy]pentanedioic acid;
- 15 2-[(phosphono)oxy]pentanedioic acid;
- 2-[[[(3-trifluoromethylbenzyl)hydroxyphosphinyl]oxy]pentanedioic acid; and a pharmaceutically acceptable salt, hydrate, or a mixture thereof.

66. The method of claim 59 wherein R_1 and R_2 are straight or branched aliphatic groups or carbocyclic and X is NR_1 .
- 20

67. The method of claim 66 wherein the NAALADase inhibitor is selected from the group consisting of:
- 2-[[methylhydroxyphosphinyl]amino]pentanedioic acid;
- 2-[[ethylhydroxyphosphinyl]amino]pentanedioic acid;
- 25 2-[[propylhydroxyphosphinyl]amino]pentanedioic acid;
- 2-[[butylhydroxyphosphinyl]amino]pentanedioic acid;
- 2-[[cyclohexylhydroxyphosphinyl]amino]pentanedioic acid;
- 2-[[[(cyclohexyl)methylhydroxyphosphinyl]amino]pentanedioic

- acid;
- 2-[[phenylhydroxyphosphinyl]amino]pentanedioic acid;
- 2-[[benzylhydroxyphosphinyl]amino]pentanedioic acid;
- 2-[[phenylethylhydroxyphosphinyl]amino]pentanedioic acid;
- 5 2-[[phenylpropylhydroxyphosphinyl]amino]pentanedioic acid;
- 2-[[phenylbutylhydroxyphosphinyl]amino]pentanedioic acid;
- 2-[[[(4-methylbenzyl)hydroxyphosphinyl]amino]pentanedioic acid;
- 2-[[[(4-fluorobenzyl)hydroxyphosphinyl]amino]pentanedioic acid;
- 2-[[[(2-fluorobenzyl)hydroxyphosphinyl]amino]pentanedioic acid;
- 10 2-[[[(pentafluorobenzyl)hydroxyphosphinyl]amino]pentanedioic acid;
- 2-[[[(methoxybenzyl)hydroxyphosphinyl]amino]pentanedioic acid;
- 2-[[[(2, 3, 4-trimethoxyphenyl)hydroxyphosphinyl]amino]pentanedioic acid;
- 15 2-[[[(1-naphthyl)hydroxyphosphinyl]amino]pentanedioic acid;
- 2-[[[(2-naphthyl)hydroxyphosphinyl]amino]pentanedioic acid;
- 2-[[[(1-naphthyl)methylhydroxyphosphinyl]amino]pentanedioic acid;
- 2-[[[(2-naphthyl)methylhydroxyphosphinyl]amino]pentanedioic acid;
- 20 2-[[[(1-naphthyl)ethylhydroxyphosphinyl]amino]pentanedioic acid;
- 2-[[[(2-naphthyl)ethylhydroxyphosphinyl]amino]pentanedioic acid;
- 25 2-[[[(1-naphthyl)propylhydroxyphosphinyl]amino]pentanedioic acid;
- 2-[[[(2-naphthyl)propylhydroxyphosphinyl]amino]pentanedioic acid;

- 2-[[[(1-naphthyl)butylhydroxyphosphinyl]amino]pentanedioic acid;
- 2-[[[(2-naphthyl)butylhydroxyphosphinyl]amino]pentanedioic acid;
- 5 2-[[[(phenylprop-2-enyl)hydroxyphosphinyl]amino]pentanedioic acid;
- 2-[[benzylhydroxyphosphinyl]amino]pentanedioic acid;
- 2-[[[(4-fluorophenyl)hydroxyphosphinyl]amino]pentanedioic acid;
- 2-[[[(hydroxy)phenylmethyl]hydroxyphosphinyl]amino]pentanedioic acid;
- 10 2-[[[(3-methylbenzyl)hydroxyphosphinyl]amino]pentanedioic acid;
- 2-[(phosphono)amino]pentanedioic acid;
- 2-[[[(3-trifluoromethylbenzyl)hydroxyphosphinyl]amino]pentanedioic acid;
- 15 2-[[methylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
- 2-[[ethylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
- 2-[[propylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
- 2-[[butylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
- 2-[[cyclohexylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
- 20 2-[[[(cyclohexyl)methylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
- 2-[[phenylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
- 2-[[phenylethylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
- 25 2-[[phenylpropylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
- 2-[[phenylbutylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
- 2-[[[(2, 3, 4-trimethoxyphenyl)-3-hydroxyphosphinyl]amino]-2-

- phenylethanoic acid;
- 2-[[[(1-naphthyl)hydroxyphosphinyl]amino]-2-phenylethanoic acid;
- 2-[[[(2-naphthyl)hydroxyphosphinyl]amino]-2-phenylethanoic acid;
- 5 2-[[[(1-naphthyl)methylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
- 2-[[[(2-naphthyl)methylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
- 10 2-[[[(1-naphthyl)ethylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
- 2-[[[(2-naphthyl)ethylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
- 2-[[[(1-naphthyl)propylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
- 15 2-[[[(2-naphthyl)propylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
- 2-[[[(1-naphthyl)butylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
- 20 2-[[[(2-naphthyl)butylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
- 2-[[phenylprop-2-enylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-methylethanoic acid;
- 25 2-[[benzylhydroxyphosphinyl]amino]-2-ethylethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-propylethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-butylethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-cyclohexylethanoic acid;

- 2-[[benzylhydroxyphosphinyl]amino]-2-(cyclohexyl)methyl
ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-benzylethanoic acid;
- 5 2-[[benzylhydroxyphosphinyl]amino]-2-phenylethylethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-phenylpropylethanoic
acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-phenylbutylethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-(2, 3, 4-
10 trimethoxyphenyl)ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-(1-naphthyl)ethanoic
acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-(2-naphthyl)ethanoic
acid;
- 15 2-[[benzylhydroxyphosphinyl]amino]-2-(1-naphthyl)methyl
ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-(2-naphthyl)methyl
ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-(1-naphthyl)ethylethanoic
20 acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-(2-naphthyl)ethylethanoic
acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-(1-naphthyl)propyl
ethanoic acid;
- 25 2-[[benzylhydroxyphosphinyl]amino]-2-(2-naphthyl)propyl
ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-(1-naphthyl)butylethanoic
acid;

- 2-[[benzylhydroxyphosphinyl]amino]-2-(2-naphthyl)butylethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-phenolprop-2-enylethanoic acid;
- 5 2-[(methylhydroxyphosphinyl)amino]hexanedioic acid;
- 2-[(benzylhydroxyphosphinyl)amino]hexanedioic acid;
- 2-[(methylhydroxyphosphinyl)amino]heptanedioic acid;
- 2-[(benzylhydroxyphosphinyl)amino]heptanedioic acid;
- 2-[(methylhydroxyphosphinyl)amino]octanedioic acid;
- 10 2-[(benzylhydroxyphosphinyl)amino]octanedioic acid;
- 2-[(methylhydroxyphosphinyl)amino]nonanedioic acid;
- 2-[(benzylhydroxyphosphinyl)amino]nonanedioic acid;
- 2-[(methylhydroxyphosphinyl)amino]decanedioic acid;
- 2-[(benzylhydroxyphosphinyl)amino]decanedioic acid;
- 15 3-[[[(2-pyridyl)methylhydroxyphosphinyl]amino]pentanedioic acid;
- 3-[[[(3-pyridyl)methylhydroxyphosphinyl]amino]pentanedioic acid;
- 3-[[[(4-pyridyl)methylhydroxyphosphinyl]amino]pentanedioic acid;
- 20 3-[[[(3-pyridyl)ethylhydroxyphosphinyl]amino]pentanedioic acid;
- 3-[[[(3-pyridyl)propylhydroxyphosphinyl]amino]pentanedioic acid;
- 3-[[[(tetrahydrofuranyl)methylhydroxyphosphinyl]amino]pentanedioic acid;
- 25 3-[[[(tetrahydrofuranyl)ethylhydroxyphosphinyl]amino]pentanedioic acid;
- 3-[[[(tetrahydrofuranyl)propylhydroxyphosphinyl]amino]

- pentanedioic acid;
- 3-[[(2-indolyl) methylhydroxyphosphinyl] amino] pentanedioic acid;
- 3-[[(3-indolyl) methylhydroxyphosphinyl] amino] pentanedioic acid;
- 5 3-[[(4-indolyl) methylhydroxyphosphinyl] amino] pentanedioic acid;
- 3-[[(3-indolyl) ethylhydroxyphosphinyl] amino] pentanedioic acid;
- 3-[[(3-indolyl) propylhydroxyphosphinyl] amino] pentanedioic acid;
- 10 3-[[(2-thienyl) methylhydroxyphosphinyl] amino] pentanedioic acid;
- 3-[[(3-thienyl) methylhydroxyphosphinyl] amino] pentanedioic acid;
- 15 3-[[(4-thienyl) methylhydroxyphosphinyl] amino] pentanedioic acid;
- 3-[[(3-thienyl) ethylhydroxyphosphinyl] amino] pentanedioic acid;
- 3-[[(3-thienyl) propylhydroxyphosphinyl] amino] pentanedioic acid;
- 20 2-[[(2-pyridyl) methylhydroxyphosphinyl] amino] -2-phenylethanoic acid;
- 2-[[(3-pyridyl) methylhydroxyphosphinyl] amino] -2-phenylethanoic acid;
- 2-[[(4-pyridyl) methylhydroxyphosphinyl] amino] -2-phenylethanoic acid;
- 25 2-[[(3-pyridyl) ethylhydroxyphosphinyl] amino] -2-phenylethanoic acid;
- 2-[[(3-pyridyl) propylhydroxyphosphinyl] amino] -2-phenylethanoic

- acid;
- 2-[[[(tetrahydrofuranyl)methylhydroxyphosphinyl]amino]-2-phenyl
ethanoic acid;
- 2-[[[(tetrahydrofuranyl)ethylhydroxyphosphinyl]amino]-2-phenyl
5 ethanoic acid;
- 2-[[[(tetrahydrofuranyl)propylhydroxyphosphinyl]amino]-2-phenyl
ethanoic acid;
- 2-[[[(2-indolyl)methylhydroxyphosphinyl]amino]-2-phenylethanoic
acid;
- 10 2-[[[(3-indolyl)methylhydroxyphosphinyl]amino]-2-phenylethanoic
acid;
- 2-[[[(4-indolyl)methylhydroxyphosphinyl]amino]-2-phenylethanoic
acid;
- 2-[[[(3-indolyl)ethylhydroxyphosphinyl]amino]-2-phenylethanoic
15 acid;
- 2-[[[(3-indolyl)propylhydroxyphosphinyl]amino]-2-phenylethanoic
acid;
- 2-[[[(2-thienyl)methylhydroxyphosphinyl]amino]-2-phenylethanoic
acid;
- 20 2-[[[(3-thienyl)methylhydroxyphosphinyl]amino]-2-phenylethanoic
acid;
- 2-[[[(4-thienyl)methylhydroxyphosphinyl]amino]-2-phenylethanoic
acid;
- 2-[[[(3-thienyl)ethylhydroxyphosphinyl]amino]-2-phenylethanoic
25 acid;
- 2-[[[(3-thienyl)propylhydroxyphosphinyl]amino]-2-phenylethanoic
acid;
- 2-[[[benzylhydroxyphosphinyl]amino]-2-(2-pyridyl)methylethanoic

acid;

2-[[benzylhydroxyphosphinyl]amino]-2-(3-pyridyl)methylethanoic
acid;

5 2-[[benzylhydroxyphosphinyl]amino]-2-(4-pyridyl)methylethanoic
acid;

2-[[benzylhydroxyphosphinyl]amino]-2-(3-pyridyl)ethylethanoic
acid;

2-[[benzylhydroxyphosphinyl]amino]-2-(3-pyridyl)propylethanoic
acid;

10 2-[[benzylhydroxyphosphinyl]amino]-2-(tetrahydrofuran-2-yl)
methylethanoic acid;

2-[[benzylhydroxyphosphinyl]amino]-2-(tetrahydrofuran-2-yl)
ethylethanoic acid;

15 2-[[benzylhydroxyphosphinyl]amino]-2-(tetrahydrofuran-2-yl)
propylethanoic acid;

2-[[benzylhydroxyphosphinyl]amino]-2-(2-indolyl)methylethanoic
acid;

2-[[benzylhydroxyphosphinyl]amino]-2-(3-indolyl)methylethanoic
acid;

20 2-[[benzylhydroxyphosphinyl]amino]-2-(4-indolyl)methylethanoic
acid;

2-[[benzylhydroxyphosphinyl]amino]-2-(3-indolyl)ethylethanoic
acid;

25 2-[[benzylhydroxyphosphinyl]amino]-2-(3-indolyl)propylethanoic
acid;

2-[[benzylhydroxyphosphinyl]amino]-2-(2-thienyl)methylethanoic
acid;

2-[[benzylhydroxyphosphinyl]amino]-2-(3-thienyl)methylethanoic

acid;

2-[[benzylhydroxyphosphinyl]amino]-2-(4-thienyl)methylethanoic acid;

2-[[benzylhydroxyphosphinyl]amino]-2-(3-thienyl)ethylethanoic acid;

2-[[benzylhydroxyphosphinyl]amino]-2-(3-thienyl)propylethanoic acid; and a pharmaceutically acceptable salt, hydrate, or a mixture thereof.

68. The method of claim 66 wherein the NAALADase inhibitor is selected from the group consisting of:

2-[(benzylhydroxyphosphinyl)amino]pentanedioic acid;

2-[(phenylhydroxyphosphinyl)amino]pentanedioic acid;

2-[[[(hydroxy)phenylmethyl]hydroxyphosphinyl]amino]pentanedioic acid;

2-[(butylhydroxyphosphinyl)amino]pentanedioic acid;

2-[[[(3-methylbenzyl)hydroxyphosphinyl]amino]pentanedioic acid;

2-[(3-phenylpropylhydroxyphosphinyl)amino]pentanedioic acid;

2-[[[(4-fluorophenyl)hydroxyphosphinyl]amino]pentanedioic acid;

2-[(methylhydroxyphosphinyl)amino]pentanedioic acid;

2-[(phenylethylhydroxyphosphinyl)amino]pentanedioic acid;

2-[[[(4-methylbenzyl)hydroxyphosphinyl]amino]pentanedioic acid;

2-[[[(4-fluorobenzyl)hydroxyphosphinyl]amino]pentanedioic acid;

2-[[[(4-methoxybenzyl)hydroxyphosphinyl]amino]pentanedioic acid;

2-[[[(2-fluorobenzyl)hydroxyphosphinyl]amino]pentanedioic acid;

2-[[[(pentafluorobenzyl)hydroxyphosphinyl]amino]pentanedioic acid;

2-[(phosphono)amino]pentanedioic acid;

2-[[[(3-trifluoromethylbenzyl)hydroxyphosphinyl]amino]pentanedioic acid; and a pharmaceutically acceptable salt, hydrate, or a mixture thereof.

5 69. The method of claim 59 wherein R_1 or R_2 is heterocyclic and X is CH_2 .

70. The method of claim 69 wherein the NAALADase inhibitor is selected from the group consisting of:

- 2-[[[(2-pyridyl)hydroxyphosphinyl]methyl]pentanedioic acid;
2-[[[(3-pyridyl)hydroxyphosphinyl]methyl]pentanedioic acid;
10 2-[[[(4-pyridyl)hydroxyphosphinyl]methyl]pentanedioic acid;
2-[[[(tetrahydrofuran-2-yl)hydroxyphosphinyl]methyl]pentanedioic acid;
2-[[[(2-indolyl)hydroxyphosphinyl]methyl]pentanedioic acid;
2-[[[(3-indolyl)hydroxyphosphinyl]methyl]pentanedioic acid;
15 2-[[[(4-indolyl)hydroxyphosphinyl]methyl]pentanedioic acid;
2-[[[(2-thienyl)hydroxyphosphinyl]methyl]pentanedioic acid;
2-[[[(3-thienyl)hydroxyphosphinyl]methyl]pentanedioic acid;
2-[[[(4-thienyl)hydroxyphosphinyl]methyl]pentanedioic acid;
3-[(2-pyridyl)hydroxyphosphinyl]-2-phenylpropanoic acid;
20 3-[(3-pyridyl)hydroxyphosphinyl]-2-phenylpropanoic acid;
3-[(4-pyridyl)hydroxyphosphinyl]-2-phenylpropanoic acid;
3-[(tetrahydrofuran-2-yl)hydroxyphosphinyl]-2-phenylpropanoic acid;
3-[(2-indolyl)hydroxyphosphinyl]-2-phenylpropanoic acid;
25 3-[(3-indolyl)hydroxyphosphinyl]-2-phenylpropanoic acid;
3-[(4-indolyl)hydroxyphosphinyl]-2-phenylpropanoic acid;
3-[(2-thienyl)hydroxyphosphinyl]-2-phenylpropanoic acid;
3-[(3-thienyl)hydroxyphosphinyl]-2-phenylpropanoic acid;

-221-

3-[(4-thienyl)hydroxyphosphinyl]-2-phenylpropanoic acid;
3-(benzylhydroxyphosphinyl)-2-(2-pyridyl)propanoic acid;
3-(benzylhydroxyphosphinyl)-2-(3-pyridyl)propanoic acid;
3-(benzylhydroxyphosphinyl)-2-(4-pyridyl)propanoic acid;
5 3-(benzylhydroxyphosphinyl)-2-(tetrahydrofuranyl)propanoic acid;
3-(benzylhydroxyphosphinyl)-2-(2-indolyl)propanoic acid;
3-(benzylhydroxyphosphinyl)-2-(3-indolyl)propanoic acid;
3-(benzylhydroxyphosphinyl)-2-(4-indolyl)propanoic acid;
10 3-(benzylhydroxyphosphinyl)-2-(2-thienyl)propanoic acid;
3-(benzylhydroxyphosphinyl)-2-(3-thienyl)propanoic acid;
3-(benzylhydroxyphosphinyl)-2-(4-thienyl)propanoic acid; and
a pharmaceutically acceptable salt, hydrate, or a mixture thereof.

15 71. The method of claim 69 wherein R_1 or R_2 is heterocyclic and X is oxygen.

72. The method of claim 71 wherein the NAALADase inhibitor is selected from the group consisting of:
2-[[[(2-pyridyl)hydroxyphosphinyl]oxy]pentanedioic acid;
20 2-[[[(3-pyridyl)hydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(4-pyridyl)hydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(tetrahydrofuranyl)hydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(2-indolyl)hydroxyphosphinyl]oxy]pentanedioic acid;
25 2-[[[(3-indolyl)hydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(4-indolyl)hydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(2-thienyl)hydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(3-thienyl)hydroxyphosphinyl]oxy]pentanedioic acid;

2-[[(4-thienyl) hydroxyphosphinyl] oxy] pentanedioic acid;
2-[[(2-pyridyl) hydroxyphosphinyl] oxy] -2-phenylethanoic acid;
2-[[(3-pyridyl) hydroxyphosphinyl] oxy] -2-phenylethanoic acid;
2-[[(4-pyridyl) hydroxyphosphinyl] oxy] -2-phenylethanoic acid;
5 2-[[(tetrahydrofuranyl) hydroxyphosphinyl] oxy] -2-phenylethanoic
acid;
2-[[(2-indolyl) hydroxyphosphinyl] oxy] -2-phenylethanoic acid;
2-[[(3-indolyl) hydroxyphosphinyl] oxy] -2-phenylethanoic acid;
2-[[(4-indolyl) hydroxyphosphinyl] oxy] -2-phenylethanoic acid;
10 2-[[(2-thienyl) hydroxyphosphinyl] oxy] -2-phenylethanoic acid;
2-[[(3-thienyl) hydroxyphosphinyl] oxy] -2-phenylethanoic acid;
2-[[(4-thienyl) hydroxyphosphinyl] oxy] -2-phenylethanoic acid;
2-[[benzylhydroxyphosphinyl] oxy] -2-(2-pyridyl) ethanoic acid;
2-[[benzylhydroxyphosphinyl] oxy] -2-(3-pyridyl) ethanoic acid;
15 2-[[benzylhydroxyphosphinyl] oxy] -2-(4-pyridyl) ethanoic acid;
2-[[benzylhydroxyphosphinyl] oxy] -2-(tetrahydrofuranyl) ethanoic
acid;
2-[[benzylhydroxyphosphinyl] oxy] -2-(2-indolyl) ethanoic acid;
2-[[benzylhydroxyphosphinyl] oxy] -2-(3-indolyl) ethanoic acid;
20 2-[[benzylhydroxyphosphinyl] oxy] -2-(4-indolyl) ethanoic acid;
2-[[benzylhydroxyphosphinyl] oxy] -2-(2-thienyl) ethanoic acid;
2-[[benzylhydroxyphosphinyl] oxy] -2-(3-thienyl) ethanoic acid;
2-[[benzylhydroxyphosphinyl] oxy] -2-(4-thienyl) ethanoic acid;
and a pharmaceutically acceptable salt, hydrate, or a mixture
25 thereof.

73. The method of claim 59 wherein R_1 or R_2 is heterocyclic and X is NR_1 .

74. The method of claim 73 wherein the NAALADase


inhibitor is a compound selected from the group consisting of:

- 2-[[[(2-pyridyl)hydroxyphosphinyl]amino]pentanedioic acid;
2-[[[(3-pyridyl)hydroxyphosphinyl]amino]pentanedioic acid;
2-[[[(4-pyridyl)hydroxyphosphinyl]amino]pentanedioic acid;
5 2-[[[(tetrahydrofuranyl)hydroxyphosphinyl]amino]pentanedioic acid;
2-[[[(2-indolyl)hydroxyphosphinyl]amino]pentanedioic acid;
2-[[[(3-indolyl)hydroxyphosphinyl]amino]pentanedioic acid;
2-[[[(4-indolyl)hydroxyphosphinyl]amino]pentanedioic acid;
10 2-[[[(2-thienyl)hydroxyphosphinyl]amino]pentanedioic acid;
2-[[[(3-thienyl)hydroxyphosphinyl]amino]pentanedioic acid;
2-[[[(4-thienyl)hydroxyphosphinyl]amino]pentanedioic acid;
2-[[[(2-pyridyl)hydroxyphosphinyl]amino]-2-phenylethanoic acid;
2-[[[(3-pyridyl)hydroxyphosphinyl]amino]-2-phenylethanoic acid;
15 2-[[[(4-pyridyl)hydroxyphosphinyl]amino]-2-phenylethanoic acid;
2-[[[(tetrahydrofuranyl)hydroxyphosphinyl]amino]-2-phenyl ethanoic acid;
2-[[[(2-indolyl)hydroxyphosphinyl]amino]-2-phenylethanoic acid;
2-[[[(3-indolyl)hydroxyphosphinyl]amino]-2-phenylethanoic acid;
20 2-[[[(4-indolyl)hydroxyphosphinyl]amino]-2-phenylethanoic acid;
2-[[[(2-thienyl)hydroxyphosphinyl]amino]-2-phenylethanoic acid;
2-[[[(3-thienyl)hydroxyphosphinyl]amino]-2-phenylethanoic acid;
2-[[[(4-thienyl)hydroxyphosphinyl]amino]-2-phenylethanoic acid;
2-[[benzylhydroxyphosphinyl]amino]-2-(2-pyridyl)ethanoic acid;
25 2-[[benzylhydroxyphosphinyl]amino]-2-(3-pyridyl)ethanoic acid;
2-[[benzylhydroxyphosphinyl]amino]-2-(4-pyridyl)ethanoic acid;
2-[[benzylhydroxyphosphinyl]amino]-2-(tetrahydrofuranyl) ethanoic acid;

2-[[benzylhydroxyphosphinyl]amino]-2-(2-indolyl)ethanoic acid;
2-[[benzylhydroxyphosphinyl]amino]-2-(3-indolyl)ethanoic acid;
2-[[benzylhydroxyphosphinyl]amino]-2-(4-indolyl)ethanoic acid;
2-[[benzylhydroxyphosphinyl]amino]-2-(2-thienyl)ethanoic acid;
5 2-[[benzylhydroxyphosphinyl]amino]-2-(3-thienyl)ethanoic acid;
2-[[benzylhydroxyphosphinyl]amino]-2-(4-thienyl)ethanoic acid;
and a pharmaceutically acceptable salt, hydrate, or a mixture
thereof.


1/6

FIG. 1




2/6

FIG.2



3/6


FIG.3




SUBSTITUTE SHEET (RULE 26)

4/6


FIG.4



5/6



6/6



SUBSTITUTE SHEET (RULE 26)

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US97/10149

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : Please See Extra Sheet.

US CL : Please See Extra Sheet.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/7, 89, 94, 95, 99, 120; 546/344; 548/144, 469; 549/79, 229; 562/24

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

CAS ONLINE, APS

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X — A	JACKSON et al. Design, synthesis, and biological activity of a potent inhibitor of the Neuropeptidase N-acetylated α -linked acidic dipeptidase. J. Med. Chem. 1996, Vol. 39, pages 619-622, especially see pages 620-621 and Table 1 on page 620.	52, 53, 56-62 ----- 63-71
X	STAUCH et al. The effects of N-acetylated alpha-linked acidic dipeptidase (NAALADase) inhibitors on [3 H] NAAG catabolism in vivo. Neuroscience Letters. 1989, Vol. 100, pages 295-300, especially see Table 1 on page 297.	55

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	T	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	X*	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
B earlier document published on or after the international filing date	Y*	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	g*	document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means		
P document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search


13 AUGUST 1997

Date of mailing of the international search report

24 SEP 1997

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

CHANA AULAKH

Telephone No. (703) 308-1235

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US97/10149

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	CARTER et al. Prostate-specific membrane antigen is a hydrolase with substrate and pharmacologic characteristics of a neuropeptidase. Proc. Natl. Acad. Sci. 1996, Vol. 93, pages 749-753, 1996, see the entire document.	1-51
A	SUBASINGHE et al. Synthesis of Acyclic and dehydroaspartic acid analogues of Ac-Asp-Glu-OH and their inhibition of rat brain N-acetylated α -linked acidic dipeptidase (NAALA Dipeptidase). J. Med. Chem. 1990, Vol. 33, pages 2734-2744, see the entire document.	54

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US97/10149

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

Please See Extra Sheet.

1. ☒ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US97/10149

A. CLASSIFICATION OF SUBJECT MATTER:
IPC (6):

A61K 38/05, 31/675, 31/67, 31/66, 31/665; C07D 209/04, 213/16, 271/07, 307/36, 33/10; C07F 9/28

A. CLASSIFICATION OF SUBJECT MATTER:
US CL :

514/7, 89, 94, 95, 99, 120; 546/344; 548/144, 469; 549/79, 229; 562/24

BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

This application contains claims directed to more than one species of the generic invention. These species are deemed to lack Unity of Invention because they are not so linked as to form a single inventive concept under PCT Rule 13.1. The species are as follows:

- I. A method of treating cancer wherein the NAALADase inhibitor is an acidic peptide analog, classifiable in class 514, subclass 7.
- II. A method of treating cancer wherein the NAALADase inhibitor is a glutamate mimic, classifiable in class 514, subclass 94.
- III. A method of treating cancer wherein the NAALADase inhibitor is a glutamate-derived hydroxyphosphinyl derivative and R is pyridyl, classifiable in class 514, subclass 89.
- IV. A method treating cancer wherein the NAALADase inhibitor is a glutamate-derived hydroxyphosphinyl derivative and R is indolyl, classifiable in class 514, subclass 95.
- V. A method of treating cancer wherein the NAALADase inhibitor is a glutamate-derived hydroxyphosphinyl derivative and R is furyl or thienyl, classifiable in class 514, subclass 99.
- VI. A method of treating cancer wherein the NAALADase inhibitor is a glutamate-derived hydroxyphosphinyl derivative and R is substituted or unsubstituted alkyl or alkenyl or cycloalkyl or cycloalkenyl or phenyl, classifiable in class 514, subclass 120.

The claims are deemed to correspond to the species listed above in the following manner:

Species I. Claims 3, 29, 54.

Species II. Claims 4, 30, 55.

Species III. 5, 10-25, 31, 36-51, 56 and 59-74.

Species IV. Claims 5, 10-25, 31, 36-51, 56 and 59-74.

Species V. Claims 5, 10-25, 31, 36-51, 56 and 59-74.

Species VI. Claims 5, 10-25, 31, 36-51, 56 and 59-74.

The following claims are generic: Claims 1, 2, 6-9, 26-28, 32-35, 52, 53, 57 and 58.

The species listed above do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, the species lack the same or corresponding special technical features for the following reasons: There is no common core which in the Markush Practice, is a significant structural element shared by all the alternatives; see PCT Administrative Instructions Annex B Part I (f) (i) (B) (1) and further, all alternatives do not belong to a recognized class of chemical compounds in the art to which the invention pertains; see *supra* (B) (2).

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☒ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.